

DAFTAR PUSTAKA

1. Vaughan D. G, Asbury, T., Riordan-eva , P. oftalmologi umum edisi 17, EGC, Jakarta. 2015;169.
2. M.W. Ansari, A. Nadeem, Anatomy of the orbit, in: Atlas of Ocular Anatomy, Springer, 2016: 1–10
3. Azhar Seedat, Shaheen Seedat, Sulaiman E. I. Moosa. Proptosis with Increased Orbital Fat in an Obese Patient. Case Reports in Radiology. 2020, 1-3.
4. Kasen R. Hutchings, Seth J. Fritzhand, Bitu Esmaeli, Kirthi Koka, Jiawei Zhao, Salmaan Ahmed, James Matthew Debnam. Graves' Eye Disease: Clinical and Radiological Diagnosis. Biomedicines. 2023;11(2)312
5. Smith, T.J.; Hegedüs, L. Graves' Disease. N. Engl. J. Med. 2016; 375: 1552–65.
6. Bao Y, Zhang Z, Li C, Ma H, Yin P, Wang Y, Luo G, Lu R. Geometric and Volumetric Measurements of Orbital Structures in CT Scans in Thyroid Eye Disease Classification. *Applied Sciences*. 2021; 11(11):4873. <https://doi.org/10.3390/app11114873>
7. Du Y, Lu BY, Chen J, He JF. Measurement of the orbital soft tissue volume in Chinese adults based on three-dimensional CT reconstruction. *J Ophthalmol*. 2019; 9721085.
8. Khizar Rana, Valerie Juniat, Wen Yong, Sandy Patel, Dinesh Selva. Normative orbital measurements in an Australian cohort on computed tomography. *Orbit*. 2023;42:1, 68
9. Shayan Soomro, Kimia Ziahosseini, Poonam Sharma. Proptosis of the eye: a potential form of Madelung's disease. *BMJ Case Reports*. 2021;14:(9), 243669
10. Yoko Kaichi, Keizo Tanitame, Hiroaki Terada, Hideki Itakura, Haruya Ohno, Masayasu Yoneda, Yuji Takahashi, Yuji Akiyama, Kazuo Awai. Thyroid-associated Orbitopathy: Quantitative Evaluation of the Orbital Fat Volume and Edema Using IDEALFSE. *European Journal of Radiology Open*. 2019;6:182-186.
11. Qingyao Ning, Xiaoyao Yu, Qi Gao, Jiajun Xie, Chunlei Yao, Kun Zhou, Juan Ye. An accurate interactive segmentation and volume calculation of orbital soft tissue for orbital reconstruction after enucleation. *BMC Ophthalmology*. 2019;19:1

12. Byun, J.S.; Moon, N.J.; Lee, J.K. Quantitative analysis of orbital soft tissues on computed tomography to assess the activity of thyroid-associated orbitopathy. *Graefe's Arch. Clin. Exp. Ophthalmol.* 2017, 255, 413–420
13. J. Singh, R. Ab Rahman, Z.A. Rajion, J. Abdullah, I. Mohamad, Orbital morphometry: a computed tomography analysis, *J. Craniofac. Surg.* 2017; 28 (1)e64–e70
14. Weaver AA, Loftis K.L, Tan J,C et al. CT Based Three-Dimensions measurement of Orbit and Eye Anthropometrty(editor Weaver AA).vol.15. 2024. 4892-97
15. Regensburg NI, Kok PH, Zonneveld FW, et al. A New and Validated CT-based method for the calculation of orbital soft tissue volumes (Editor : Regensburg NI).2015;1758-62
- 16.1Regensburg NI, Wiersinga WM, van Velthoven ME, Berendschot TT, Zonneveld FW, Baldeschi L, et al: Age and gender-specific reference values of orbital fat and muscle volumes in Caucasians. *Br J Ophthalmol* 2011;95:1660-1663
17. Kim IT, Choi BJ. Normal range of exophthalmos values on orbit computerized tomography in Koreans. *Ophthalmologica.* 2001;[215\(3\)](#):156–162. doi:10.1159/000050850
18. Bagheri A, Shahraki K, Arabi A, Kashkouli MB.J Normal Exophthalmometry Values in Iranian Population: A Meta-analysis. *Ophthalmic Vis Res.* 2021 Jul 29;16(3):470-477. doi: 10.18502/jovr.v16i3.9441. eCollection 2021 Jul-Sep
19. Sibagariang R. Volume Lemak Retro-Orbita Normal Pada Orang Dewasa Berdasarkan Pengukuran CT Scan. Jakarta 2011.
20. Andrades P, Cuevas P, Hernandez R, Danilla b S, Villalobos R. Characterization of the orbital volume in normal population.*J of Cranio-Maxillo-Facial Surgery.*2018.46:597.