

(RESEARCH ARTICLE)



# The relationship between the use of cosmetics and the occurrence of acne vulgaris in female medical students at the faculty of medicine UKI in the year 2016

Ago Harlim <sup>1,\*</sup>, Daniar Hapsari <sup>2</sup> and Charity Harlim <sup>3</sup>

<sup>1</sup> *Dermatology and Venereology Division, Medical Faculty, Christian University of Indonesia, Indonesia.*

<sup>2</sup> *Medical Faculty, Christian University of Indonesia, Indonesia.*

<sup>3</sup> *Medical Faculty, Tarumanegara University, Indonesia.*

Magna Scientia Advanced Research and Reviews, 2024, 10(01), 086–092

Publication history: Received on 16 November 2023; revised on 02 January 2024; accepted on 05 January 2024

Article DOI: <https://doi.org/10.30574/msarr.2024.10.1.0176>

## Abstract

Acne vulgaris is a chronic inflammation of the pilosebaceous follicles caused by various factors, typically characterized by distinctive clinical features and usually occurring in adolescents and young adults. The use of cosmetics can lead to the development of acne in women, as the ingredients used may be comedogenic or acneogenic. The current use of cosmetics is undeniably an essential need for women. The aim of this study is to determine the relationship between the use of different types of cosmetics and the occurrence of AV. The variables examined in this research include compact powder, loose powder, sunblock, foundation, and blush. This study is an observational research with a cross-sectional design involving a sample of 100 female medical students from the Faculty of Medicine at the Christian University of Indonesia, divided into two respondent groups: one with acne and one without acne. The chi-square test is used for the analysis. From the research results, it can be concluded that, of the five variables studied, only the use of compact powder showed a significant correlation.

**Keywords:** Acne vulgaris; Compact powder; Sunblock; Foundation; Blush

## 1. Introduction

The current use of cosmetics is an essential need for women; however, cosmetics can lead to skin diseases such as Acne Vulgaris (AV). AV is commonly found in adolescents and young adults<sup>1-3</sup>. The occurrence of AV is also influenced by other factors such as the content of cosmetics, skin type (sensitive skin), or the frequency of use<sup>4</sup>. There are four pathogenesises of acne, namely increased sebum production, hyperproliferation of pilosebaceous follicles, colonization of *Propionibacterium acnes* (PA), and the inflammatory process<sup>1-5</sup>. Clinical symptoms of acne include comedones, papules, pustules, nodules, and cysts<sup>6</sup>. Given the widespread use of cosmetics and the prevalence of AV, the purpose of this study is to prove the relationship between the use of different types of cosmetics (loose powder, compact powder, sunblock, foundation, blush) and the occurrence of AV.

## 2. Material and methods

The research design used is observational with a cross-sectional design. Inclusion criteria are female students willing to participate as research respondents, aged between 18-23 years, and have used cosmetics for at least 2 weeks. Exclusion criteria are female students undergoing acne treatment by a general practitioner, dermatologist, or purchasing acne medication from a store/pharmacy in the last 2 weeks. The minimum sample size used is 50 samples. Data is collected through questionnaire completion, followed by a physical examination according to the clinical presentation of acne vulgaris. Acne vulgaris is categorized as moderate (acne <20) and severe (acne >20), and types of

\* Corresponding author: Ago Harlim

cosmetics include compact powder, loose powder, sunblock, foundation, and blush. Data processing is done using SPSS software for univariate and bivariate analysis using chi-square. Research ethics involve obtaining informed consent, prioritizing the interests of respondents, and not imposing any charges.

### 3. Result

Univariate analysis describes the characteristics of 100 respondents, including age, class, acne issues, and the use of loose powder, compact powder, sunblock, foundation, and blush among the 100 respondents.

**Tabel 1** Distribution of Respondent Characteristics

		Frequency	Percentage
Age	18 years	18	18.00
	19 years	11	11.00
	20 years	13	13.00
	21 years	49	49.00
	22 years	9	9.00
	23 years	0	0.00
Class	2013	69	69.00
	2014	2	2.00
	2015	3	3.00
	2016	26	26.00
Experiencing Acne Issues	Not experiencing	50	50.00
	Moderate acne (< 20 acne)	48	48.00
	Severe acne(> 20 acne)	2	2.00
Use of loose powder	Never	36	36.00
	Once a day	48	48.00
	Twice a day	11	11.00
	Three times a day	5	5.00
Use of Sunblock	Never	48	48.00
	Once a day	46	46.00
	> More than Once a day	6	6.00
Use of foundation	Never	66	66.00
	Once a day	28	28.00
	Twice a day	6	6.00
	> More than twice a day	0	0.00
<i>Use of Blush on</i>	Never	73	73.00
	Once a day	25	25.00
	> More than Once a day	2	2.00
Total		100	100

Bivariate analysis was conducted to examine the relationship between the use of cosmetics, including compact powder, loose powder, sunblock, foundation, and blush on, with acne vulgaris using cross-tabulation and the chi-square ( $\chi^2$ ) method.

**Table 2** Cross-tabulation between the Use of Compact Powder and the Occurrence of Acne Vulgaris

		Acne Vulgaris						
		No Acne		Moderate acne		Severe acne		Total
		n	%	n	%	n	%	
Compact Powder	Never	25	25	21	21	0	0	46
	1x a day	14	14	16	16	0	0	30
	2x a day	7	7	5	5	1	1	13
	3x a day	4	4	6	6	0	0	10
	4x a day	0	0	0	0	1	1	1
Total		50	50	48	48	2	2	100

**Table 3** Chi-Square Tests

Value	Asymp. Sig. (2-sided)
N	100
df	8
Pearson Chi-Square	54.082
Critical Value	15.507
Conclusion	Significant

The result of the analysis of compact powder yielded a  $\chi^2$  value of 54.082, which is greater than the  $\chi^2$  table value. Additionally, the p-value is less than 0.05, leading to the conclusion that there is a significant relationship between the use of compact powder and the occurrence of acne vulgaris.

**Table 4** Cross-tabulation between the Use of Loose Powder and the Occurrence of Acne Vulgaris

		Acne Vulgaris						
		No acne		Moderate acne		Severe acne		Total
		n	%	n	%	n	%	
Loose Powder	Never	22	22	14	14	0	0	36
	1x a day	19	19	27	27	2	2	48
	2x a day	6	6	5	5	0	0	11
	3x a day	3	3	2	2	0	0	5
Total		50	50	48	48	2	2	100

**Table 5** Chi-Square Tests

Value		Asymp. Sig. (2-sided)
N	100	0.466
df	6	
Pearson Chi-Square	5.627	
Critical Value	12.592	
Conclusion		Not Significant

After the analysis, the calculated  $\chi^2$  value is (5.627), which is less than the  $\chi^2$  table value, and the p-value is 0.466, where the p-value is greater than 0.05. Therefore, it can be concluded that there is no significant relationship between the use of loose powder and the occurrence of acne vulgaris.

**Table 6** Cross-tabulation between the Use of Sunblock and the Occurrence of Acne Vulgaris

		Acne Vulgaris						Total
		No Acne		Moderate acne		Severe acne		
		n	%	n	%	n	%	
Sun block	Never	27	27	21	21	0	0	48
	1x a day	20	20	24	24	2	2	46
	>1x a day	3	3	3	3	0	0	6
Total		50	50	48	48	2	2	100

**Table 7** Chi-Square Tests

Value		Asymp. Sig. (2-sided)
N	100	0.483
df	4	
Pearson Chi-Square	3.467	
Critical Value	9.488	
Conclusion		Not Significant

The analysis resulted in a calculated  $\chi^2$  value of 3.467, as  $\chi^2$  hitung <  $\chi^2$  tabel, and the obtained p-value is 0.483. When the p-value is greater than 0.05, it can be concluded that there is no significant relationship between the use of foundation and the occurrence of acne vulgaris.

**Table 8** Cross-tabulation between the Use of Foundation and the Occurrence of Acne Vulgaris

		Acne Vulgaris						Total
		No Acne		Moderate acne		Severe acne		
		n	%	n	%	n	%	
Foundation	Never	34	34	32	32	0	0	66
	1x a day	12	12	14	14	2	2	28
	2x a day	4	4	2	2	0	0	6
Total		50	50	48	48	2	2	100

**Table 9** Chi-Square Tests

Value		Asymp. Sig. (2-sided)
N	100	0.193
df	4	
Pearson Chi-Square	6.088	
Critical Value	9.488	
Conclusion		Not Significant

The analysis of the use of foundation resulted in a  $\chi^2$  table value of 9.488. With a calculated  $\chi^2$  value of 6.088, where  $\chi^2_{hitung} < \chi^2_{tabel}$ , and a p-value of 0.193, it can be concluded that there is no significant relationship between the use of foundation and the occurrence of acne vulgaris, as the p-value is greater than 0.05.

**Table 10** Cross-tabulation between the Use of Blush On and the Occurrence of Acne Vulgaris

		Akne Vulgaris						
		No Acne		Moderate acne		Severe acne		Total
		n	%	n	%	n	%	
Blush On	Never	40	40	33	33	0	0	73
	1x a day	10	10	13	13	2	2	25
	>1x a day	0	0	2	2	0	0	2
Total		50	50	48	48	2	2	100

**Table 11** Chi-Square Tests

Value		Asymp. Sig. (2-sided)
N	100	0.057
df	4	
Pearson Chi-Square	9.164	
Critical Value	9.488	
Conclusion		Not Significant

The analysis of the use of compact powder yielded a  $\chi^2$  value of 6.088, with a calculated  $\chi^2$  value lower than the  $\chi^2$  table value (9.488), and a p-value of 0.193. With a p-value greater than 0.05, it can be concluded that there is no significant relationship between the use of foundation and the occurrence of acne vulgaris.

#### 4. Discussion

In the pathogenesis of acne vulgaris, increased sebum production and enlarged sebaceous follicle size occur. Sebum production is regulated by androgen hormones, which also play a role in changes to sebocyte and keratinocyte cells, leading to the development of microcomedones and comedones that progress into inflammatory lesions. Individuals with acne also experience hyperplasia of pilosebaceous follicles, possibly due to decreased linoleic acid in acne patients<sup>7-9</sup>. This leads to thickening of the granulosum layer, increased keratohyalin, increased lipid content, resulting in gradual thickening and blockage<sup>7</sup>. These blockages are also influenced by the use of compact powder because the particle size of compact powder is extremely small, allowing it to enter the pores. Additionally, the occurrence of blockages is also influenced by the presence of lanolin as a binding agent that is acne-prone<sup>10</sup>. Furthermore, the tropical climate in

Indonesia, situated in a tropical region, causes the skin to tend to sweat and become oily. This can lead to clogged pores, especially when combined with the use of compact powder<sup>11,12</sup>.

Loose powder rarely causes acne because the particle size of loose powder is larger compared to compact powder. The particle size of loose powder is around 200 mesh, which is larger than that of compact powder (250 mesh), preventing it from entering and clogging the facial pores<sup>13</sup>. The composition of loose powder, including ingredients like lanolin, petrolatum, etc, is generally less than what is found in compact powder<sup>14,15</sup>.

The use of sunscreen does not have an impact on the occurrence of acne. This can happen because the majority of users apply it infrequently, typically once a day. Infrequent use of cosmetics results in lower absorption, minimizing the likelihood of acne development. The content of ingredients in sunscreen also plays a role, such as anhydrous preparations, emulsions, and oil-free preparations. Oil-free or non-greasy sunscreens minimize pore blockage, reducing the chances of acne formation<sup>16,17</sup>.

Foundations come in different types. Some types of foundation contain a significant amount of acne-causing or comedogenic ingredients, and using them on sensitive skin can lead to acne development. Using a foundation with fewer acne-causing or comedogenic ingredients is less likely to cause acne, as it does not clog facial pores<sup>18-20</sup>.

The frequency of use and skin type, whether normal or dry, rarely lead to acne. Additionally, cleanliness factors also play a role. Individuals who diligently clean their face after activities or after using cosmetics can minimize the occurrence of acne on the face<sup>18,19</sup>.

Blush comes in various types, including Loose or Compact Powder Rouges, Anhydrous cream rouge, Emulsion cream, and liquid rouges. If loose powder blush is used, it tends to be less likely to cause acne because the particle size is larger compared to compact powder types<sup>19-21</sup>. This is because the particle size of loose powder blush is larger, preventing it from clogging pores. Liquid rouge blush types also rarely cause acne. This is because the solvent is water or hydroalcoholic, making it safe for the skin. Another factor is the frequency of blush application, with the majority not using it every day. The less frequent the use of cosmetics, the less likely acne will occur, as the absorption into facial pores is reduced<sup>22,23</sup>.

---

## 5. Conclusion

From the results of a study on the influence of cosmetic use on the occurrence of acne vulgaris in female medical students at UKI FK in 2016, it can be concluded that there is a significant relationship only between the use of compact powder and the occurrence of acne vulgaris (AV). The use of loose powder, sunblock, foundation, and blush on was not found to be associated with the occurrence of AV.

---

## Compliance with ethical standards

### *Acknowledgement*

Thank you to all students in the Faculty of Medicine Class 2016 who have been involved and helped during the data collection process. Thanks are also conveyed to colleagues who have jointly completed this research.

### *Disclosure of conflict of interest*

No conflict of interest to be disclosed.

### *Statement of informed consent*

Informed consent was obtained from all individual participants included in the study. Ethical Clearance was obtained from Faculty of Medicine, Universitas Kristen Indonesia.

---

## References

- [1] Goh C, Cheng C, Agak G, et al. Acneiform Disorders. In: Kang S, Amagai M, L. Bruckner A et al. eds. Fitzpatrick's dermatology. 9<sup>th</sup> Ed. New York: McGraw-Hill, 2019.p.1391-1418

- [2] White GM. Recent findings in the epidemiologic evidence, classification, and subtypes of acne vulgaris. *Journal of the American Academy of Dermatology*. 1998 Aug, 39(2):S34-S37
- [3] Lynn DD, Umari T, Dunnick CA, Dellavalle RP. The epidemiology of acne vulgaris in late adolescence. *Adolesc Health Med Ther*. 2016, 7:13-25.
- [4] Harlim A, Pritarianti P A. The Relation Between Face Cleansing with Acne Vulgaris Case. *J Adv Res Dermatol Cosmetics*. 2023, 2:1-4
- [5] Do TT, Zarkhin S, Orringer JS, Nemeth S, Hamilton T, Sachs D, et al. Computer-assisted alignment and tracking of acne lesions indicate that most inflammatory lesions arise from comedones and de novo. *Journal of the American Academy of Dermatology*. 2008 Apr, 58(4):603-8.
- [6] Khunger N, Kumar C. A clinico-epidemiological study of adult acne: Is it different from adolescent acne?. *Indian J Dermatol Venereol Leprol*. 2012, 78(3):335.
- [7] Thiboutot D, Gilliland K, Light J, Lookingbill D. Androgen metabolism in sebaceous glands from subjects with and without acne. *Arch Dermatol*. 1999 Sep, 135(9):1041-5.
- [8] Lucky AW, Biro FM, Simbartl LA, Morrison JA, Sorg NW. Predictors of severity of acne vulgaris in young adolescent girls: results of a five-year longitudinal study. *J Pediatr*. 1997 Jan, 130(1):30-9.
- [9] Katoulis A, Kakepis E, Kintziou H, Kakepis M, Stavrianeas N. Comedogenicity of cosmetics: a review. *Acad Dermatol Venereol*. 1996 Sep, 7(2):115-9.
- [10] Ghani H, Rahman R, Liu K, Cubelli S. An Investigation of Makeup Ingredients and their Effects on Acne Cosmetics with Dermatologic Practice Recommendations. *J of Skin*. 2021 Sep 13, 5(5):474-81.
- [11] Kligman AM, Mills OH. "Acne cosmetica". *Arch Dermatol*. 1972 Dec, 106(6):843-50.
- [12] Khanna N, Gupta SD. Acneiform eruptions after facial beauty treatment. *Int J Dermatology*. 1999 Mar, 38(3):196-9.
- [13] Pribadi FW. Prarencana produksi bedak antiacne. Available from: <http://repository.wima.ac.id/3838/2/BAB%201.pdf>. January 24, 2017.
- [14] Yoo J, Kim H. Use of Beauty Products among U.S. Adolescents: An Exploration of Media Influence. *Journal of Global Fashion Marketing*. 2010 Aug, 1(3):172-81.
- [15] Singh S, Mann BK, Tiwary NK. Acne Cosmetica Revisited: A Case-Control Study Shows a Dose-Dependent Inverse Association between Overall Cosmetic Use and Post-Adolescent Acne. *Dermatology*. 2013, 226(4):337-41.
- [16] Andriana, R., Effendi, A., dan Berawi, K. N. The Correlation of Cosmetic Usage to Acne Vulgaris Case. Available from: <http://jukeunila.com/wp-content/uploads/2016/05/18-Risa-Andriana.pdf>. October 17, 2016.
- [17] Chilicka K, Rusztowicz M, Rogowska AM, Szyguła R, Nowicka D. Efficacy of Oxybrasion and Cosmetic Acids on Selected Skin Parameters in the Treatment with Acne Vulgaris. *CCID*. 2023 May, Volume 16:1309-17.
- [18] Zahra Ghodsi S, Orawa H, Zouboulis CC. Prevalence, Severity, and Severity Risk Factors of Acne in High School Pupils: A Community-Based Study. *Journal of Investigative Dermatology*. 2009 Sep, 129(9):2136-41.
- [19] Matsuoka Y, Yoneda K, Sadahira C, Katsuura J, Moriue T, Kubota Y. Effects of skin care and makeup under instructions from dermatologists on the quality of life of female patients with acne vulgaris. *The Journal of Dermatology*. 2006 Nov, 33(11):745-52.
- [20] Perera MPN, Peiris WMDM, Pathmanathan D, Mallawaarachchi S, Karunathilake IM. Relationship between acne vulgaris and cosmetic usage in Sri Lankan urban adolescent females. *J of Cosmetic Dermatology*. 2018 Jun, 17(3):431-6.
- [21] Conforti C, Giuffrida R, Fadda S, Fai A, Romita P, Zalaudek I, et al. Topical dermocosmetics and acne vulgaris. *Dermatol Ther*. 2021 Jan, 34(1):e14436.
- [22] Suh D, Oh H, Lee SJ, Kim HJ, Ryu HJ. Relationship between acne and the use of cosmetics: Results of a questionnaire study in 539 Korean individuals. *J of Cosmetic Dermatology*. 2021 Jul, 20(7):2172-8.
- [23] Zhao J, Wang Y, Jiang L, Mu YZ. The application of skin care product in acne treatment. *Dermatol Ther*. 2020 Nov, 33(6):e14287.