

Frica Ronauli Batubara dkk  
(The Relationship between  
Physical Fitness Level with Body  
Mass Index of Medical  
Students at Indonesian  
Christian University, Jakarta)

*by* Library Referensi

---

**Submission date:** 03-Jul-2024 05:24PM (UTC+0700)

**Submission ID:** 2412010844

**File name:** Medical\_Students\_at\_Indonesian\_Christian\_University,\_Jakarta.pdf (278.54K)

**Word count:** 4486

**Character count:** 24035



# The Relationship between Physical Fitness Level with Body Mass Index of Medical Students at Indonesian Christian University, Jakarta

Frica Ronauli Batubara <sup>a\*</sup>, Raden Revianto Ananto Putro <sup>a</sup>  
and Wendy Hendrika <sup>a</sup>

<sup>a</sup> Faculty of Medicine, Universitas Kristen Indonesia, Jakarta, Indonesia.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijtdh/2024/v45i71564>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119031>

Original Research Article

Received: 20/04/2024

Accepted: 22/06/2024

Published: 25/06/2024

## ABSTRACT

**Aims:** This research aims to understand the relationship between physical fitness level and body mass index (BMI) of students at the Faculty of Medicine, Indonesian Christian University, especially the class of 2021.

**Study Design:** The research design used in this study was observational with a cross-sectional approach.

**Place and Duration of Study:** This research will be conducted at the Faculty of Medicine, Indonesian Christian University Laboratory in December 2022.

**Methodology:** The population in this study were all students from the Faculty of Medicine, class of 2021. The research sample consisted of 57 students from the Faculty of Medicine, class of 2021,

\*Corresponding author: Email: [friscabatubara@yahoo.co.id](mailto:friscabatubara@yahoo.co.id);

**Cite as:** Batubara, Frica Ronauli, Raden Revianto Ananto Putro, and Wendy Hendrika. 2024. "The Relationship Between Physical Fitness Level With Body Mass Index of Medical Students at Indonesian Christian University, Jakarta". *International Journal of TROPICAL DISEASE & Health* 45 (7):96-103. <https://doi.org/10.9734/ijtdh/2024/v45i71564>.

who were selected using a sampling technique, namely consecutive sampling. The physical fitness test uses the Harvard Step Test.

**Results:** From the research results, it was found that the physical fitness level of 57 students registered at the Faculty of Medicine, Indonesian Christian University could be categorized as follows: 44 students (77.2%) were in the less ability, 8 students (14.0%) included in the moderate ability, and 5 students (8.8%) were included in the good ability. Apart from that, an analysis of the body mass index (BMI) of students at the Faculty of Medicine, Indonesian Christian University, Class of 2021, showed that the largest proportion consisted of individuals who were classified as having a normal weight, namely 31 people (54.4%), obesity, 12 people (21.1%), underweight as many as 6 people (10.5%), overweight as many as 4 people (7.0%) and Obesity II as many as 4 people (7.0%). Based on bivariate analysis, the results showed that there was no significant relationship between Physical Fitness and Body Mass Index. This was concluded by obtaining a chi-square value, namely  $p = 0.740$ .

**Conclusion:** From the research results, it can be concluded that the Body Mass Index is not a determining factor that guarantees physical fitness, likewise other moderate variables such as age and gender also have no relationship with physical fitness.

*Keywords: Physical fitness; body mass index; relationship.*

## 1. INTRODUCTION

### 1.1 Background

Physical fitness refers to a state where a person can carry out activities with optimal effectiveness and good efficiency without feeling high fatigue. Physical fitness plays an important role in a person being involved in physical activities and sports, thus having a positive influence on their cognitive, psychomotor, and affective development [1].

It is sought that people who have a high level of physical fitness will demonstrate increased receptivity to instruction, resulting in better academic performance. Physical fitness refers to the capacity to perform tasks effectively, even under challenging conditions, which cannot be achieved by individuals who have poor physical fitness. Physical fitness is a fundamental and crucial asset for humans [2]. Physical fitness assessment can be done through a fitness test. The experiment began with respondents engaging in a fitness assessment known as the Harvard Step Test. This test aims to evaluate students' physical performance abilities by measuring their pulse rate with their current physical fitness condition [3]. The main purpose of conducting fitness test experiments, especially the Harvard Step Test, is to assess the respondents' aptitude, physical fitness, and work capacity. Next, understand the correlation between physical fitness level and Body Mass Index (BMI) [3-7].

A person's physical fitness level can naturally increase, but it can also decrease independently

when a person does minimal physical activity. Body mass index (BMI) is recognized as a significant determinant of a person's level of physical fitness [8]. Body Mass Index (BMI) refers to a direct way to assess the nutritional status of individuals aged 18 years and over. It calculates the ratio of body weight to height, without distinguishing the contribution of muscle mass, bone, fluid, or fat to the overall weight [9]. According to the researchers' writing, this study intends to examine the correlation between physical fitness level and body mass index for students enrolled at the Faculty of Medicine, Indonesian Christian University.

### 1.2 Research Problem

The research problem is whether a relationship exists between physical fitness level and body mass index among students at the Faculty of Medicine, Indonesian Christian University Class of 2021.

### 1.3 Research Purposes

This research aims to determine the correlation between physical fitness level and body mass index in students at the Faculty of Medicine, Indonesian Christian University Class of 2021.

### 1.4 Benefits of Research

#### 1.4.1 Benefits for health facilities

It is hoped that it will help and educate clinicians to be able to determine the patient's fitness condition with the Harvard Step Test and it is hoped that in the future it can be applied in primary health facilities.

#### 1.4.2 Benefits to society

This research helps share explanations with the public so that it can be a means of education in maintaining and monitoring physical fitness by practicing the Harvard Step Test.

#### 1.4.3 General research benefits

It is hoped that the results of this research can be used as a reference in developing this research further, and It is hoped that the Harvard Step Test can become a reliable test for testing physical fitness

## 2. MATERIALS AND METHODS

### 2.1 Material

#### 2.1.1 Definition of physical fitness

The level of physical fitness can be assessed based on an individual's ability to carry out daily activities without feeling tired. Physical fitness is generally associated with an individual's capacity and ability to perform routine tasks consistently, efficiently, and effectively over a long period without experiencing fatigue, thereby enabling them to maintain their energy to engage in additional activities [10]. Physical fitness refers to the ability to carry out daily activities with energy and attention. Therefore, individuals can maintain sufficient energy levels to engage in recreational activities and manage unexpected crises effectively without experiencing excessive fatigue [11]. If a person can perform his daily tasks with high efficiency and minimal fatigue. Individuals demonstrate a good level of physical fitness. A person can still derive pleasure from their leisure time [12].

#### 2.1.2 The physical fitness test uses the harvard step test

The Harvard Step Test is a cardiovascular endurance assessment that evaluates the physiological capacity of the heart and lungs. This assessment can measure an individual's aptitude to sustain prolonged physical activity without experiencing fatigue. Participants were instructed to climb up and down the bench at a height of 40 cm. Participants were instructed to perform a series of repetitions on a bench, alternating between up and down movements, for five minutes. This exercise continues until the respondent reaches a state of exhaustion, rendering them unable to participate further in

the test. Fatigue occurs when the respondent cannot maintain his stride for 15 seconds. After the respondent sits down and the test is completed, their pulse will be counted within 1, 2, and 3 minutes [13,14]. There are many examples of terminology, such as cardiorespiratory endurance, cardiovascular endurance, aerobic power, cardiovascular endurance, cardiorespiratory endurance, and aerobic fitness, which have the same definition [15,16].

The Harvard Step Test classification or what can be called the physical fitness index (IKJ) is divided into 3, namely poor ability, moderate ability, and good ability [17,18].

#### 2.1.3 Body mass index

Body Mass Index (BMI) is a widely used approach to assess an individual's weight status and potential health problems. This involves calculating an individual's BMI by dividing their body mass in kilograms by the square of their height in meters. Additionally, body mass index (BMI) is a direct, economical, and non-intrusive approach. Body Mass Index (BMI) fails to account for variations in muscle mass and body fat distribution [19,20].

According to the World Health Organization (WHO), the classification of body mass index (BMI) consists of five categories. These categories are as follows: underweight, indicated by a BMI of less than 18.5; normal body weight, indicated by a BMI ranging from 18.5 to 22.9; overweight with associated risks, indicated by a BMI ranging from 23 to 24.9; class I obesity, indicated by a BMI ranging from 25 to 29.9; and class II obesity, indicated by a BMI equal to or greater than 30 [21].

**Table 1. Harvard step test classification**

Classification	Score
Poor Ability	<50
Moderate Ability	50 - 80
Good Ability	>80

**Table 2. BMI classification**

Classification	BMI
Underweight	<18,5
Normal Body Weight	18,5 – 22,9
Being Overweight with Risks	>23 – 24,9
Obesity I	25 – 29,9
Obesity II	≥30

5

## 2.2 Methods

### 2.2.1 Research design

The research design used in this study was observational with a cross-sectional approach.

### 2.2.2 Place and time

This research will be carried out at the Laboratory of the Faculty of Medicine, Indonesian Christian University on December 14 2022

### 2.2.3 Population and sample

The research focused on the student population of the Indonesian Christian University Faculty of Medicine who were actively registered in the Class of 2021.

This research used a portion of the 138 students currently registered at the Faculty of Medicine, Indonesian Christian University Class of 2021 which was calculated using a sample formula, namely the Slovin formula. After calculating using the Slovin formula, the required sample was 57 people. The chosen methodology uses a non-probability approach, specifically using a consecutive sampling design to determine people from the population who meet certain conditions as samples.

### 2.2.4 Inclusion and exclusion criteria

Inclusion Criteria

- a. Active student at the Faculty of Medicine, Indonesian Christian University Class of 2021
- b. Willing to do research
- c. Aged 18-21 Years
- d. Male and female gender

Exclusion Criteria: There were no exclusion criteria in this study.

### 2.2.5 Data collection

Before taking the test, participants are given a pre-test briefing that covers the protocol and related aspects related to the Harvard Step Test. The experimental procedure involved traversing a 40 cm platform or ladder cyclically, synchronized with a metronome set at intervals of 120 beats per minute. The above process is carried out continuously until the respondent's strength decreases, provided that the duration of the process does not exceed 300 seconds. The

test is considered invalid if the execution of the steps is not in sync with the rhythm of the metronome or if the participant experiences a fall during the process of setting up and lowering the bench. The Indonesian Christian University laboratory will be used to carry out measurements.

### 2.2.6 Data analysis

#### 2.2.6.1 Univariate analysis

Univariate analysis aims to explain the attributes of each variable evaluated through the use of frequency distribution tables.

#### 2.2.6.2 Bivariate analysis

Bivariate analysis uses cross-tabulation as a tool to test potential correlations or relationships between two variables, especially the independent variable and the dependent variable. This research uses the Pearson statistical correlation coefficient to analyze research data. The chi-square test is a statistical method used to determine the significance of the relationship between categorical variables. A correlation is considered significant when the p-value is less than 0.05, indicating a statistically significant association between two dependent variables. When the p-value exceeds 0.05, this indicates that no statistically significant correlation was obtained between the two variables under consideration

## 3. RESULTS AND DISCUSSION

### 3.1 Results

This research aims to see whether or not there is a relationship between the level of physical fitness and the Body Mass Index (BMI) of students at the Faculty of Medicine, Indonesian Christian University Class of 2021. The research was conducted on December 14 2022 at the UKI FK Laboratory. This research sample was taken using the Slovin formula method and the results obtained were 57 people. The results of the analysis from this research are as follows.

#### 3.1.1 Univariate analysis: Distribution of respondents based on body mass index (BMI)

Frequency Distribution Based on Body Mass Index can be presented in Table 3:

5 Based on the frequency data in Table 3, it can be seen that the BMI frequency distribution of 2021

FK UKI student respondents is dominated by students with a normal weight classification, 31 people (54.4%). Obesity I was 12 people (21.1%), Underweight was 6 people (10.5%), Overweight was 4 people (7.0%) and Obesity II was 4 people (7.0%).

**3.1.2 Univariate analysis: Frequency distribution of physical fitness**

The frequency distribution of respondents based on the results of the physical fitness test using the Harvard Step Test is presented in Table 4:

**3.1.3 Bivariate analysis: Relationship between physical fitness and body mass index**

The results of bivariate analysis using the chi-square test to see whether there is a relationship between body mass index and physical fitness are presented in Table 5.

Based on the data from the analysis in Table 5, it can be seen that the Chi-Square calculation results obtained ( $p=0.740$ ) which can be interpreted as meaning that the relationship between physical fitness level and BMI is not

proven to be related where the value shows  $>0.05$ .

**3.2 Discussion**

Based on research that has been conducted regarding the relationship between physical fitness level and Body Mass Index (BMI) among 57 students at the Faculty of Medicine, Indonesian Christian University Class of 2021, calculated using the Slovin formula, it is known that the average age of the students used as a sample is 19 years, the average body mass index is 25.25 kg/m<sup>2</sup>, and the average physical fitness as measured through the Harvard Step Test regarding the Physical Fitness Index (IKJ) score is 44.12.

Furthermore, the Body Mass Index (BMI) of each student was categorized into five different groups, namely underweight, normal, overweight, obesity level I, and obesity level II. A total of 6 people (10.2%) showed a BMI that was included in the thin category, while 31 people (54.4%) were classified as having normal weight. In addition, 4 people (7.0%) were categorized as overweight, 12 people (21.1%) were classified as obese group I, and 4 people (7.0%) were classified as obese group II.

**Table 3. Frequency distribution based on body mass index**

BMI Classification	Number of Respondent	Percentage (%)
Underweight	66	10.3
Normal Body Weight	31	54.4
Being Overweight with Risks	4	7.0
Obesity I	12	21.1
Obesity II	4	7.0
<b>Total</b>	<b>57</b>	<b>100</b>

**Table 4. The frequency distribution of respondents based on the results of the physical fitness test using the harvard step test**

Physical Fitness Classification	Number of Respondent	Percentage (%)
Poor Ability	44	77.2
Moderate Ability	8	14.0
Good Ability	5	8.8
<b>Total</b>	<b>57</b>	<b>100</b>

**Table 5. The results of bivariate analysis using the chi-square test to see whether there is a relationship between body mass index and physical fitness**

Chi-Square Tests	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	5.165 <sup>a</sup>	8	.740

a. 23 cells (92.0%) have an expected count of less than 5. The minimum expected count is .07.

1 The physical fitness of students at the Faculty of Medicine, Indonesian Christian University Class of 2021 was measured using the Harvard Step Test and the results of the Physical Fitness Index (IKJ) were obtained. Next, each student is grouped into groups based on their level of physical fitness, namely: low, medium, and high. There were 44 students (77.2%) in the low group, 8 students (14.0%) in the medium group, and 5 students (8.8%) in the high group. This shows that students at the Faculty of Medicine, Indonesian Christian University, Class of 2021, have poor physical fitness on average.

The findings of this research, as determined by the Pearson chi-square correlation test, obtained a significance value (p) between physical fitness level and Body Mass Index (BMI) of (p=0.740). This shows that the level of physical fitness and Body Mass Index (BMI) are not related. A 2023 study in India by Chinmoyee Baruah Hazra et al. revealed a significant relationship between physical fitness and Body Mass Index (BMI), as assessed through the Harvard Step Test experiment, and the calculation of Body Mass Index (BMI) [22,23].

This research is in line with a previous study conducted in 2021 by Aprianto D et al. which examined the correlation between body mass index and the level of physical fitness among staff at the Ministry of Health. This study used the Pearson Chi-square correlation test to examine the relationship between Body Mass Index (BMI) and the level of physical fitness of Ministry of Health employees. The analysis findings show that there is no significant correlation between BMI and physical fitness level (p=0.159) [24,25].

The differences between the two studies can be influenced by several factors, one of which is the number of samples used, the samples used by Chinmoyee Baruah Hazra et al. had as many as 100 while in this study there were only 57 people, and apart from that, according to Aprianto D et al. [25]. Activities before the pre-test can also influence the results of this test. One of the obstacles in this research is the absence of measurements of various variables that could potentially influence the results of this test. For example, factors such as daily exercise intensity and level of physical activity can influence a person's physical fitness. Apart from that, lifestyles such as smoking, genetic heredity, nutrition, and pre-test student activities can also influence the results of this research.

## 5 4. CONCLUSION

Based on the research results, it can be concluded that Body Mass Index does not have a significant relationship with Physical Fitness. This is proven by the results of bivariate analysis with a p-value = 0.740 > sig 0.05.

## 5. SUGGESTION

It is recommended that future researchers carry out additional investigations, using more accurate and comprehensive measurement tools, to evaluate potential confounding variables, such as physical activity and exercise, which influence the amount of physical fitness and can also add a pre-test questionnaire before conducting the Harvard experiment. Step Test. Apart from that, efforts are also needed to improve physical fitness, which is one way to involve yourself in sustainable and rhythmic sports activities that involve most of the body's muscles. Examples of such activities include running, swimming, and gymnastics.

## 8 DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## CONSENT AND ETHICAL APPROVAL

It is not applicable.

## ACKNOWLEDGEMENTS

We would like to thank the faculty of medicine for facilitating us in conducting research, especially the students of the faculty of medicine class of 2021 who were willing to be sampled in this research.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Rozi F, Safitri SR, Syukriadi A. Evaluation of students' physical fitness levels at physical education lectures at IAIN salatiga. *Edutrained Journal: Journal of*

- Education and Training. 2021, July 6;5(1):13–8.
2. Gempur Santoso. Performance Ability Using the Harvard Step Test. *WAHANA*. 2020, Jun 1;72(1):67–70. Available:<http://jurnal.unipasby.ac.id/index.php/whn/article/view/2371>
  3. Aprianto DP, Nurwahyuni A. The relationship between body mass index (BMI) and the physical fitness level of ministry of health employees. 2021;9. Available: <http://ejournal.uika-bogor.ac.id/index.php/Hearty/issue/archive>
  4. Prihantini NN, Adintia M, Utomo TY. Relationship between body mass index and low-density lipoprotein and triglycerides in type 2 diabetes mellitus patients. *Asian J. Med. Health*. 2023, Apr 21;21(7):65-74. [cited 2024 Jun. 13] Available:<https://journalajmah.com/index.php/AJMAH/article/view/843>
  5. Hurissi E, Allami M, Alhazmi A, Alsomaili H, Gharawi S, Kamili A, Homadi J, Ageeli F, Moafa A, Abdelmola A, Zanquti A, Alhazmi AH, Ghulaysi S. Prevalence of obesity and its association with academic performance and other factors among jazan university medical students: A cross-sectional study. *J. Adv. Med. Med. Res*. 2024, Jan 5;36(1):1-9. [cited 2024 Jun. 13] Available:<https://journaljamr.com/index.php/JAMMR/article/view/5347>
  6. Vargas PA, Flores M, Robles E. Sleep quality and body mass index in college students: The role of sleep disturbances. *Journal of American college health*. 2014, Nov 17;62(8):534-41.
  7. Grasdalsmoen M, Eriksen HR, Lønning KJ, Sivertsen B. Physical exercise and body-mass index in young adults: A national survey of Norwegian university students. *BMC public health*. 2019, Dec;19:1-9.
  8. Ilyas I, Almunawar A. Physical fitness profile of volleyball extracurricular students. *indonesian journal of sports and health*. 2020, December 31;1(1):37–45. Available:<http://jurnal.stokbinaguna.ac.id/index.php/JOK/article/view/297>
  9. Munipiddin M, Tamim MH, Nopiana R. The Relationship between nutritional status and physical fitness level of class X Male students at MA NW Kalijaga. *Porkes Journal*. 2018, December 31;1(2):52–61. Available:<http://ejournal.hamzanwadi.ac.id/index.php/porkes/article/view/1414>
  10. Pranata D, Kumaat N. The influence of sports and physical training models on adolescent physical fitness: Literature review. *Journal of Sports Medicine*. 2022;10(02):107–16.
  11. Wiranata Y, Inayah I. Comparison of body mass calculation using the body mass index (BMI) and bioelectrical impedance analysis (BIA) methods. *Journal of Health Management of the RSDr Soetomo Foundation*. 2020, April 23;6(1):43. Available:<http://jurnal.stikes-yrsds.ac.id/index.php/JMK/article/view/280>
  12. Hazra CB. Correlation of body composition (Body Mass Index, Waist-To-Hip Ratio, Waist-To-Height Ratio, Neck Circumference) with fitness index using harvard step test. *Med Rxiv*. 2023, January 1;2023.07.13.23292635. Available:<http://medrxiv.org/content/early/2023/07/16/2023.07.13.23292635.abstract>
  13. Utami D. Factors influencing body mass index in adolescents aged 15-18 years at SMAN 14 Tangerang. *Journal of Medical and Health Sciences*. 2017;4(3).
  14. Khurde N, Jibhkate A, Udhan V, Khurde S. A gender-based comparative cross-sectional study of physical fitness index using Harvard's step test in the medical students of Western India. *Natl J Physiol Pharm Pharmacol*. 2021;(0):1.
  15. Mahfud I, Gumantan A, Fahrizqi EB. Analysis of BMI (body mass index) of Indonesian teknocrat university football ukm athletes. *Satria: Journal of Sports Athleticism in Teaching and Recreation on Interdisciplinary Analysis*. 2020;3(1):9-13.
  16. Yhuwono Y. The relationship between body mass index and blood pressure in the elderly in Pesucen Village, Banyuwangi. *Indonesian Journal of Public Health*. 2018;13(1).
  17. Kamaruddin I. Body Mass Index (BMI) on cardiovascular endurance. *Sportive: Journal Of Physical Education, Sport and Recreation*. 2020;3(2):117-22.
  18. Niswatin D, Cahyawati WA, Rosida L. Literature review: Relationship between body mass index (BMI) and muscle mass in the elderly. *Homeostasis*. 2021, Apr 30;4(1):171-80.
  19. Abineno AP, Malinti E. Relationship between body mass index and blood pressure in adults. *Indonesian Journal of Nursing and Health Sciences*. 2022, Apr 28;3(1):37-42.



20. Lengkana AS, Muhtar T. Physical fitness learning. CV Greetings Noble People; 2021, Mar 11.
21. JM Journal, Syaquy A. The relationship between body mass index and physical fitness of UNJA medical study program students. Jambi Medical Journal: Journal of Medicine and Health. 2017, Jun 2;5(1).
22. Rousdyanto MH, Ratimiasih Y, Ardiyanto A. Correlation between physical activity and body mass index (BMI) and the physical fitness level of Class XI students for the 2020/2021 academic year at Sma Negeri 3 Demalang. Journal of Primary and Secondary Education. 2021, Jul 30;1(1):15-21.
23. Aprianto DP, Nurwahyuni A. The relationship between body mass index (BMI) and the physical fitness level of ministry of health employees. Hearty: Journal of Public Health. 2021, Aug 25;9(2):49-57.
24. Sari RK, Hadi JP, Wijyaningrum L. The relationship between body mass index and physical fitness level by measuring Vo2 max in diving fishermen in Kedung Cowek subdistrict, Surabaya. Hang Tuah Medical Journal. 2019, May 17;16(2):196-203.
25. Juliyanty NK, Indonesiani SH, Suryanditha PA. Relationship between physical activity level and body mass index on cardiorespiratory endurance in students at the Faculty of Medicine and Health Sciences, Warmadewa University. Aesculapius Medical Journal. 2022, Oct 21;2(3):143-9.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119031>

# Frica Ronauli Batubara dkk (The Relationship between Physical Fitness Level with Body Mass Index of Medical Students at Indonesian Christian University, Jakarta)

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

20%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://repository.uki.ac.id">repository.uki.ac.id</a> Internet Source	5%
2	Submitted to Forum Komunikasi Perpustakaan Perguruan Tinggi Kristen Indonesia (FKPPTKI) Student Paper	3%
3	<a href="http://www.science.gov">www.science.gov</a> Internet Source	2%
4	<a href="http://www.unud.ac.id">www.unud.ac.id</a> Internet Source	2%
5	<a href="https://eprints.unm.ac.id">eprints.unm.ac.id</a> Internet Source	1%
6	Submitted to Universitas Dian Nuswantoro Student Paper	1%
7	<a href="http://journal.unj.ac.id">journal.unj.ac.id</a> Internet Source	1%
8	<a href="http://reunir.unir.net">reunir.unir.net</a> Internet Source	

1 %

9

Submitted to University of Hong Kong

Student Paper

1 %

10

science.manuscript2send.com

Internet Source

1 %

11

scales.arabpsychology.com

Internet Source

1 %

12

ijrti.org

Internet Source

1 %

13

www.ejournal.warmadewa.ac.id

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On