

THE EDUCATION PROGRAMS ENHANCING PHYSICAL ACTIVITY: SYSTEMATIC LITERATURE REVIEW

by Library Referensi

Submission date: 18-Jul-2025 10:40AM (UTC+0700)

Submission ID: 2716625387

File name: artikel_IJMAES_terbit.pdf (954.69K)

Word count: 4077

Character count: 22392



International Journal of Medical and Exercise Science

(Multidisciplinary, Peer Reviewed and Indexed Journal)

ORIGINAL ARTICLE

THE EDUCATION PROGRAMS ENHANCING PHYSICAL ACTIVITY: SYSTEMATIC LITERATURE REVIEW

Search engine:
www.ijmaes.org

Citra Puspa Juwita^{1*}, Weeke Budhyanti², Bernadetha Nadeak³

Corresponding Author:

¹Physiotherapy Program, Faculty of Vocational Studies, Universitas Kristen Indonesia, Jakarta, Indonesia, E-mail: citra.simatupang@uki.ac.id

Co Authors:

²Physiotherapy Program, Faculty of Vocational Studies, Universitas Kristen Indonesia, Jakarta, Indonesia

³Magister of Education Management, Postgraduate Program, Universitas Kristen Indonesia, Jakarta, Indonesia

ABSTRACT

Background: Doing regular physical activity has many benefits, but the reality is people do not do it. Education is one of the efforts to promote sufficient physical activity in society. The study aims to provide a systematic review of the impact of educational program enhancing in physical activity. **Methods:** A systematic literature review using preferred reporting items for systematic reviews. A precise search of the journal database was identified and thematic analysis was used to summarize the findings. The study using three databases, conducted from 2012 to 2022, language in English, the research design was an experimental study, and using educational program interventions. We used Boolean with keywords: Physical activities AND education. The articles obtained are downloaded and analysis by the authors. The analysis was conducted by comparing the increase in physical activity after the intervention. **Results:** Nine articles from the database were eligible for review, it was found that education was provided in many models, such: (1) consultation; (2) distributing physical activity videos; (3) environmental modification; (4) assisting physical education; (5) improving healthy eating and active living policies; (6) specific exercise; (7) face to face training and educational booklet; (8) The Pender's HPM (nine 4-h training sessions and consulting support via telephone contact and social media group; and (9) based on socio-ecological model and conducted on 4 levels. Comprehensive education using a socio-ecological model is suitable for improving the changing physical activity (2612,15 MET-min/week). **Conclusions:** Participants who were given a comprehensive education program and behavior change theory produced high changes in physical activity.

Keywords: Education, behavior, exercise, activity, training.

Received on 18th February 2025; Revised on 24th February 2025; Accepted on 28th February 2025
DOI:10.36678/IJMAES.2025.V11I01.016

INTRODUCTION

Physical activity is one of many factors that affect non-communicable diseases. Sufficient physical activity may improve health¹, improve individual mental health, and well-being². In 2020, there were 3.9 million³ and in 2015 there were 5.3 million deaths related to sedentary lifestyles. Other data shows that between 2001-2016, one in four adults did not do as much physical activity as they should⁴, especially during the Covid pandemic, where movement is limited which results in people being affected by joint disease⁵. WHO recommends that all countries set physical activity targets in national guidelines and tools, thereby helping to support all age groups to maintain their health⁶.

Physical activity promotion is encouraged worldwide, including in developing countries, Indonesia has made efforts to promote sufficient physical activity among its citizens through education, sports-promoting campaigns, increasing provision of physical activity facilities, encouraging motor vehicle-free areas to facilitate physical activities, Health Community Movement campaign, and encouraging connectivity between modes of public transportation⁷. It is said that sufficient physical activity is adjusted to age and health condition, as 5-17 years old individuals should exercise three times a week for 60 minutes, and above 18 years old individual should exercise 150-300 minutes of moderate intensity per week, and individuals with poor health condition may suit times and types of exercises according to their physical fitness⁸. Providing appropriate physical activity with self-efficacy has been shown to result in longer adherence⁹, another benefit is that it results in

physical, psychological, social, spiritual, and total health¹⁰.

Promoting physical activity requires resources and collaboration with all stakeholders, including education for the community¹¹. Community education about active lifestyle is a more popular solution, compared with other promotional activities, as it may be done anywhere, anytime, and to anyone with affordable resources. Many educational interventions have been carried out to increase physical activity in the community, but the summary is still needed. This research aims to provide a systematic review of educational programs that result in changes in physical activity.

METHODS

This research is a systematic literature review with Preferred Reporting Items for Systematic Reviews & Met Analyses (PRISMA) 2009¹².

Literature search strategy: Precise criteria and strategies for searching on journal databases were identified and thematic analysis was used to summarise findings. A systematic review was conducted between June and August 2022 using three electronic databases. The criteria include a) from three databases such as PubMed, Web of Science, Scopus, and others published, b) conducted from 2012 to 2022, c) using the English language, d) using experiment study; and e) using education program as intervention. We used Boolean with the keyword: Physical activities AND education. The article found will be downloaded and analysis. Eligibility criteria are stated from the research question through the Problem, Intervention, Comparative, Outcome, and Studies (PICOS) assessment¹².

Data extraction: Data was extracted and verified by the researcher using PICOS forms. The article obtained that matches the criteria will be summarized in a table, which contains: (1) author's name, (2) title, (3) country and year of research, (4) intervention models, (5) sampling method and number of samples, (6) outcomes related to physical activity, and (7) pre and post-intervention results and change results. Critical reviews will be analysed one by one through in-depth discussions between researchers.

Bias Risk Evaluation: To reviewer (CPJ, WB, BN) independently assesses each article using the Robvis tools approach, i.e. randomization bias, intervention bias, lost data, outcome measurement, and result selection 13. From

five domains from nine eligible articles were produced 31% with some concern bias and 69% low risk bias%.

RESULTS

The result of three journal databases and other sources was 157 articles that meet the criteria. After reading all of the text, 15% of the articles were excluded due to duplication, 69% were excluded as not meeting the PICOS criteria, and 24 articles were eligible. As not all the articles can open, only 15 articles may be analyzed. By our analysis of full-text articles, we found 6 articles rejected from the research objectives, thus only 9 articles may be analyzed qualitatively (see picture 1).




Figure 1. Flowchart Article Selection Process

Participants of the research study consisted of children and adults with varying health conditions, such as health, autism, and low

back pain. As the results of eligible articles, it was found that apart from providing knowledge, education was also provided in

other forms, i.e: (1) consultation resulted in a change of 178 MET-min/week, (2) distributing physical activity videos using private WhatsApp groups and videos resulted in a change of 31.19 MET-min/week, (3) environmental modification produces changes of 60,2 MET-min/week; (4) Assisting physical education resulted in a change of 45,1 point; (5) Caregiver to improve healthy eating and active living policies resulted in a change of 1,24 point; (6) Low-Load Motor Control (LMC) and High Load Lifting (HLL) exercise resulted in a

mean of change of 4,2 point of LMC and 2,5 point of HLL; (7) Training face to face and educational booklet resulted in a change of 1263.68 MET-min/week; (8) The Pender's HPM (nine 4-h training sessions and consulting support via phone contact and social media group) resulted in a change of 37 MET-min/week, dan; (9) Based on socioecological model and conducted on 4 levels of personal, social, organizational, and political resulted in a change of 2612.15 MET-min/week.

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
1	Rakhshani T, Khiyali Z, Masrurpur F, Khani Jeihooni A ¹⁴	Effect of Educational Intervention on improvement of physical activities of middle-aged woman	Iran, 2017	Education al (Consultation, group discussion, evaluation, reaching goal)	Two-stage cluster sampling method, 160, middle-aged women	Physical activities = International Physical Activity Questionnaire (IPAQ)	Pre= 548.30 MET-min/week Post= 726.30 MET-min/week $\Delta = 178$ MET-min/week
2	Yarımkaç E, Esentürk OK, İlhan EL, Karasu N ¹⁵ .	A WhatsApp-delivered intervention promote physical	Turkey, April 27 – June 21, 2020	A private WhatsApp group and videos	Random sampling, 42, children	Physical activity= Leisure Time Exercise Questio	Pre= 20.04 MET-min/week Post= 51.23 MET-

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
		activity in young children with autism spectrum disorder				nnaire (LTEQ)	min/week Δ = 31.19 MET-min/week
3	Morais LDC, Paravidino VB, Mediano MFF, et al ¹⁶ .	Effectiveness of a school-based randomized controlled trial aimed at increasing physical activity time in adolescents	Brazil, 2016	PG: general basis for a healthy lifestyle, through educational games, group debates and culinary classes focused PEG: PG + environment	Random sampling, 2511, adolescents	Physical activity= physical activity questionnaire for the adolescent	Pre = 219.9 MET-min/week Post = 280.1 MET-min/week Δ = 60.2 MET-min/week
4	Gill M, Roth SE, Chan-Golston AM, et al ¹⁷	Evaluation of an intervention to increase physical activity in low-income,	Los Angeles; October 2014, January & March 2015	Assisting physical education teacher using PE curriculum (SPARK PE)	561 (287 intervention and 274 control), middle schools	Moderate-to-vigorous physical activity (MVPA)= System for Observing	Pre =16.0 MET-min/week Post = 61.1 MET-min/week

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
		urban middle schools				g Fitness Instruction Time (SOFIT)	$\Delta = 45.1$ MET-min/week
5	Slining M, Wills S, Fair M, et al ¹⁸	LiveWell in early childhood: results from a two-year pilot intervention to improve nutrition and physical activity policies, systems and environments among early childhood education programs in South Carolina	Greenville, USA. June 2016 and August 2018	ECE center directors and caregivers to improve healthy eating and active living policies and practices	Randomly, 120, 3-5-year-old	PA policy= Environment and Policy Assessment and Observation (EPAO)	Pre = 11.76 MET-min/week Post = 13 MET-min/week $\Delta = 1.24$ MET-min/week
6	Aasa B, Berglund L, Michaelso	Individualized Low-Load Motor Control	Sweden, 2014	LMC exercises: low load lifting	Randomization, 70 participants, 25-60	Patient-specific function	LMC Pre = 3.8 MET-

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
	n P, Aasa U ¹⁹	Exercises and Education Versus a High Load Lifting Exercise and Education to Improve Activity, Pain Intensity, and Physical Performance in Patients with Low Back Pain: A Randomized Controlled Trial		exercise and high load lifting exercise	years	al scale	min/week Post= 8 MET-min/week HLL Pre= 4.8 MET-min/week Post= 7.3 MET-min/week Δ = 4.2 and 2.5 MET-min/week
7	Khodaveisi M, Azizpour B, Jadidi A, Mohammadi Y ²⁰	Education based on the health belief model to improve the level of physical activity	Iran, February 2018 to February 2019	Training face-to-face and educational booklet	convenience sampling method from different faculties, 130 participants	Physical activity= International Physical Activity Questionnaire (IPAQ).	Pre= 2,679.47 MET-min/week Post= 3,943.15 MET-

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
					ts, aged 25–50 year		min/week Δ = 1,263.68 MET-min/week
8	Vahedian Shahroodi M, Belin Tavakoly Sany S, Hosseini Khaboshan Z, Esmaeily H, Jafari A, Tajfard M ²¹	Effect of a theory-based educational intervention for enhancing nutrition and physical activity among Iranian women: a randomised control trial	Iran, April 2016 to February 2017	The Pender's HPM nine 4-h training sessions and consulting support via phone contact and social media group	202, 18-50 years	Physical activity= International Physical Activity Questionnaire (IPAQ-L)	Mean intervention = 1840 MET-min/week Mean control= 1803 MET-min/week Mean Δ = 37 MET-min/week
9	Tehrani H, Majlessi F, Shojaeizadeh D, Sadeghi R, Kabootarkhani	Applying Socioecological Model to Improve Women's Physical Activity: A Randomized Control	Iran, NA	Education al multimedia and conducted on 4 levels of personal, social,	360, 22-40 years	Physical activity= international physical activity questionnaire (IPAQ)	Pre = 992.17 MET-min/week Post = 3604.32 MET-min/week

No	Study	Title	Country And Year of Research	Form Of Intervention	Sampling Method and Number of Samples	Outcomes Related to Physical Activity	Pre And Post Intervention Results and Change Results (Δ)
	MH ²²	Trial		organizational, and political			$\Delta = 2612.15$ MET-min/week

Table 1 The Characteristic Article

9

DISCUSSION

The purpose of this study was to determine the review of the impact of educational programs on physical activity based on research that has been done. The results of the Systematic literature review found that participants consisted of children, adolescents, and adults. Health conditions in research subjects from nine articles also varied, some were healthy, autistic, eye pain, and low back pain. The results of the review found that those who produced high physical activity change outcomes were given a comprehensive intervention and a behavior change approach.

The expected result of the educational program is a change in better behavior, in this case, physical activity. The intervention in the article review found that two interventions that use behavior change theory, namely the health belief models theory²⁰ and the social-ecological model theory^{18,22}. Health belief models are theories that focus on individuals²³, three factors that play a role, namely the readiness of the individual, encouragement

from outside the individual, and how the behavior itself. While social ecological theory, in addition to individuals, this theory focuses on interpersonal relationships, organizations, and policies that play a role in creating behavior²⁴. The subjects in the three articles that used the behavior theory approach were in poor health and different age groups, but two articles were in the adult age group and healthy conditions. The outcome of physical activity that we can see in the two articles that use the theoretical approach results in high physical activity changes from other interventions and looks higher in those that use social ecological theory.

Physical activity in different age groups and health conditions is a limitation of this study because it cannot be compared. A healthy person will have higher physical activity compared to a sick person, as well as the general adult group has higher physical activity than children and the elderly. This limitation can be taken into consideration in future research.

The results of the review of this article also found that the interventions provided included

various methods, namely (1) literacy in the form of booklets and policies, assisting, curriculum (2) face-to-face, namely seminars, FGDs, consulting, classes, and debates (3) technology using WhatsApp, social media, phones, and multimedia, (4) exercise, and; (5) provision of infrastructure. All interventions in the articles use literacy methods, six articles that use face-to-face methods, four articles use technology methods, two articles use physical exercise, and two articles use activity infrastructure. Interventions that use all methods are obtained in one article^{22,25}, with the highest outcome results, namely a change of 2612.15 MET-min/week.

Physical activity is a behavior in the community that is difficult to change, so it requires great effort, not only through health education, but there must be health promotion efforts which include advocacy, assimilation, and community empowerment.

CONCLUSION

Sufficient physical activity behavior is a lifestyle that needs to be implemented in society to live a healthy life. One form of effective and efficient promotion of physical activity is by providing educational programs. Based on our literature review found that comprehensive educational interventions produced high changes in physical activity in healthy participants. The level of change in physical activity in the community needs to take into account the participant's health condition, the next literature may be reviewed enhancing of physical activity among people with similar health conditions and age.

Limitation of Study: In this research, only nine eligible articles were obtained from three database sources and search for relevant articles from other sources. It is necessary to add additional database to obtain more specific educational programs based on similar health conditions and age.

REFERENCES

1. Wendra, Rochmah EN, Fadhurrohman W. The Effect of Physical Activity on the Muscle Strength of Obese People' Backs. *Poltekita J Ilmu Kesehatan*. 2023;17(2):487-492.
2. Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219-229. doi:10.1016/S0140-6736(12)61031-9
3. Strain T, Brage S, Sharp SJ, et al. Use of the prevented fraction for the population to determine deaths averted by existing prevalence of physical activity: a descriptive study. *Lancet Glob Heal*. 2020;8(7):e920-e930. doi:10.1016/S2214-109X(20)30211-4
4. Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. *Lancet Glob Heal*. 2018;6(10):e1077-e1086. doi: [https://doi.org/10.1016/S2214-109X\(18\)30357-7](https://doi.org/10.1016/S2214-109X(18)30357-7)
5. Napitupulu RM, Juwita CP, Manik JWH. Joint complaints in the elderly during the COVID-19. *Int J Community Med Public Heal*. 2022;9(7):2803-2807.
6. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451-1462. doi:10.1136/bjsports-2020-102955
7. Presiden. Instruksi Presiden Republik Indonesia Nomor 1 Tahun 2017 Tentang Gerakan Masyarakat Hidup Sehat.; 2017:1-10. <https://ci.nii.ac.jp/naid/40021243259/>

8. OMS. WHO Guidelines on Physical Activity and Sedentary Behaviour.; 2020. https://apps.who.int/iris/bitstream/handle/10665/325147/WHO-NMH-PND-2019_4-eng.pdf?sequence=1&isAllowed=y%0Ahttp://www.who.int/iris/handle/10665/311664%0Ahttps://apps.who.int/iris/handle/10665/325147
9. Damayanti R, Juwita CP, Besral B, Aras D. Enhancing physical exercise adherence by promoting self-efficacy in elderly with osteoarthritis knee. *J Phys Educ Sport*. 2024;24(4):1009-1015. doi:10.7752/jpes.2024.04115
10. Sterkowicz-Przybycień K, Kulesza M. Physical activity and wellness in Polish older women practicing Pilates and nonexercising women: a cross-sectional analysis. *J Phys Educ Sport*. 2023; 23(9):2435-2444. doi:10.7752/jpes.2023.09280
11. Arnell S, Jerlinder K, Geidne S, Lundqvist LO. Experiences of stakeholder collaboration when promoting participation in physical activity among adolescents with autism spectrum disorder. *Disabil Rehabil*. 2022;44(9):1728-1736. doi:10.1080/09638288.2021.1887944
12. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med*. 2009; 151 (4): 65-94. doi:10.1016/j.jclinepi.2009.06.006
13. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Research Synthesis Methods*. doi:10.1002/jrsm.1411
14. Rakhshani T, Khiyali Z, Masrurpour F, Khani Jeihooni A. Effect of educational intervention on improvement of physical activities of middle-aged women. *BMC Womens Health*. 2021;21(1):1-8. doi:10.1186/s12905-021-01494-z
15. Yarım kaya E, Esentürk OK, İlhan EL, Karasu N. A WhatsApp-delivered intervention to promote physical activity in young children with autism spectrum disorder. *Int J Dev Disabil*. 2022; 68(5):732-743. doi:10.1080/20473869.2021.1887436
16. Morais LDC, Paravidino VB, Mediano MFF, et al. Effectiveness of a school-based randomized controlled trial aimed at increasing physical activity time in adolescents. *Eur J Public Health*. 2021;31(2):367-372. doi:10.1093/eurpub/ckab025
17. Gill M, Roth SE, Chan-Golston AM, et al. Evaluation of an Intervention to Increase Physical Activity in Low-Income, Urban Middle Schools. *J Sch Health*. 2019;89(9):705-714. doi:10.1111/josh.12808
18. Slining M, Wills S, Fair M, et al. LiveWell in early childhood: results from a two-year pilot intervention to improve nutrition and physical activity policies, systems and environments among early childhood education programs in South Carolina. *BMC Public Health*. 2021;21(1):1-9. doi:10.1186/s12889-021-10975-7
19. Aasa B, Berglund L, Michaelson P, Aasa U. Individualized low-load motor control exercises and education versus a high-load lifting exercise and education to improve activity, pain intensity, and physical performance in Patients with Low BACK pain: A Randomized controlled trial. *J Orthop Sports Phys Ther*. 2015;45(2):77-85. doi:10.2519/jospt.2015.5021
20. Khodaveisi M, Azizpour B, Jadidi A,

- Mohammadi Y. Education based on the health belief model to improve the level of physical activity. *Phys Act Nutr.* 2021;25(4):17-23. doi:10.20463/pan.2021.0022
21. Vahedian Shahroodi M, Belin Tavakoly Sany S, Hosseini Khaboshan Z, Esmaeily H, Jafari A, Tajfard M. Effect of a theory-based educational intervention for enhancing nutrition and physical activity among Iranian women: a randomised control trial. *Public Health Nutr.* 2021;24(18): 6046-6057. doi:10.1017/S1368980021002664
 22. Tehrani H, Majlessi F, Shojaeizadeh D, Sadeghi R, Kabootarkhani MH. Applying socioecological model to improve women's physical activity: A randomized control trial. *Iran Red Crescent Med J.* 2016;18(3):1-5. doi:10.5812/ircmj.21072
 23. Marvuglia A, Koppelaar R, Rugani B. The effect of green roofs on the reduction of mortality due to heatwaves: Results from the application of a spatial microsimulation model to four European cities. *Ecol Modell.* 2020;438:109351. doi:https://doi.org/10.1016/j.ecolmodel.2020.109351
 24. Unicef. Brief on the Social Ecological Model.; 2020.
 25. Razi P, Marsofely RL, Marleni WA. Comparison of the Effectiveness of Counseling with Social Media , Animation and Short Films in Improving Perception and Motivation to Do Physical Activity in Adolescents of Health Promotion, Poltekkes Kemenkes Jambi, Jambi, Indonesia Departement of Hea. *Poltekita J Ilmu Kesehat.* 2023;17(3):792-797.

Citra Puspa Juwita, Weeke Budhyanti, Bernadetha Nadeak (2025). The Education Programs Enhancing Physical Activity: Systematic Literature Review, *ijmaes*; 11(1); 2287-2298.

THE EDUCATION PROGRAMS ENHANCING PHYSICAL ACTIVITY: SYSTEMATIC LITERATURE REVIEW

ORIGINALITY REPORT

23%

SIMILARITY INDEX

18%

INTERNET SOURCES

15%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to KPJ International College of
Nursing and Health Science

Student Paper

5%

2

bmcpublichealth.biomedcentral.com

Internet Source

2%

3

zenodo.org

Internet Source

1%

4

www.frontiersin.org

Internet Source

1%

5

Submitted to Herzing University

Student Paper

1%

6

repositorio.unican.es

Internet Source

1%

7

irep.iium.edu.my

Internet Source

1%

8

N. Ammon, S. Mason, J.M. Corkery. "Factors
impacting antiretroviral therapy adherence
among human immunodeficiency virus-
positive adolescents in Sub-Saharan Africa: a
systematic review", Public Health, 2018

Publication

1%

9

bmcmwomenshealth.biomedcentral.com

Internet Source

1%

10

repository.stikim.ac.id

Internet Source

1%

11	efsupit.ro Internet Source	1 %
12	journals.sagepub.com Internet Source	1 %
13	Miranda Westfall, Sarah E. Roth, Monique Gill, Alec M. Chan-Golston, Lindsay N. Rice, Catherine M. Crespi, Michael L. Prelip. "Exploring the Relationship Between MyPlate Knowledge, Perceived Diet Quality, and Healthy Eating Behaviors Among Adolescents", American Journal of Health Promotion, 2020 Publication	1 %
14	www.dovepress.com Internet Source	1 %
15	www.wjgnet.com Internet Source	1 %
16	Barwais, Faisal A, Thomas F Cuddihy, and L Tomson. "Physical activity, sedentary behavior and total wellness changes among sedentary adults: a 4-week randomized controlled trial", Health and Quality of Life Outcomes, 2013. Publication	1 %
17	repository.uki.ac.id Internet Source	1 %
18	ircmj.neoscriber.org Internet Source	<1 %
19	academic.oup.com Internet Source	<1 %
20	www.obesityandenergetics.org Internet Source	<1 %

Submitted to Udayana University

21

Student Paper

<1 %

22

pure.amc.nl

Internet Source

<1 %

23

www.researchgate.net

Internet Source

<1 %

24

Citra Puspa Juwita, Dra. Rita Damayanti, Besral, Djohan Aras. "Self-efficacy resources program for behavior changes", Nursing Management, 2023

Publication

<1 %

25

Debbie Van Biesen, Tine Van Damme, Natalia Morgulec-Adamowicz, Aleksandra Buchholz, Momna Anjum, Séan Healy. "A Systematic Review of Digital Interventions to Promote Physical Activity in People With Intellectual Disabilities and/or Autism", Adapted Physical Activity Quarterly, 2023

Publication

<1 %

26

journals.humankinetics.com

Internet Source

<1 %

27

journals2.ums.ac.id

Internet Source

<1 %

28

www.mdpi.com

Internet Source

<1 %

29

periodicosapm.emnuvens.com.br

Internet Source

<1 %

30

www.sportedu.org.ua

Internet Source

<1 %

31

Junyu Wang, Yuan Yuan, Ting Xie, Ligong Zhang, Hong Xu, Shu-Cheng Lin, Yong Yang, Dong Zhu, Jie Zhuang. "Optimal dose and type

<1 %

of exercise to improve motor symptoms in adults with Parkinson's disease: A network meta-analysis", Journal of Science and Medicine in Sport, 2025

Publication

32

Santos, João Francisco. "Um Projeto para Vila Nova de Cerveira: O Percurso pela Construção em Madeira", Universidade do Porto (Portugal), 2024

Publication

<1 %

33

Hadi Tehrani, Fershteh Majlessi, Davoud Shojaeizadeh, Roya Sadeghi, Marzieh Hasani Kabootarkhani. "Applying Socioecological Model to Improve Women's Physical Activity: A Randomized Control Trial", Iranian Red Crescent Medical Journal, 2016

Publication

<1 %

34

Jérémy Raffin, Marie Fourteau, Kelly Virecoulon Giudici, Yves Rolland, Bruno Vellas, Philipe de Souto Barreto. "Cross-sectional and longitudinal associations between physical activity and intrinsic capacity in healthy older adults from the MAPT study", Archives of Gerontology and Geriatrics, 2025

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On