



← Back to Submissions

1440 / **Marliana et al.** / Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life

Library

Workflow

Publication

Submission

Review

Copyediting

Production

Submission Files

Q Search

▶		8783	Article Submission_Lina Marlina_JPHS.docx	October 9, 2025	Article Text
---	---	------	---	-----------------	--------------

Download All Files

Pre-Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
------	------	------------	---------	--------

No Items



← Back to Submissions

1440 / **Marliana et al.** / Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life

Library

Workflow

Publication

Submission

Review

Copyediting

Production

Round 1

Round 1 Status

Submission accepted.

Notifications

[JPHS] Editor Decision	2025-11-10 05:33 PM UTC
[JPHS] Editor Decision	2026-02-28 01:34 PM UTC
[JPHS] Editor Decision	2026-05-23 01:11 PM UTC

Reviewer's Attachments

Q Search

8923

1440-Article Text-6981-8783-2-20251009.docx

October 17, 2025

Revisions

Q Search

Upload File

▶

9556

1440-7113-8923-5-20251017_Revision.docx

November 19, 2025

Article Text

▶

10323

1440-7113-8923-5-20251017_Revision_Update.docx

December 16, 2025

Article Text

Review Discussions

[Add discussion](#)

Name	From	Last Reply	Replies	Closed
Revisions Required	vikas_gupta 2025-11-25 12:44 PM UTC	lina25 2025-12-16 07:03 AM UTC	1	☐
Missing Revision and Response to Reviewer 2	vikas_gupta 2025-12-26 12:12 PM UTC	-	0	☐
▶ Revision according to the suggest of reviewer 1 and reviewer 2	lina25 2026-01-20 01:57 PM UTC	-	0	☐

The Relationship between Earphone Use Patterns and The Risk of Hearing Loss which Influences The Quality of Life

^{1,2}Lina Marlina*, ¹Sena Pongtuluran, ¹Fransiskus Harf Poluan

Corresponding Author: *lina.marlina@uki.ac.id

¹ Department of Ear Nose and Throat, Faculty of Medicine, Universitas Kristen Indonesia

² General Hospital Universitas Kristen Indonesia, Jakarta, Indonesia

Commented [L1]: The title of this article is quite clear and informative. It accurately describes the focus of the research, which links earphone use patterns to the risk of hearing loss and its impact on quality of life. However, the title could be slightly more concise, for example, "Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life." This would have made the title more concise and easier to understand.

ARTICLE INFO

Article history

Received XX July 2022

Revised XX August 2022

Accepted XX August 2022

ABSTRACT

In today's modern life, most people are constantly exposed to noise every day, potentially disrupting their hearing function. One such factor is the use of hearing devices, namely earphones. Approximately 1.1 billion young adults and adolescents are at risk of experiencing hearing loss in their lifetime. If someone is at risk of hearing loss, it can undoubtedly impact their quality of life. Participants in this study were students at the Christian University of Indonesia in 2019, 2020, and 2021. The purpose of this study was to determine whether risky earphone use patterns are associated with quality of life. Data for this study were collected using a cross-sectional observational analytical methodology with consecutive sampling technique. The study sample was assessed with a questionnaire created using Google Forms. The questionnaire, which is divided into two parts: the Earphone Use Pattern Questionnaire (PPE) and the Hearing Handicap Inventory for Adults (HHIA), to determine the patient's quality of life, was then analyzed using the Chi-Square and Spearman tests. Of the 225 respondents, there were 62.2% risky earphone usage patterns and 37.8% non-risky earphone users. According to the HHIA score findings, 74.2% of people had no disabilities, 9.8% had mild to moderate disabilities, and 16.0% had severe disabilities, with $p = 0.015$ ($p < 0.05$) obtained through the use of the Chi-Square statistical test. There is a relationship between earphone usage patterns and the risk of hearing loss which affects the quality of life of students of

the Faculty of Medicine, Universitas Kristen Indonesia Class of 2019, 2020, and 2021.

Keywords

Earphone use patterns
Hearing loss
Hearing Handicap
Inventory for Adults
Quality of Life

This is an open-access article under the [CC-BY-SA](#) license.



Introduction

The ability to hear is at the heart of all human communication, both with each other and with the world around us [1]. Hearing loss is one of the most common disabilities worldwide, affecting more than 1.5 billion people, according to the World Health Organization (WHO) [2]. Hearing loss is common but often goes unrecognized by patients and doctors, leading to inadequate treatment. The prevalence of clinically significant hearing loss has been found to double every decade [3]. In Southeast Asia, the prevalence of hearing loss is estimated at 27%, which equates to approximately 156 million people in the total population. In contrast, the prevalence of hearing loss in Indonesia nationally is 2.6% of the total population, and this figure is expected to increase over time [4]

In today's modern life, most people are constantly exposed to noise that can potentially disrupt hearing function [5]. One of them is the use of hearing devices, namely earphones, which are one of the causes of hearing loss. WHO estimates that 1.1 billion adolescents and young adults are at risk of hearing loss as a direct result of exposure to loud noises from electronic devices. This risk can be associated with the increased volume of sound produced by these devices [6]. The use of earphones is an important indicator of increased risk, especially noise-induced hearing loss, because earphones can bring the noise source closer to the ear so that the ear will receive much greater exposure [7]. A study by Rumampuk in

Commented [L2]: The introduction of this article provides good context by citing global data on hearing loss and linking it to earphone use. The authors state that approximately 1.1 billion young people are at risk of hearing loss due to exposure to loud noise, particularly from electronic devices such as earphones. They also mention the importance of quality of life and how hearing loss can impact it.

Revision Suggestions:

1. While informative, the introduction could be more focused by reducing some overly technical information about ear anatomy to more directly address the research objectives.
2. The use of earphones could be clarified as a primary risk factor in the context of this study to make it more relevant to the research focus.

Indonesia found that 26.7% of students who use earphones experience hearing loss at some level, with 6.7% experiencing moderate hearing loss [8]. Based on data from WHO, there are around 430 million people or more than 5% of the world's population who have hearing loss that impacts their quality of life and based on data, most people who experience hearing loss are caused by the habit of listening to music at high volumes and other noisy environments for long periods of time [9]. Based on the above background, the researcher wants to know the relationship between earphone usage patterns and the risk of hearing loss, which affects the quality of life of students of the Faculty of Medicine, Indonesian Christian University, Class of 2019-2021.

Literature Review

A. Ear Anatomy

Human ear development begins with the appearance of the otic placode and vestibulocochlear ganglia at three weeks of gestation. This is followed by the appearance of the external canal, which begins to develop from the first branchial cleft in the fourth week, then forms a hole through ectodermal proliferation. By the 28th week, the epithelial core has canalized from medial to lateral, resulting in a fully patent external canal [10].

The ear can be divided into three parts: the outer ear, the middle ear, and the inner ear [11]. The outer ear consists of the auricle (also known as the pinna) and the external auditory canal, as well as the lateral surface of the tympanic membrane, which is part of the outer ear [12]. The middle ear consists of the tympanic membrane and three auditory ossicles: the malleus, incus, and stapes. These three ossicles allow sound waves to travel from the outer ear to the inner ear by connecting the tympanic membrane to the inner ear [13]. The inner ear is a system of canals and a cavity within the petrous part of the temporal bone. The inner ear is formed by the bony labyrinth (osseous labyrinth), which contains the membranous labyrinth. The bony labyrinth consists of three components: the semicircular canals, the vestibule, and the bony cochlea. The membranous labyrinth is located within the bony labyrinth and is a system of interconnected channels lined with epithelium and containing endolymph. The membranous labyrinth consists of the membranous semicircular canals, the utricle, the saccule, the endolymphatic duct, the endolymphatic sac, and the reuniens duct. The cochlear duct contains the organ of Corti, which is the organ of hearing. There are sensory nerve endings in the ampullae of the semicircular canals (crista ampullaris) and in the utricle and saccule (macula sacculi and utriculi) which function as static and kinetic senses.




Figure 2.1 Ear Anatomy [3]

B. Physiology of Hearing

Hearing and balance are functions performed by the human ear. The transduction mechanism in the ear is responsible for detecting and analyzing sound. This mechanism is responsible for converting sound waves into electrochemical impulses. If the anatomy is abnormal, hearing will not function properly even if the nerves are intact. Hearing is the process by which sound vibrations from the surrounding environment are converted into action potentials in the cochlea and auditory nerve. Sound is produced when an object is made to vibrate, such as when a guitar string is plucked. The resulting vibrational pressure is called a sound wave. Therefore, the ear is designed to detect and categorize various sonic qualities, including pitch (the frequency of the sound wave) and loudness (the perceived intensity of the sound). Hertz is the standard system of frequency measurement (Hz, cycles per second). The human ear can generally detect sounds between 1,000 and 4,000 hertz, but younger ears can detect sounds between 20,000 and 200,000 hertz. The decibel (dB) is the unit used to measure the loudness of sound, and the audible range that humans can tolerate on the decibel scale is from 0 to 130 dB (where the sound can become painful). To enter the central nervous system, each of these physical properties must first undergo some kind of transformation. Air vibrations are first converted into vibrations of the tympanic membrane, which is the first step in the process. Once they reach these structures, the vibrations are captured by the middle ear and ossicles. Then, in the inner ear and cochlea, these vibrations are converted into vibrations of fluid, and these fluid vibrations stimulate areas known as the basilar membrane and the organ of Corti. Ultimately, these vibrations are converted into nerve impulses, which are then transmitted to the nervous system [14]

The primary function of the organ of Corti is auditory signal transduction. Sound waves enter the ear through the auditory canal and cause vibrations in the tympanic membrane.

Movement of the tympanic membrane causes subsequent vibrations in the auditory ossicles, the three middle ear bones, which transfer energy to the cochlea through the oval window. As the oval window moves, the waves travel through the perilymph fluid in the scala tympani and then the scala vestibuli of the cochlea. As fluid moves through these structures, the basilar membrane, located between the scala media and scala tympani, shifts against the tectorial membrane, respectively [15]

C. Hearing Loss

Hearing loss is a condition that occurs when the transmission of sound from the outer ear to the brain is disrupted. The disruption can occur at any stage, before, after, or after and before the cochlea. Hearing loss can be conductive, sensorineural, or mixed [16]. Conductive hearing loss is caused by impaired transmission of sound waves to the cochlea, which ultimately causes problems primarily in the outer ear and auditory canal. The problem may originate in the external canal due to obstruction from debris, wax, or foreign objects, or may result from a congenital abnormality such as aural atresia, which causes incomplete development of the outer ear. Perforation of the tympanic membrane can also cause conductive hearing loss. Perforation can be associated with trauma from using cotton swabs to clean the ears, barotrauma from diving, or as a sequela of otitis media [17]. Sensorineural hearing loss (SNHL) is the most common type and accounts for the majority of all hearing loss. Hearing loss resulting from neurological, otologic, or vascular disorders is referred to as SNHL. Sensorineural hearing loss can be caused by damage to the auditory nerve, the vestibulocochlear nerve, or the central auditory processing area of the brain [18].

Noise can damage the sensory hair cells of the inner ear through direct mechanical pressure and intense sound pressure as well as by activation of stress-induced molecular pathways, including the formation of reactive oxygen species and excess calcium. Assessed by the decibel level and duration of exposure this can cause noise-induced hearing loss which can also be known as Noise Induced Hearing Loss (NIHL) [19]. NIHL is hearing loss caused by exposure to sounds that are too strong so that they can damage hearing cells, this kind of hearing loss is often described as "noise-induced". NIHL can be classified into three, namely, Temporary Threshold Shift (TTS), Permanent Threshold Shift (PTS), and Acoustic Trauma. Four factors contribute to the effects of noise, namely, sound level (volume), audio spectrum, distribution of exposure time, cumulative noise exposure over days, weeks or years [20]. The term "transient threshold shift" is used to describe objective changes in hearing acuity that can be measured on audiometry immediately after an episode of loud noise exposure and which return to pre-exposure levels after a few days to two weeks. A temporary threshold shift is subjectively characterized by decreased hearing sensitivity, a feeling of fullness in the ear,

tinnitus (ringing), and the perception of hearing loss. Prolonged or repeated exposure to noise can lead to the death of sensory hair cells and result in permanent hearing loss, referred to as a "permanent threshold shift." Hair cell death may be followed by a gradual loss of spiral ganglion neurons over months or years [21].

D. Earphones

To listen to audio, people usually need a device called earphones, which convert electrical energy into sound waves. Earphones are most often used for media consumption and digital communication via gadgets such as smartphones and computers. [22]. Consumers are increasingly turning to earphones due to advances in audiovisual and communication technology. Earphones are now commonly used to listen to music on mobile phones and other audio devices, thanks to these innovations. People, especially teenagers, have made listening to music through headphones or earphones a regular part of their lives. Most teenagers wear earphones or headphones, typically to listen to music for extended periods of time. They do so even while engaging in other activities, such as traveling, exercising, or even sleeping. Prolonged use can cause continuous noise, which can damage hearing [6].

Recent advances in audiovisual technology have had a significant impact on the creation of modern music players and the hearing aids used to listen to music on these devices. Earphones, headsets, headphones, and other types of hearing aids are just a few of the many types that have been produced, with varying capacities and sound output functions [23]. According to one study, 88.2% of 1,407 children used personal music players, also known as MP3 players, and 27.4% of these children listened to music at the highest possible volume for extended periods without interruption. Another study found that 94.3% of those participating used personal music players daily for between one and three hours over three years. Conversely, earphones are the most common form of amplification for the hearing impaired. This demonstrates the widespread use of earphones among today's youth [23].

E. The Impact of Earphone Use Patterns on Hearing Loss

While earphone use is currently more effective than loudspeakers, it's important to note that earphone use can impact hearing response and chronic noise. Even at the same volume, listening to music through earphones produces a higher overall noise intensity than listening to the same music without earphones. When someone listens to music with earphones, the sound source is physically closer to them than when listening without earphones. Earphones also don't fully distribute sound and don't completely block out external noise, which can contribute to hearing loss. Earphone use patterns can be divided into four components: frequency, duration, noise volume, and length of earphone use. These four components are the primary assessments for determining earphone use patterns, which are then classified into two categories: risky and non-risky earphone use patterns [24].

Hearing loss is one of the effects caused by earphone use, resulting from improper use. Continuous exposure is a risk factor for hearing loss. Approximately 5 to 10% of earphone users are at risk of hearing loss and even permanent hearing loss if they listen to music for more than one hour a day at high volume [23]

Due to the length of earphone use of more than three years and daily use, earphones require attention, as literature indicates that prolonged exposure to loud noise can cause noise-induced hearing loss (NIHL) [25]. Research by Setianiet (2018) found that 77.8% of those with a history of noise exposure from earphones experienced NIHL, and 22.2% experienced NIHL with more than five years of use. Therefore, using earphones for five years or more can increase the risk of NIHL [26]. Noise-induced hearing loss (NIHL) is a form of sensory deficit that can lead to sensorineural hearing loss due to continuous exposure to loud noise over a prolonged period [27].

F. The Impact of Hearing Loss on Quality of Life

According to the World Health Organization (WHOQOL), quality of life is defined as an individual's perception of their life in society within the context of their culture and existing value systems, related to goals, expectations, standards, and concerns. Based on this definition, quality of life can be defined as life satisfaction or the fulfillment of life's needs based on a person's perceived physical, psychological, and social conditions [28], [29]. Factors that can influence a person's quality of life include psychological conditions. For example, psychological conditions can arise from the impact of an individual's physical limitations, which can lead to anxiety disorders, which can reduce the sufferer's ability to work and socialize with others. This can indirectly impact a person's quality of life. Beyond psychological factors, there are social relationship factors such as social support, which is the presence of others who can be relied upon to provide assistance, encouragement, acceptance, and attention, thereby improving and influencing the well-being or quality of life for the individual concerned [30]

Hearing loss can lead to a potential handicap or disability, which can limit the individual's ability to perform various tasks in life. This can have psychological impacts, such as communication barriers, leading to anxiety, depression, and even impaired social interaction. Psychological and social relationships are factors that can impact a person's quality of life [31]

Material And Methods

This study uses a cross-sectional observational analytical research design, the researcher wants to prove the extent of the closeness of the relationship between a variable and another variable, where in this study the researcher wants to know the relationship

Commented [L3]: The method used was a cross-sectional observational study design with consecutive sampling. The authors collected data using two questionnaires: the Earphone Usage Pattern Questionnaire (EUPQ) and the Hearing Handicap Inventory for Adults (HHIA). Data analysis was performed using the Chi-square test and Spearman's correlation test.

Strengths:

1. The study design used was appropriate for identifying the relationship between earphone usage patterns and hearing loss.
2. The selection of two relevant questionnaire instruments (EUPQ and HHIA) supported the study's objective of measuring earphone usage patterns and their impact on quality of life.

Recommendations for Revision:

1. The sample used was students from the Faculty of Medicine, which could be biased because it is not representative of the general population. The authors recommend adding more sample variation to increase the generalizability of the study results.
2. Further explanation of the inclusion and exclusion criteria would improve the transparency of the research method.

between earphone use patterns and the risk of hearing loss which can affect the quality of life of students of the Faculty of Medicine, Indonesian Christian University Class of 2019-2021.

This research was conducted from December to January 2023 at the Faculty of Medicine, Christian University of Indonesia, on Jl. Mayjen Sutoyo No. 2, RT. 2/RW. 11, Cawang Village, Kramat Jati District in East Jakarta City. The population of this study consisted of students from the Faculty of Medicine at Universitas Kristen Indonesia who graduated between the 2019-2021 intake. The sample consisted of all students who met the inclusion and exclusion criteria and had provided informed consent. The sampling technique used in this study was non-probability sampling, namely, consecutive sampling, which was used to determine the research sample based on predetermined criteria and objectives. The research instrument consisted of two questionnaires: the Earphone Usage Pattern Questionnaire (EUPQ), which was used to examine respondents' habits of using earphones, and the Hearing Handicap Inventory for Adults (HHIA) questionnaire, which was used to determine the quality of life of earphone user respondents.

Data collected through the questionnaire were processed using IBM SPSS (Statistics for Social Science) and Microsoft Office Excel. Categorical data were displayed as percentages, while numerical data were displayed as means and standard deviations. All data were presented in tabular form. An inferential analysis was then performed to analyze the relationship between EUPQ and HHIA using the Chi-square test. The Pearson correlation test was used to analyze the relationship between EUPQ scores and HHIA, provided the data were normally distributed; otherwise, the Spearman test was employed.

Results

A. Respondent Profile

This study used an online questionnaire created by Google Forms and administered to students at the Faculty of Medicine, Universitas Kristen Indonesia, between December 2022 and January 2023. The aim of this study was to determine the relationship between earphone use patterns and the risk of hearing loss, which impacts quality of life, as measured by the Hearing Handicap Inventory for Adults (HHIA) questionnaire, adapted into Indonesian.

The study population comprised 225 students from the 2019, 2020, and 2021 intakes at the Faculty of Medicine, Universitas Kristen Indonesia.

The study lasted three months and was divided into two parts. In the first phase, participants were asked to complete a questionnaire about their regular earphone use patterns, and then they were asked to complete the HHIA questionnaire. The data was then analyzed, and a thesis was prepared.

Commented [L4]: The results showed that 62.2% of respondents used earphones in a risky manner, while 37.8% did not. Based on the results of the HHIA questionnaire, the majority of respondents (74.2%) did not experience hearing loss, but 16% of respondents experienced moderate to severe hearing loss. Statistical tests showed a significant relationship between risky earphone use patterns and hearing loss that impacted quality of life ($p = 0.015$).

Strengths:

1. The study results are presented clearly through tables and statistics that support the conclusion that risky earphone use patterns are associated with hearing loss that impacts quality of life.
2. The presentation of data through tables helps readers easily understand the distribution of respondents and the questionnaire results.

Revision Suggestions:

1. Although the data are presented clearly, the presentation of the results could be supplemented with graphs that more visually illustrate the relationship between variables.
2. The author could add more detailed information regarding respondent characteristics, for example, age or gender, which may have influenced the study results.

Table 1: EUPQ Distribution Overview

Assessment Criteria		<i>f</i>	%
Length of Use of Earphones	0-3 years	52	23,1
	>3 years	173	76,9
Duration of Earphone Use	<1 hour	97	43,1
	1-2 hours	66	29,3
	>2-4 hours	36	16,0
	>4-6 hours	14	6,2
	>6-8 hours	5	2,2
	>8 hours	7	3,1
Frequency of Earphone Use	1-2 days	115	51,1
	3-4 days	62	27,6
	5-6 days	17	7,6
	Every Days	31	13,8
Earphone Usage Volume	<60%	140	62,2
	60-80%	69	30,7
	>80%	16	7,1
Earphone Usage Patterns	No Risk	85	37,8
	Risk	140	62,2
Total		225	100,0

Based on Table 1, 52 respondents used earphones for 0-3 years, representing 23.1% of the total, while 173 respondents used earphones for more than 3 years, representing 76.9% of the total. Based on these findings, the majority of students using earphones at the Faculty of Medicine, Universitas Kristen Indonesia (UKI) have used them since their teenage years, with some even using them since junior high school. A total of 97 respondents (43.1%) used earphones for more than 1 hour, 66 respondents (29.3%) used earphones for 1-2 hours, 36 respondents (16.0%) used earphones for more than 2-4 hours, 14 respondents (6.2%) for more than 4-6 hours, 5 respondents (2.2%) for more than 6-8 hours, and 7 respondents (3.1%) for more than 8 hours. From this table, it can be seen that students of the Faculty of Medicine, Christian University of Indonesia, Class of 2019, 2020, and 2021 mostly use earphones for more than one hour per day.

In total, 115 respondents (51.1%) reported using earphones 1-2 days a week, while 62 (27.6%) used them 3-4 days a week, 17 (7.6%) used them 5-6 days a week, and 31 (13.8%) used them daily. Here, we see that first-, second-, and third-year medical students at the Christian University of Indonesia used earphones between two and three days a week. 140 (62.2%) respondents used earphones with a volume of <60% (≤ 9 clicks of 0 volume), 69 (30.7%) used earphones with a volume of 60-80% (10-12 clicks of 0 volume), and 16 (7.1%) used earphones with a volume of >80% (> 12 clicks of 0 volume). In this case, students at the Faculty of Medicine, Indonesian Christian University, Class of 2019, 2020, and 2021 mostly used earphones with a volume of <60% (≤ 9 clicks from 0% volume). 85 respondents (37.8%) had earphone usage patterns that did not pose a risk, while 140 (62.2%) had earphone usage patterns that did.

Title: Paper Formatting for IISTR (max. 12 words) (First author, et al.)

From what we can tell from this study, the majority of medical students at the Christian University of Indonesia (UI) in the 2019, 2020, and 2021 intakes are at risk of hearing loss. Heavy, long-term earphone use (>3 years), daily use (>1 hour), weekly use (>4 days), and high volume use (>60%) all contribute to risky earphone use.

Table 2: HHIA Score Results Overview

Indicator		f	%
Hearing Handicap Inventory Adults Score Results	No Handicap	167	74,2
	Mild Moderate Handicap	22	9,8
	Severe Handicap	36	16
	Total	225	100,0

Table 2 shows the scores from the Hearing Handicap Inventory for Adults (HHIA) questionnaire distributed to groups at risk for hearing loss. Of the respondents who completed the questionnaire, 167 (74.2%) had no handicap, 22 (9.8%) had a mild-moderate handicap, and 36 (16%) had a severe handicap.

Table 3: The relationship between EUPQ and HHIA

Earphone Usage Patterns	Hearing Handicap Inventory Adults (HHIA)						p-value
	No Handicap		Mild Moderate Handicap		Severe Handicap		
	<i>f</i>	%	<i>f</i>	%	<i>f</i>	%	
No Risk	71	83,5	8	9,4	6	7,1	0,015
Risk	96	68,6	14	10,0	30	21,4	
Total	167	74.2	22	9.8	36	16	

Table 3 shows that 96 respondents (68.6%) had earphone use patterns that put them at risk for hearing loss, 14 (10.0%) had mild-moderate handicaps, and 30 (21.4%) had severe handicaps. Meanwhile, 71 respondents (83.5%) were not at risk for hearing loss, 8 (9.4%) had mild-moderate handicaps, and 6 (7.1%) had severe handicaps. Statistical tests revealed a statistically significant relationship between risky and non-risky earphone use and handicaps that can affect respondents' quality of life, with a p-value of 0.015 ($p < 0.05$). Therefore, it can be concluded that there is a significant relationship between earphone use patterns and the risk of hearing loss. This relationship affects a person's quality of life. To determine the significance of the correlation and the degree of correlation between the independent and dependent variables, an analysis was conducted based on numerical data from the earphone usage pattern questionnaire and HHIA scores using the Pearson correlation test if the data were normally distributed and alternatively using the Spearmann correlation test if the data were not normally distributed. A normality test was conducted to determine the normality of the data distribution, using the Kolmogorov-Smirnov test because the sample size was >50.

Table 4: Normality Test Results

Variable	Mean	SD	Median	Min	Max	p-value
Earphone Usage Patterns	9,48	3,24	9,0	4,0	18,0	<0,001^a
HHIA	16,14	21,42	6,0	2,0	96.0	<0,001

Description:

^aKolmogorov-Smirnov

Table 4 shows the results of the Kolmogorov-Smirnov test on earphone usage patterns, with a p value of <0.05 and HHIA of <0.05.

Table 5: Spearman Test Results

Variable	r	P-value
EUPQ		
HHIA	0,202	0,002 ^b

Description:

^bSpearman's Test

A Spearman's correlation test followed this. Table 4.5 shows a weak, significant relationship between earphone use patterns and the risk of hearing loss, which impacts students' quality of life.

B. Discussion

1. Distribution of Respondents' Earphone Usage Patterns

From the results of the research conducted by researchers, a questionnaire was completed to assess earphone usage patterns, which included information on duration, frequency, duration, and volume. It was found that the majority of respondents (173 respondents) had used earphones for more than three years, totaling 76.9%. This indicates that students from the Faculty of Medicine, Christian University of Indonesia (2019-2021) have been listening to music through earphones since they were in school. This aligns with research conducted by Fatimah Hamzah and Nurul, whose respondents were students from the Faculty of Medicine, Muhammadiyah University of Makassar, with an average duration of earphone use of more than three years (95.7%) [32]. According to the ENT-KL textbook, hearing loss due to prolonged exposure to excessively loud noise can lead to sensorineural hearing loss [33]. According to the World Health Organization (WHO), the largest number of earphone users worldwide are adolescents aged 12 to 24, who constitute part of the urban population [34]. Relationship between Sitting Position and Lower Back Pain.

Commented [L5]: This article's discussion links the research findings to previous studies, highlighting the importance of earphone usage patterns in increasing the risk of hearing loss. The author also explains how factors such as the duration and volume of earphone use contribute to the risk of hearing loss. This discussion emphasizes the importance of awareness of the negative impacts of unwise earphone use.

Strengths:

1. A fairly robust discussion, linking the research findings to existing literature, provides readers with a better understanding of the topic.
2. The author also cites several other studies that support the findings of this study, enriching the discussion.

Revision Suggestions:

1. The discussion could be more in-depth by further exploring the biological mechanisms involved in listening to music at high volume through earphones and how this affects hearing in the long term.
2. A discussion of other factors influencing quality of life, such as social support or mental health, which may interact with the results of this study could also be added.

The use of earphones has become commonplace in modern culture, especially among young people, who are more likely to use them as part of their daily routine than older generations.²⁸ Earphones can be used as a means of listening to music and communicating because they can produce clearer sound than listening through the earphone speakers themselves. Furthermore, earphones are easy to carry anywhere, the volume can be adjusted, and they can provide privacy for the user. Regarding the duration of earphone use, the respondents in this study mostly used earphones for less than 1 hour, amounting to 97 people (47.1%). This result is in line with research conducted by Martanegara, Ilman, it was found that most respondents in the study used earphones for <1 hour amounting to 61.34%. Noise-induced hearing loss can occur in the general public, especially teenagers, if they are often exposed to the use of earphones at high volume and continue to use them for an average of one to three hours per day [26]. More frequent use of earphones can be caused by the attachment to using earphones at high intensity because the use of earphones can be used in all things such as when driving, sitting relaxing, doing activities, this earphone is very helpful so that even though doing activities, the person can still communicate or listen to music [22].

This study also found that 51.5% of the sample, or 115 students, from the Faculty of Medicine, Universitas Kristen Indonesia, Class of 2019, 2020, and 2021, mostly used earphones once or twice a week. A low frequency of earphone use can be determined when respondents wear their earphones from 1-2 days per week. This finding is consistent with that obtained from a study conducted in the past by Fatimah Hamzah and Nurul on students of the Faculty of Medicine, Universitas Muhammadiyah Makassar. The researchers found that almost half of the students (47.8%) used earphones for 1-2 days per week.

Based on the results of this study, as many as 140 respondents (62.2% of the total) often use earphone volume with > 60% on earphones. This is different when compared to research by Velaro, Adrian on Medical Faculty Students, University of North Sumatra, most of the respondents in their study used earphones with a volume of 60-80%.²⁶ Respondents who use earphones with high volume usually occur because users are not satisfied listening to them at low volume, so they choose to increase the volume of the sound produced by the earphones even though they are aware of the dangers of using earphones with high volume. This occurs even though respondents are aware of the dangers of using earphones with high volume [35]. If the use of earphones with high volume has become a habit for its users, there will be a trauma effect on the sound receptors due to the use of earphones with high volume intensity and cause a decrease in hearing function.

2. HHIA Score Overview

Information about hearing loss, expectations, and motivation can be collected using the Hearing Handicap Inventory for Adults (HHIA). The World Health Organization (WHO) defines

disability or handicap as "a disadvantage to a person due to an impairment or impairment that limits or prevents the fulfillment of a normal role," taking into account the person's age, gender, and social and cultural context. The HHIA is a questionnaire used to assess the extent of handicap in adults with hearing loss. The HHIA is just one of many self-report instruments available to measure the functional impact of hearing loss on a person's daily life. HHIA results can be used to address some of the psychological impacts of hearing loss. The higher the HHIA score, the greater the recognition that hearing loss is a significant disability for the respondent due to the greater limitations or obstacles experienced by the sufferer in fulfilling their daily roles, resulting in a diminished quality of life.

This tool can assist audiologists by providing additional information regarding: (1) helping to substantiate complaints of hearing loss that are not apparent on conventional audiometric testing results; (2) facilitating candidate decisions regarding the use of amplification (hearing aids); (3) assisting in the counseling process; (4) serving as a guide for determining client-centered rehabilitation programs; and (5) serving as a measurement criterion in rehabilitation evaluations, including the benefits of post-surgical hearing aids.⁴⁶ Hearing loss can impact quality of life and limit activities or participation in daily life. The impact of this hearing loss can be assessed using the HHIA questionnaire [36]. In this study, the researchers used the HHIA questionnaire, which had been adapted into Indonesian and had been tested for validity and reliability.

The HHIA questionnaire in this study was administered to groups of people at and not at risk of developing hearing loss due to earphone use. 225 people (100%) completed the HHIA questionnaire. The results showed that 74.2% had no handicap, 9.8% had mild-moderate handicap, and 16% had severe handicap. In a previous study by Marcus and Dayse, the majority of respondents (70.8%) had severe handicap, 22.9% had mild-moderate handicap, and 6.3% had no handicap [37]. The HHIA questionnaire varies widely and depends on life experiences, health expectations, and even the individual's ability to adapt. People with hearing loss will experience communication difficulties and experience social and emotional difficulties in their daily lives. This will lead to different perceptions of their quality of life [36].

3. Relationship between Earphone Use Patterns and HHIA Scores

This study found a statistically significant relationship between earphone use patterns and the risk of hearing loss, which impacts the quality of life of students at the Faculty of Medicine, Christian University of Indonesia, in the 2019, 2020, and 2021 intakes ($p = 0.015$) of the 2019, 2020, and 2021 intakes ($p < 0.05$). Furthermore, the Spearman correlation test revealed a weakly significant relationship between earphone use patterns and the risk of

hearing loss, which impacted students' quality of life ($p = 0.002b$). Similarly, research by Marcus and Dayse confirmed that hearing loss can significantly impact quality of life [37].

The correlation between the two variables supports this finding. Incorrect use of earphones can lead to a risk of hearing loss and negatively impact quality of life, as measured by the HHIA questionnaire. Earphone use patterns can increase the risk of hearing loss, which can lead to physical, social, and emotional changes in individuals. These changes can persist and disrupt daily activities, significantly impacting quality of life, especially for those at risk for hearing loss.

Conclusion

This study shows that risky earphone usage patterns are significantly related to the quality of life of students from the Faculty of Medicine, Christian University of Indonesia, class of 2019, 2020, and 2021. Of the 225 respondents, the majority (62.2%) used earphones with risky patterns, and the results of the Chi-Square statistical test ($p = 0.015$; $p < 0.05$) confirmed a significant relationship between earphone usage patterns and hearing loss that affects quality of life. These findings indicate that unwise earphone use can increase the risk of hearing loss, which has an impact on the social and emotional aspects of students. Therefore, preventive efforts are needed in the form of education on how to use earphones safely and increasing awareness of the importance of maintaining hearing health from a young age.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- [1] C. J. Plack, *The sense of hearing*. Taylor and Francis, 2023. doi: 10.4324/9781003303329/SENSE-HEARING-CHRISTOPHER-PLACK.
- [2] W. H. Organization, *World report on hearing*. 2021. Accessed: Oct. 07, 2025. [Online]. Available: https://books.google.com/books?hl=en&lr=&id=zMRqEAAQBAJ&oi=fnd&pg=PR5&dq=WHO.+World+Report+On+Hearing.+2021&ots=cHAtI_6aMx&sig=5ks-DjL_goiQWico-iRullwvc-8
- [3] C. L. Nieman and E. S. Oh, "Hearing loss," *acpjournals.org* CL Nieman, *ES OhAnnals Intern. Med.* 2020•*acpjournals.org*, vol. 173, no. 11, pp. ITC81-ITC96, Dec. 2020, doi: 10.7326/AITC202012010.
- [4] B. Putri, R. Halim, H. N.-J. Kesmas, and undefined 2021, "Studi kualitatif gangguan pendengaran akibat bising/noise induced hearing loss (NIHL) pada marshaller di Bandar Udara Sultan Thaha Kota Jambi tahun 2020," *Mail. Putri, R Halim, HS NasutionJurnal Kesmas Jambi, 2021*•*mail.online-journal.unja.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://mail.online-journal.unja.ac.id/jkmj/article/view/12400>
- [5] N. Sarah, ... F. L.-J. (Jurnal K., and undefined 2016, "Hubungan Penggunaan Earphone Dengan Gangguan Pendengaran Pada Siswa Sma Negeri 9 Manado," *ejournal.unsrat.ac.id* NA Sarah, *F Lintong, JF RumampukJJK (Jurnal Kedokt. Klin. 2016*•*ejournal.unsrat.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.unsrat.ac.id/index.php/jkk/article/view/14363>
- [6] A. Budiarmanto, A. Asnawati, H. Huldani, ... S. K.-, and undefined 2024, "GAMBARAN TINGKAT PENGETAHUAN MAHASISWA PSKPS FK ULM ANGKATAN 2020 TENTANG FAKTOR RISIKO NOISE-INDUCED HEARING LOSS (NIHL)," *ppjp.ulm.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ppjp.ulm.ac.id/journals/index.php/hms/article/view/13178>

Commented [L6]: The conclusion could be stronger by adding further recommendations regarding policies or interventions that can be implemented by educational institutions or related agencies to reduce the risk of hearing loss due to unsafe earphone use.

- [7] P. Berliana, S. Dan, S. Keman, D. Kesehatan, L. Fakultas, and K. Masyarakat, "Pengaruh Penggunaan Pelindung Telinga dan Earphone Terhadap Noise Induced Hearing Loss Dan Tinitus Pada. Pekerja Bengkel," *e-journal.unair.ac.idPB Syah, S KemanJurnal Kesehat. Lingkungan, 2017•e-journal.unair.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://e-journal.unair.ac.id/JKL/article/download/9148/5147>
- [8] "Susiyanti E, Imanto M. Efek Pengunaan Earphone sebagai... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Susiyanti+E%2C+Imanto+M.+Ef+ek+Pengunaan+Earphone+sebagai+Faktor+Resiko+Kejadian+Noise+Induced+Hearing+Loss.+Majority.+2020%3B+9%282%29%3A+63-67.+&btnG= (accessed Oct. 07, 2025).
- [9] E. Yuliyani, D. Yudhanto, ... W. A.-I. J., and undefined 2022, "Peranan program acara stasiun TVRI NTB sebagai salah satu media edukasi kesehatan pendengaran," *Acad. Yuliyani, D Yudhanto, WS Aff. E Triani, F Libr. DAIS PrajnyaswariINDRA J. Pengabdi. Kpd. Masyarakat, 2022•academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/93056133/125.pdf>
- [10] L. Moneta, L. Q.-D.-O.-H. and N. Surgery, and undefined 2017, "Embryology and anatomy of the ear," *Elsevier*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S1043181017300398>
- [11] R. Maynard and N. Downes, *Anatomy and histology of the laboratory rat in toxicology and biomedical research*. 2019. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=2kGHDwAAQBAJ&oi=fnd&pg=PP1&dq=Maynard+RL,+Downes+N.+The+Ear.+In:+Anatomy+and+Histology+of+the+Laboratory+Rat+in+Toxicology+and+Biomedical+Research+%5BInternet%5D.+Elsevier%3B+2019.+p.+293-302.+Available+from:+https://linkinghub.elsevier.com/retrieve/pii/B978012811837500023X&ots=-lqlh67SG&sig=eYuR9X9RkEvkTDRjfkG-C7whl70>
- [12] A. Szymanski and Z. Geiger, "Anatomy, head and neck, ear," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/article/nbk/nbk470359>
- [13] A. Szymanski, J. Toth, M. Ogorevc, and Z. Geiger, "Anatomy, head and neck, ear tympanic membrane," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk448117>
- [14] "Sánchez López de Nava A, Lasrado S. Physiology, Ear. 2022 - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Sánchez+López+de+Nava+A%2C+Lasrado+S.+Physiology%2C+Ear.+2022&btnG= (accessed Oct. 07, 2025).
- [15] H. White, M. Helwany, ... A. B.-S., and undefined 2023, "Anatomy, head and neck, ear organ of corti," *ncbi.nlm.nih.govHJ White, M Helwany, AR Biknevičius, DC PetersonStatPearls [Internet], 2023•ncbi.nlm.nih.gov*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.ncbi.nlm.nih.gov/books/NBK538335/>
- [16] R. Sataloff and P. Roehm, *Hearing loss*. 2024. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=lg0iEQAAQBAJ&oi=fnd&pg=PP1&dq=+Hearing+Loss&ots=Y1HvwBOzEA&sig=EjaY-3-SDMk5upL0768pSMieSHE>
- [17] T. Sooriyamoorthy and O. De Jesus, "Conductive hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk563267>
- [18] R. Tanna, J. Lin, and O. De Jesus, "Sensorineural hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk565860>
- [19] S. C.-C. E. and H. Health and undefined 2019, "Noise-induced hearing loss," *account.cehh.press.lshrm.ac.ukS ChadhaCommunity Ear Hear. Heal. 2019•account.cehh.press.lshrm.ac.uk*, Accessed: Oct. 07, 2025. [Online]. Available: <https://account.cehh.press.lshrm.ac.uk/index.php/s-j-cehh/article/view/31>
- [20] "Chan Y, Goddard JC. Preface. In: KJ Lee's Essential... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Chan+Y%2C+Goddard+JC.+Pref+ace.+In%3A+KJ+Lee's+Essential+Otolaryngology%3A+Head+and+Neck+Surgery%2C+12e+%5BInternet%5D.+New+York%2C+NY%3A+McGraw-Hill+Education%3B+2019.+Available+from%3A+accesssurgery.mhmedical.com%2Fcontent.aspx%3Faid%3D1172366487&btnG= (accessed Oct. 07, 2025).
- [21] L. Cunningham, D. T.-N. E. J. of Medicine, and undefined 2017, "Hearing loss in adults," *Mass Med. SocLL Cunningham, DL TucciNew Engl. J. Med. 2017•Mass Med. Soc*, vol. 377, no. 25, pp. 2465–2473, Dec. 2017, doi: 10.1056/NEJMRA1616601.
- [22] M. Erlanda, P. Negara, I. Triansyah, D. Hasni, and B. Y. Febrianto, "Hubungan Intensitas Penggunaan Earphone dengan Derajat Gangguan Pendengaran pada Mahasiswa Fakultas

- Kedokteran Universitas Baiturrahmah," *journal.scientific.id*MEP Negara, I Triansyah, D Hasni, BY FebriantoScientific Journal, 2022•*journal.scientific.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://journal.scientific.id/index.php/sciena/article/view/49>
- [23] A. Velaro, D. Z.-O. R. L. Indonesiana, and undefined 2021, "The correlation between earphone usage patterns with the incidence rate and severity of tinnitus," *orli.or.id*AJ Velaro, D Zahar. *Rhino Laryngol. Indones.* 2021•*orli.or.id*, Accessed: Oct. 07, 2025. [Online]. Available: <http://orli.or.id/index.php/orli/article/view/432>
- [24] M. Zhang, A. Jeske, S. Y. (yrs), and undefined 2017, "Changes in the frequency and intensity of Tinnitus using the Suppressive Noise Spectrum," *m.karmaimpex.com*M Zhang, A Jeske, S Youngage (yrs), 2017•*m.karmaimpex.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://m.karmaimpex.com/hjed/pdf/ated-aid1002.pdf>
- [25] B. Rosita, "Pemakaian Earphone dalam Mendengar Musik yang Menimbulkan Bising di Kalangan Milenial," 2019, Accessed: Oct. 07, 2025. [Online]. Available: <https://osf.io/asmb.v1/>
- [26] L. Setiani, N. Syakila, Y. Y.-J. K. N. Medika, and undefined 2018, "Hubungan Lama Paparan Penggunaan Earphone Musik Terhadap Terjadinya Gangguan Pendengaran Akibat Bising pada Mahasiswa Fakultas Kedokteran," *jknamed.com*L Setiani, N Syakila, Y YusniJurnal Kedokt. Nanggroe Med. 2018•*jknamed.com*, Accessed: Oct. 07, 2025. [Online]. Available: <http://www.jknamed.com/jknamed/article/view/51>
- [27] T. Wang, T. Chang, R. Tyler, Y. Lin, W. L.-B. sciences, and undefined 2020, "Noise induced hearing loss and tinnitus—new research developments and remaining gaps in disease assessment, treatment, and prevention," *mdpi.com*TC Wang, TY Chang. R Tyler, YJ Lin, WM Liang, YW Shau, WY Lin, YW Chen, CD LinBrain Sci. 2020•*mdpi.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.mdpi.com/2076-3425/10/10/732>
- [28] "Pratiwi Y. Pengaruh Dukungan Sosial Terhadap Kualitas... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Pratiwi+Y.+Pengaruh+Dukungan+Sosial+Terhadap+Kualitas+Hidup+Lanjut+Usia+di+Pusar+Santunan+Keluarga+%28PUSAKA%29+Kecamatan+Pancoran+Jakarta+Selatan+%5Bskripsi%5D.+Jakarta%3A+Fakultas+Dakwah+dan+Komunikasi+Universitas+Islam+Negeri+Syarif+Hidayatullah%3B+2017&btnG= (accessed Oct. 07, 2025).
- [29] "WHO. WHOQOL User Manual Programme On Mental Health... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=WHO.+WHOQOL+User+Manual+Programme+On+Mental+Health+Division+Of+Mental+Health+And+Prevention+Of+Substance+Abuse+World+Health+Organization.+2020&btnG= (accessed Oct. 07, 2025).
- [30] D. Maya *et al.*, "Kualitas hidup lansia ditinjau dari sabar dan dukungan sosial," *ejournal.umm.ac.id*DMP Sari, CYD Lestari, EC Putra, F Nashorijurnal Ilm. Psikol. Ter. 2018•*ejournal.umm.ac.id*, vol. 06, no. 02, pp. 2540–8291, 2018, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.umm.ac.id/index.php/jipt/article/view/5341>
- [31] A. Nurfaizi, "MENINGKATKAN KEMAMPUAN DETEKSI DINI DAN PENATALAKSANAAN GANGGUAN PENDENGARAN DI KALIBARU KULON BANYUWANGI," *academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/108530358/14428.pdf>
- [32] "Hamzah NF. Pengaruh Perilaku Penggunaan Earphone... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Hamzah+NF.+Pengaruh+Perilaku+Penggunaan+Earphone+Terhadap+Gangguan+Telinga+Pada+Mahasiswa+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar+%5Bskripsi%5D.+Makassar%3A+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar%3B+2021&btnG= (accessed Oct. 07, 2025).
- [33] "Soepardi EA, Iskandar N, Bashiruddin J, Restuti RD.... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Soepardi+EA%2C+Iskandar+N%2C+Bashiruddin+J%2C+Restuti+RD.+Buku+Ajar+Ilmu+Kesehatan%3A+Telinga%2C+Hidung%2C+Tenggorok%2C+Kepala+%26+Leher.+7th+ed.+Jakarta%3A+BP+FKUI%3B+2012.+&btnG= (accessed Oct. 07, 2025).
- [34] N. Yudiasari, S. W.-P. J. K. Masyarakat, and undefined 2021, "Pengembangan Media Promosi Kesehatan Tentang Bahaya Penggunaan Earphone Mahasiswa Universitas Airlangga Di Surabaya," *103.245.72.38*N Yudiasari, S WidatiPreventif J. Kesehat. Masyarakat, 2021•*103.245.72.38*, vol. 12, pp. 57–86, 2021, Accessed: Oct. 07, 2025. [Online]. Available: <http://103.245.72.38/index.php/preventif/article/view/182>
- [35] S. Levey, B. Fligor, ... C. G.-C. I. in, and undefined 2012, "The effects of noise-induced hearing loss on children and young adults," *pubs.asha.org*S Levey, BJ Fligor, C Ginocchi, L KagimbiContemporary Issues Commun. Sci. Disord. 2012•*pubs.asha.org*, vol. 39, no. Fall, pp. 76–83, Oct. 2012, doi: 10.1044/CICSD_39_F_76.

- [36] J. Bashiruddin, W. Alviandi, B. Bramantyo, and R. Ismail, "Praktik Klinis: Hearing Handicap Inventory For Adults Adaptasi Bahasa Indonesia Sebagai Alat Ukur Kualitas Hidup Pasien Gangguan Pendengaran," 2019, Accessed: Oct. 07, 2025. [Online]. Available: <https://scholar.ui.ac.id/en/publications/buku-panduanpetunjuk-panduan-praktik-klinis-hearing-handicap-inve>
- [37] M. Voola, D. T.-V.-T. M. Journal, and undefined 2020, "Quality of life handicap measured in patients with profound unilateral or bilateral deafness," *tasmanmedicaljournal.com* M Voola, D Távora-Viera *Tasman Med. Journal*, 2020 • *tasmanmedicaljournal.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://tasmanmedicaljournal.com/2020/12/quality-of-life-handicap-measured-in-patients-with-profound-unilateral-or-bilateral-deafness/>

Authors



Lina Marlina    is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. She is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery. She is a member of the Indonesian Medical Association (IDI). Her professional interests include ENT and ENT and clinical research. She is actively involved in academic research and public health programs. She can be contacted at lina.marlina@uki.ac.id.



Sena Pongtuluran    is a general practitioner who recently graduated from the Faculty of Medicine, Christian University of Indonesia, Jakarta. She is currently pursuing a Master's degree in Hospital Administration (MARS) at Pelita Harapan University (UPH). Her interests include clinical medicine, community health, and healthcare management. She is passionate about contributing to healthcare innovation and evidence-based medical practice. She can be contacted at senapongtuluran@gmail.com



Fransiskus Harf Poluan    is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. He is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery at Mayapada Hospital, Kuningan, Jakarta. He is actively involved in academic research and public health programs. He can be contacted at fransiskus.poluan@uki.ac.id

Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life

^{1,2}Lina Marlina*, ¹Sena Pongtuluran, ¹Fransiskus Harf Poluan

Corresponding Author: *lina.marlina@uki.ac.id

¹ Department of Ear Nose and Throat, Faculty of Medicine, Universitas Kristen Indonesia

² General Hospital Universitas Kristen Indonesia, Jakarta, Indonesia

ARTICLE INFO

Article history

Received XX July 2022

Revised XX August 2022

Accepted XX August 2022

ABSTRACT

In today's modern life, most people are constantly exposed to noise every day, potentially disrupting their hearing function. One such factor is the use of hearing devices, namely earphones. Approximately 1.1 billion young adults and adolescents are at risk of experiencing hearing loss in their lifetime. If someone is at risk of hearing loss, it can undoubtedly impact their quality of life. Participants in this study were students at the Christian University of Indonesia in 2019, 2020, and 2021. The purpose of this study was to determine whether risky earphone use patterns are associated with quality of life. Data for this study were collected using a cross-sectional observational analytical methodology with consecutive sampling technique. The study sample was assessed with a questionnaire created using Google Forms. The questionnaire, which is divided into two parts: the Earphone Use Pattern Questionnaire (PPE) and the Hearing Handicap Inventory for Adults (HHIA), to determine the patient's quality of life, was then analyzed using the Chi-Square and Spearman tests. Of the 225 respondents, there were 62.2% risky earphone usage patterns and 37.8% non-risky earphone users. According to the HHIA score findings, 74.2% of people had no disabilities, 9.8% had mild to moderate disabilities, and 16.0% had severe disabilities, with $p = 0.015$ ($p < 0.05$) obtained through the use of the Chi-Square statistical test. There is a relationship between earphone usage patterns and the risk of hearing loss which affects the quality of life of students of the Faculty of Medicine, Universitas Kristen Indonesia Class of 2019, 2020, and 2021.

Keywords

Earphone use patterns
Hearing loss
Hearing Handicap
Inventory for Adults
Quality of Life

This is an open-access article under the [CC-BY-SA](#) license.



Introduction

The ability to hear is at the heart of all human communication, both with each other and with the world around us [1]. Hearing loss is one of the most common disabilities worldwide, affecting more than 1.5 billion people, according to the World Health Organization (WHO) [2]. While common, hearing loss often goes unrecognized by patients and doctors, leading to inadequate treatment. The prevalence of clinically significant hearing loss has been found to double every decade. [3]. In Southeast Asia, the prevalence of hearing loss is estimated at 27%, which equates to approximately 156 million people in the total population. In contrast, the prevalence of hearing loss in Indonesia nationally is 2.6% of the total population, and this figure is expected to increase over time. [4]

In today's modern life, most people are constantly exposed to noise that can potentially disrupt hearing function [5]. The use of personal audio devices, specifically earphones, is a significant and growing risk factor for hearing loss in the context of this study. WHO estimates that 1.1 billion adolescents and young adults are at risk of hearing loss as a direct result of exposure to loud noises from electronic devices. This risk is primarily associated with the increased volume of sound produced by these devices [6]. The use of earphones is a significant indicator of increased risk, especially for noise-induced hearing loss (NIHL), as earphones can bring the noise source closer to the ear, resulting in the listener receiving significantly greater

exposure [7]. A study by Rumampuk in Indonesia found that 26.7% of students who use earphones experience some level of hearing loss, with 6.7% experiencing moderate hearing loss [8]. According to data from the WHO, approximately 430 million people, or more than 5% of the global population, experience hearing loss that affects their quality of life. The habit of listening to music at high volumes via earphones and other noisy environments for long periods of time is a primary cause [9]. Based on the above background, the researcher wants to know the relationship between earphone usage patterns (including volume and duration) and the risk of hearing loss which affects the quality of life of students of the Faculty of Medicine, Indonesian Christian University, Class of 2019-2021.

Literature Review

A. Ear Anatomy

Human ear development begins with the appearance of the otic placode and vestibulocochlear ganglia at three weeks of gestation. This is followed by the appearance of the external canal, which begins to develop from the first branchial cleft in the fourth week, then forms a hole through ectodermal proliferation. By the 28th week, the epithelial core has canalized from medial to lateral, resulting in a fully patent external canal [10].

The ear can be divided into three parts: the outer ear, the middle ear, and the inner ear [11]. The outer ear consists of the auricle (also known as the pinna) and the external auditory canal, as well as the lateral surface of the tympanic membrane, which is part of the outer ear [12]. The middle ear consists of the tympanic membrane and three auditory ossicles: the malleus, incus, and stapes. These three ossicles allow sound waves to travel from the outer ear to the inner ear by connecting the tympanic membrane to the inner ear [13]. The inner ear is a system of canals and a cavity within the petrous part of the temporal bone. The inner ear is formed by the bony labyrinth (osseous labyrinth), which contains the membranous labyrinth. The bony labyrinth consists of three components: the semicircular canals, the vestibule, and the bony cochlea. The membranous labyrinth is located within the bony labyrinth and is a system of interconnected channels lined with epithelium and containing endolymph. The membranous labyrinth consists of the membranous semicircular canals, the utricle, the saccule, the endolymphatic duct, the endolymphatic sac, and the reuniens duct. The cochlear duct contains the organ of Corti, which is the organ of hearing. There are sensory nerve endings in the ampullae of the semicircular canals (crista ampullaris) and in the utricle and saccule (macula sacculi and utriculi) which function as static and kinetic senses.




Figure 2.1 Ear Anatomy [3]

B. Physiology of Hearing

Hearing and balance are functions performed by the human ear. The transduction mechanism in the ear is responsible for detecting and analyzing sound. This mechanism is responsible for converting sound waves into electrochemical impulses. If the anatomy is abnormal, hearing will not function properly even if the nerves are intact. Hearing is the process by which sound vibrations from the surrounding environment are converted into action potentials in the cochlea and auditory nerve. Sound is produced when an object is made to vibrate, such as when a guitar string is plucked. The resulting vibrational pressure is called a sound wave. Therefore, the ear is designed to detect and categorize various sonic qualities, including pitch (the frequency of the sound wave) and loudness (the perceived intensity of the sound). Hertz is the standard system of frequency measurement (Hz, cycles per second). The human ear can generally detect sounds between 1,000 and 4,000 hertz, but younger ears can detect sounds between 20,000 and 200,000 hertz. The decibel (dB) is the unit used to measure the loudness of sound, and the audible range that humans can tolerate on the decibel scale is from 0 to 130 dB (where the sound can become painful). To enter the central nervous system, each of these physical properties must first undergo some kind of transformation. Air vibrations are first converted into vibrations of the tympanic membrane, which is the first step in the process. Once they reach these structures, the vibrations are captured by the middle ear and ossicles. Then, in the inner ear and cochlea, these vibrations are converted into vibrations of fluid, and these fluid vibrations stimulate areas known as the basilar membrane and the organ of Corti. Ultimately, these vibrations are converted into nerve impulses, which are then transmitted to the nervous system [14]

The primary function of the organ of Corti is auditory signal transduction. Sound waves enter the ear through the auditory canal and cause vibrations in the tympanic membrane.

Movement of the tympanic membrane causes subsequent vibrations in the auditory ossicles, the three middle ear bones, which transfer energy to the cochlea through the oval window. As the oval window moves, the waves travel through the perilymph fluid in the scala tympani and then the scala vestibuli of the cochlea. As fluid moves through these structures, the basilar membrane, located between the scala media and scala tympani, shifts against the tectorial membrane, respectively [15]

C. Hearing Loss

Hearing loss is a condition that occurs when the transmission of sound from the outer ear to the brain is disrupted. The disruption can occur at any stage, before, after, or after and before the cochlea. Hearing loss can be conductive, sensorineural, or mixed [16]. Conductive hearing loss is caused by impaired transmission of sound waves to the cochlea, which ultimately causes problems primarily in the outer ear and auditory canal. The problem may originate in the external canal due to obstruction from debris, wax, or foreign objects, or may result from a congenital abnormality such as aural atresia, which causes incomplete development of the outer ear. Perforation of the tympanic membrane can also cause conductive hearing loss. Perforation can be associated with trauma from using cotton swabs to clean the ears, barotrauma from diving, or as a sequela of otitis media [17]. Sensorineural hearing loss (SNHL) is the most common type and accounts for the majority of all hearing loss. Hearing loss resulting from neurological, otologic, or vascular disorders is referred to as SNHL. Sensorineural hearing loss can be caused by damage to the auditory nerve, the vestibulocochlear nerve, or the central auditory processing area of the brain [18].

Noise can damage the sensory hair cells of the inner ear through direct mechanical pressure and intense sound pressure as well as by activation of stress-induced molecular pathways, including the formation of reactive oxygen species and excess calcium. Assessed by the decibel level and duration of exposure this can cause noise-induced hearing loss which can also be known as Noise Induced Hearing Loss (NIHL) [19]. NIHL is hearing loss caused by exposure to sounds that are too strong so that they can damage hearing cells, this kind of hearing loss is often described as "noise-induced". NIHL can be classified into three, namely, Temporary Threshold Shift (TTS), Permanent Threshold Shift (PTS), and Acoustic Trauma. Four factors contribute to the effects of noise, namely, sound level (volume), audio spectrum, distribution of exposure time, cumulative noise exposure over days, weeks or years [20]. The term "transient threshold shift" is used to describe objective changes in hearing acuity that can be measured on audiometry immediately after an episode of loud noise exposure and which return to pre-exposure levels after a few days to two weeks. A temporary threshold shift is subjectively characterized by decreased hearing sensitivity, a feeling of fullness in the ear,

tinnitus (ringing), and the perception of hearing loss. Prolonged or repeated exposure to noise can lead to the death of sensory hair cells and result in permanent hearing loss, referred to as a “permanent threshold shift.” Hair cell death may be followed by a gradual loss of spiral ganglion neurons over months or years [21].

D. Earphones

To listen to audio, people usually need a device called earphones, which convert electrical energy into sound waves. Earphones are most often used for media consumption and digital communication via gadgets such as smartphones and computers. [22]. Consumers are increasingly turning to earphones due to advances in audiovisual and communication technology. Earphones are now commonly used to listen to music on mobile phones and other audio devices, thanks to these innovations. People, especially teenagers, have made listening to music through headphones or earphones a regular part of their lives. Most teenagers wear earphones or headphones, typically to listen to music for extended periods of time. They do so even while engaging in other activities, such as traveling, exercising, or even sleeping. Prolonged use can cause continuous noise, which can damage hearing [6].

Recent advances in audiovisual technology have had a significant impact on the creation of modern music players and the hearing aids used to listen to music on these devices. Earphones, headsets, headphones, and other types of hearing aids are just a few of the many types that have been produced, with varying capacities and sound output functions [23]. According to one study, 88.2% of 1,407 children used personal music players, also known as MP3 players, and 27.4% of these children listened to music at the highest possible volume for extended periods without interruption. Another study found that 94.3% of those participating used personal music players daily for between one and three hours over three years. Conversely, earphones are the most common form of amplification for the hearing impaired. This demonstrates the widespread use of earphones among today's youth [23].

E. The Impact of Earphone Use Patterns on Hearing Loss

While earphone use is currently more effective than loudspeakers, it's important to note that earphone use can impact hearing response and chronic noise. Even at the same volume, listening to music through earphones produces a higher overall noise intensity than listening to the same music without earphones. When someone listens to music with earphones, the sound source is physically closer to them than when listening without earphones. Earphones also don't fully distribute sound and don't completely block out external noise, which can contribute to hearing loss. Earphone use patterns can be divided into four components: frequency, duration, noise volume, and length of earphone use. These four components are the primary assessments for determining earphone use patterns, which are then classified into two categories: risky and non-risky earphone use patterns [24].

Hearing loss is one of the effects caused by earphone use, resulting from improper use. Continuous exposure is a risk factor for hearing loss. Approximately 5 to 10% of earphone users are at risk of hearing loss and even permanent hearing loss if they listen to music for more than one hour a day at high volume [23]

Due to the length of earphone use of more than three years and daily use, earphones require attention, as literature indicates that prolonged exposure to loud noise can cause noise-induced hearing loss (NIHL) [25]. Research by Setianiet (2018) found that 77.8% of those with a history of noise exposure from earphones experienced NIHL, and 22.2% experienced NIHL with more than five years of use. Therefore, using earphones for five years or more can increase the risk of NIHL [26]. Noise-induced hearing loss (NIHL) is a form of sensory deficit that can lead to sensorineural hearing loss due to continuous exposure to loud noise over a prolonged period [27].

F. The Impact of Hearing Loss on Quality of Life

According to the World Health Organization (WHOQOL), quality of life is defined as an individual's perception of their life in society within the context of their culture and existing value systems, related to goals, expectations, standards, and concerns. Based on this definition, quality of life can be defined as life satisfaction or the fulfillment of life's needs based on a person's perceived physical, psychological, and social conditions [28], [29]. Factors that can influence a person's quality of life include psychological conditions. For example, psychological conditions can arise from the impact of an individual's physical limitations, which can lead to anxiety disorders, which can reduce the sufferer's ability to work and socialize with others. This can indirectly impact a person's quality of life. Beyond psychological factors, there are social relationship factors such as social support, which is the presence of others who can be relied upon to provide assistance, encouragement, acceptance, and attention, thereby improving and influencing the well-being or quality of life for the individual concerned [30]

Hearing loss can lead to a potential handicap or disability, limiting the individual's ability to perform various tasks in life. This can have psychological impacts, such as communication barriers, leading to anxiety, depression, and even impaired social interaction. Psychological and social relationships are factors that can impact a person's quality of life [31]

Material And Methods

This study uses a cross-sectional observational analytical research design, the researcher wants to prove the extent of the closeness of the relationship between a variable and another variable, where in this study the researcher wants to know the relationship

between earphone use patterns and the risk of hearing loss which can affect the quality of life of students of the Faculty of Medicine, Indonesian Christian University Class of 2019-2021.

This research was conducted from December to January 2023 at the Faculty of Medicine, Christian University of Indonesia, on Jl. Mayjen Sutoyo No. 2, RT. 2/RW. 11, Cawang Village, Kramat Jati District in East Jakarta City. The study population consisted of students from the Faculty of Medicine at Universitas Kristen Indonesia who graduated during the 2019-2021 intake. The sample consisted of all students who met the inclusion and exclusion criteria as in Table 1, and had given their consent.

Table 1. Inclusion and Exclusion Criteria

Criteria	Inclusion Criteria	Exclusion Criteria
Participants	Students from the 2019–2021 intake of UKI from any faculty.	Students who have previously been diagnosed with a hearing disorder (e.g., presbycusis, otosclerosis, chronic otitis media).
	Willing to participate and sign the informed consent form.	Students who are currently taking medication known to be ototoxic (e.g., certain antibiotics or chemotherapy agents).
	Own and regularly use earphones/headphones for leisure or study purposes.	Students who did not complete the entire questionnaire.

The sampling technique used in this study was non-probability sampling, namely, consecutive sampling, which was used to determine the research sample based on predetermined criteria and objectives. The research instrument consisted of two questionnaires: the Earphone Usage Pattern Questionnaire (EUPQ), which was used to examine respondents' habits of using earphones, and the Hearing Handicap Inventory for Adults (HHIA) questionnaire, which was used to determine the quality of life of earphone user respondents.

Data collected through the questionnaire were processed using IBM SPSS (Statistics for Social Science) and Microsoft Office Excel. Categorical data were displayed as percentages, while numerical data were displayed as means and standard deviations. All data were presented in tabular form. An inferential analysis was then performed to analyze the relationship between EUPQ and HHIA using the Chi-square test. The Pearson correlation test was used to analyze the relationship between EUPQ scores and HHIA, provided the data were normally distributed; otherwise, the Spearman test was employed.

Results

A. Respondent Profile

This study used an online questionnaire created by Google Forms and administered to students at the Faculty of Medicine, Universitas Kristen Indonesia, between December 2022 and January 2023. The aim of this study was to determine the relationship between earphone use patterns and the risk of hearing loss, which impacts quality of life, as measured by the Hearing Handicap Inventory for Adults (HHIA) questionnaire, adapted into Indonesian.

The study population comprised 225 students from the 2019, 2020, and 2021 intakes at the Faculty of Medicine, Universitas Kristen Indonesia.

The study lasted three months and was divided into two parts. In the first phase, participants were asked to complete a questionnaire about their regular earphone use patterns, and then they were asked to complete the HHIA questionnaire. The data was then analyzed, and a thesis was prepared.

Table 1: EUPQ Distribution Overview

Assessment Criteria		<i>f</i>	%
Length of Use of Earphones	0-3 years	52	23,1
	>3 years	173	76,9
Duration of Earphone Use	<1 hour	97	43,1
	1-2 hours	66	29,3
	>2-4 hours	36	16,0
	>4-6 hours	14	6,2
	>6-8 hours	5	2,2
	>8 hours	7	3,1
Frequency of Earphone Use	1-2 days	115	51,1
	3-4 days	62	27,6
	5-6 days	17	7,6
	Every Days	31	13,8
Earphone Usage Volume	<60%	140	62,2
	60-80%	69	30,7
	>80%	16	7,1
Earphone Usage Patterns	No Risk	85	37,8
	Risk	140	62,2
Total		225	100,0




Figure 1. The Chart of EUPQ Distribution Overview

Based on Table 1, 52 respondents used earphones for 0-3 years, representing 23.1% of the total, while 173 respondents used earphones for more than 3 years, representing 76.9% of the total. Based on these findings, the majority of students using earphones at the Faculty of Medicine, Universitas Kristen Indonesia (UKI) have used them since their teenage years, with some even using them since junior high school. A total of 97 respondents (43.1%) used earphones for more than 1 hour, 66 respondents (29.3%) used earphones for 1-2 hours, 36 respondents (16.0%) used earphones for more than 2-4 hours, 14 respondents (6.2%) for more than 4-6 hours, 5 respondents (2.2%) for more than 6-8 hours, and 7 respondents (3.1%) for more than 8 hours. From this table, it can be seen that students of the Faculty of Medicine, Christian University of Indonesia, Class of 2019, 2020, and 2021 mostly use earphones for more than one hour per day.

In total, 115 respondents (51.1%) reported using earphones 1-2 days a week, while 62 (27.6%) used them 3-4 days a week, 17 (7.6%) used them 5-6 days a week, and 31 (13.8%) used them daily. Here, we see that first-, second-, and third-year medical students at the Christian University of Indonesia used earphones between two and three days a week. 140 (62.2%) respondents used earphones with a volume of <60% (≤ 9 clicks of 0 volume), 69 (30.7%) used earphones with a volume of 60-80% (10-12 clicks of 0 volume), and 16 (7.1%) used earphones with a volume of >80% (>12 clicks of 0 volume). In this case, students at the Faculty of Medicine, Indonesian Christian University, Class of 2019, 2020, and 2021 mostly used earphones with a volume of <60% (≤ 9 clicks from 0% volume). 85 respondents (37.8%) had earphone usage

patterns that did not pose a risk, while 140 (62.2%) had earphone usage patterns that did. From what we can tell from this study, the majority of medical students at the Christian University of Indonesia (UI) in the 2019, 2020, and 2021 intakes are at risk of hearing loss. Heavy, long-term earphone use (>3 years), daily use (>1 hour), weekly use (>4 days), and high volume use (>60%) all contribute to risky earphone use.

Table 2: HHIA Score Results Overview

Indicator		<i>f</i>	%
Hearing Handicap Inventory Adults Score Results	No Handicap	167	74,2
	Mild Moderate Handicap	22	9,8
	Severe Handicap	36	16
	Total	225	100,0

Table 2 shows the scores from the Hearing Handicap Inventory for Adults (HHIA) questionnaire distributed to groups at risk for hearing loss. Of the respondents who completed the questionnaire, 167 (74.2%) had no handicap, 22 (9.8%) had a mild-moderate handicap, and 36 (16%) had a severe handicap.

Table 3: The relationship between EUPQ and HHIA

Earphone Usage Patterns	Hearing Handicap Inventory Adults (HHIA)						p-value
	No Handicap		Mild Moderate Handicap		Severe Handicap		
	<i>f</i>	%	<i>f</i>	%	<i>f</i>	%	
No Risk	71	83,5	8	9,4	6	7,1	0,015
Risk	96	68,6	14	10,0	30	21,4	
Total	167	74.2	22	9.8	36	16	

Table 3 shows that 96 respondents (68.6%) had earphone use patterns that put them at risk for hearing loss, 14 (10.0%) had mild-moderate handicaps, and 30 (21.4%) had severe handicaps. Meanwhile, 71 respondents (83.5%) were not at risk for hearing loss, 8 (9.4%) had mild-moderate handicaps, and 6 (7.1%) had severe handicaps. Statistical tests revealed a statistically significant relationship between risky and non-risky earphone use and handicaps that can affect respondents' quality of life, with a p-value of 0.015 ($p < 0.05$). Therefore, it can be concluded that there is a significant relationship between earphone use patterns and the risk of hearing loss. This relationship affects a person's quality of life. To determine the significance of the correlation and the degree of correlation between the independent and dependent variables, an analysis was conducted based on numerical data from the earphone usage pattern questionnaire and HHIA scores using the Pearson correlation test if the data were normally distributed and alternatively using the Spearmann correlation test if the data

were not normally distributed. A normality test was conducted to determine the normality of the data distribution, using the Kolmogorov-Smirnov test because the sample size was >50.

Table 4: Normality Test Results

Variable	Mean	SD	Median	Min	Max	p-value
Earphone Usage Patterns	9,48	3,24	9,0	4,0	18,0	<0,001 ^a
HHIA	16,14	21,42	6,0	2,0	96,0	<0,001

Description:

^aKolmogorov-Smirnov

Table 4 shows the results of the Kolmogorov-Smirnov test on earphone usage patterns, with a p value of <0.05 and HHIA of <0.05.

Table 5: Spearman Test Results

Variable	r	P-value
EUPQ		
HHIA	0,202	0,002 ^b

Description:

^bSpearman's Test

A Spearman's correlation test followed this. Table 4.5 shows a weak, significant relationship between earphone use patterns and the risk of hearing loss, which impacts students' quality of life.

B. Discussion

1. Distribution of Respondents' Earphone Usage Patterns

From the results of the research conducted by researchers, a questionnaire was completed to assess earphone usage patterns, which included information on duration, frequency, duration, and volume. It was found that the majority of respondents (173 respondents) had used earphones for more than three years, totaling 76.9%. This indicates that students from the Faculty of Medicine, Christian University of Indonesia (2019-2021) have been listening to music through earphones since they were in school. This aligns with research conducted by Fatimah Hamzah and Nurul, whose respondents were students from the Faculty of Medicine, Muhammadiyah University of Makassar, with an average duration of earphone use of more than three years (95.7%) [32]. According to the ENT-KL textbook, hearing loss due to prolonged exposure to excessively loud noise can lead to sensorineural hearing loss [33].

Commented [L1]: This article's discussion links the research findings to previous studies, highlighting the importance of earphone usage patterns in increasing the risk of hearing loss. The author also explains how factors such as the duration and volume of earphone use contribute to the risk of hearing loss. This discussion emphasizes the importance of awareness of the negative impacts of unwise earphone use.

Strengths:

1. A fairly robust discussion, linking the research findings to existing literature, provides readers with a better understanding of the topic.
2. The author also cites several other studies that support the findings of this study, enriching the discussion.

Revision Suggestions:

1. The discussion could be more in-depth by further exploring the biological mechanisms involved in listening to music at high volume through earphones and how this affects hearing in the long term.
2. A discussion of other factors influencing quality of life, such as social support or mental health, which may interact with the results of this study could also be added.

According to the World Health Organization (WHO), the largest number of earphone users worldwide are adolescents aged 12 to 24, who constitute part of the urban population [34].

The use of earphones has become commonplace in modern culture, especially among young people, who are more likely to use them as part of their daily routine than older generations. Earphones can be used as a means of listening to music and communicating because they can produce clearer sound than listening through the earphone speakers themselves. Furthermore, earphones are easy to carry anywhere, the volume can be adjusted, and they can provide privacy for the user. Regarding the duration of earphone use, the respondents in this study mostly used earphones for less than 1 hour, amounting to 97 people (47.1%). This result is in line with research by Martanegara and Ilman, who found that most respondents in the study used earphones for <1 hour, accounting for 61.34%. Noise-induced hearing loss can occur in the general public, especially teenagers, if they are often exposed to the use of earphones at high volume and continue to use them for an average of one to three hours per day [26]. More frequent use of earphones can be caused by the attachment to using earphones at high intensity because the use of earphones can be used in all things, such as when driving, sitting relaxing, doing activities, this earphone is very helpful so that even while doing activities, the person can still communicate or listen to music [22].

This study also found that 51.5% of the sample, or 115 students, from the Faculty of Medicine, Universitas Kristen Indonesia, Class of 2019, 2020, and 2021, mostly used earphones once or twice a week. A low frequency of earphone use can be determined when respondents wear their earphones from 1-2 days per week. This finding is consistent with that obtained from a study conducted in the past by Fatimah Hamzah and Nurul on students of the Faculty of Medicine, Universitas Muhammadiyah Makassar. The researchers found that almost half of the students (47.8%) used earphones for 1-2 days per week.

Based on the results of this study, as many as 140 respondents (62.2% of the total) often use earphone volume with > 60% on earphones. This is different when compared to research by Velaro, Adrian on Medical Faculty Students, University of North Sumatra, most of the respondents in their study used earphones with a volume of 60-80%. 26 Respondents who use earphones with high volume usually occur because users are not satisfied listening to them at low volume, so they choose to increase the volume of the sound produced by the earphones even though they are aware of the dangers of using earphones with high volume. This occurs even though respondents are aware of the dangers of using earphones with high volume [35]. If the use of earphones with high volume has become a habit for its users, there will be a trauma effect on the sound receptors due to the use of earphones with high volume intensity and cause a decrease in hearing function.

2. HHIA Score Overview

Information about hearing loss, expectations, and motivation can be collected using the Hearing Handicap Inventory for Adults (HHIA). The World Health Organization (WHO) defines disability or handicap as "a disadvantage to a person due to an impairment or impairment that limits or prevents the fulfillment of a normal role," taking into account the person's age, gender, and social and cultural context. The HHIA is a questionnaire used to assess the extent of handicap in adults with hearing loss. The HHIA is just one of many self-report instruments available to measure the functional impact of hearing loss on a person's daily life. HHIA results can be used to address some of the psychological impacts of hearing loss. The higher the HHIA score, the greater the recognition that hearing loss is a significant disability for the respondent due to the greater limitations or obstacles experienced by the sufferer in fulfilling their daily roles, resulting in a diminished quality of life.

This tool can assist audiologists by providing additional information regarding: (1) helping to substantiate complaints of hearing loss that are not apparent on conventional audiometric testing results; (2) facilitating candidate decisions regarding the use of amplification (hearing aids); (3) assisting in the counseling process; (4) serving as a guide for determining client-centered rehabilitation programs; and (5) serving as a measurement criterion in rehabilitation evaluations, including the benefits of post-surgical hearing aids.⁴⁶ Hearing loss can impact quality of life and limit activities or participation in daily life. The impact of this hearing loss can be assessed using the HHIA questionnaire [36]. In this study, the researchers used the HHIA questionnaire, which had been adapted into Indonesian and had been tested for validity and reliability.

The HHIA questionnaire in this study was administered to groups of people at and not at risk of developing hearing loss due to earphone use. 225 people (100%) completed the HHIA questionnaire. The results showed that 74.2% had no handicap, 9.8% had mild-moderate handicap, and 16% had severe handicap. In a previous study by Marcus and Dayse, the majority of respondents (70.8%) had severe handicap, 22.9% had mild-moderate handicap, and 6.3% had no handicap [37]. The HHIA questionnaire varies widely and depends on life experiences, health expectations, and even the individual's ability to adapt. People with hearing loss will experience communication difficulties and experience social and emotional difficulties in their daily lives. This will lead to different perceptions of their quality of life [36].

3. Relationship between Earphone Use Patterns and HHIA Scores

This study found a statistically significant relationship between earphone use patterns and the risk of hearing loss, which impacts the quality of life of students at the Faculty of Medicine, Christian University of Indonesia, in the 2019, 2020, and 2021 intakes ($p = 0.015$) of the 2019, 2020, and 2021 intakes ($p < 0.05$). Furthermore, the Spearman correlation test

revealed a weakly significant relationship between earphone use patterns and the risk of hearing loss, which impacted students' quality of life ($p = 0.002b$). Similarly, research by Marcus and Dayse confirmed that hearing loss can significantly impact quality of life [37].

The correlation between the two variables supports this finding. Incorrect use of earphones can lead to a risk of hearing loss and negatively impact quality of life, as measured by the HHIA questionnaire. Earphone use patterns can increase the risk of hearing loss, which can lead to physical, social, and emotional changes in individuals. These changes can persist and disrupt daily activities, significantly impacting quality of life, especially for those at risk for hearing loss.

The results demonstrating a higher HHIA score in respondents with risky earphone use patterns can be explained through well-established biological mechanisms. Exposure to sounds above 85 dB for long durations causes damage to outer hair cells (OHCs) in the cochlea. These OHCs function as mechanical amplifiers, improving frequency selectivity and sound sensitivity. High sound pressure levels lead to excessive mechanical vibration, which triggers oxidative stress and mitochondrial dysfunction in cochlear cells. Over time, this results in apoptosis and irreversible loss of sensory cells, leading to permanent threshold shifts and progressive hearing impairment.

Furthermore, loud music especially when transmitted directly into the ear canal via in-ear earphones causes temporary threshold shifts that, with repeated exposure, accumulate into permanent hearing loss. Research indicates that high-volume listening can induce excessive glutamate release, resulting in excitotoxicity at the ribbon synapses between inner hair cells and auditory nerve fibers. This "hidden hearing loss" may develop even before pure-tone audiometry detects deficits, and can manifest as difficulties in speech.

Conclusion

This study shows that students with risky earphone use patterns—such as long listening duration, high volume, and frequent weekly use—are more likely to experience hearing difficulties based on their HHIA scores. Although most students reported no handicap, a meaningful number already showed mild to severe hearing problems, indicating early signs of noise-induced hearing loss. Unsafe listening habits can gradually damage the cochlea and affect communication, concentration, and overall quality of life. Because many students have used earphones since adolescence, early prevention is important. To reduce the risk of hearing loss, educational institutions and related agencies should take more active steps, such as: 1) Providing education about safe listening habits (e.g., 60–60 rule); 2) Offering regular hearing screenings for students; 3) Creating campus policies that encourage healthy gadget use; 4) Running digital campaigns to raise awareness of safe earphone practices; 5) Partnering with health professionals to offer counseling for at-risk students. In summary, unsafe earphone use is a growing concern among young adults, and schools and health agencies should work together to promote safer listening habits and protect students' long-term hearing health.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- [1] C. J. Plack, *The sense of hearing*. Taylor and Francis, 2023. doi: 10.4324/9781003303329/SENSE-HEARING-CHRISTOPHER-PLACK.
- [2] W. H. Organization, *World report on hearing*. 2021. Accessed: Oct. 07, 2025. [Online]. Available: https://books.google.com/books?hl=en&lr=&id=zMRqEAAQBAJ&oi=fnd&pg=PR5&dq=WHO.+World+Report+On+Hearing.+2021&ots=cHAtI_6aMx&sig=5ks-DjL_goiQWico-iRullwvc-8
- [3] C. L. Nieman and E. S. Oh, "Hearing loss," *acpjournals.org* CL Nieman, *ES OhAnnals Intern. Med.* 2020•*acpjournals.org*, vol. 173, no. 11, pp. ITC81–ITC96, Dec. 2020, doi: 10.7326/AITC202012010.
- [4] B. Putri, R. Halim, H. N.-J. Kesmas, and undefined 2021, "Studi kualitatif gangguan pendengaran akibat bising/noise induced hearing loss (NIHL) pada marshaller di Bandar Udara Sultan Thaha Kota Jambi tahun 2020," *Mail. Putri, R Halim, HS NasutionJurnal Kesmas Jambi, 2021*•*mail.online-journal.unja.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://mail.online-journal.unja.ac.id/jkmj/article/view/12400>
- [5] N. Sarah, ... F. L.-J. (Jurnal K., and undefined 2016, "Hubungan Penggunaan Earphone Dengan Gangguan Pendengaran Pada Siswa Sma Negeri 9 Manado," *ejournal.unsrat.ac.id* NA Sarah, F Lintong, JF RumampukJJK (Jurnal Kedokt. Klin. 2016•*ejournal.unsrat.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.unsrat.ac.id/index.php/jkk/article/view/14363>
- [6] A. Budiartman, A. Asnawati, H. Huldani, ... S. K.-, and undefined 2024, "GAMBARAN TINGKAT PENGETAHUAN MAHASISWA PSKPS FK ULM ANGKATAN 2020 TENTANG FAKTOR RISIKO NOISE-INDUCED HEARING LOSS (NIHL)," *ppjp.ulm.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ppjp.ulm.ac.id/journals/index.php/hms/article/view/13178>
- [7] P. Berliana, S. Dan, S. Keman, D. Kesehatan, L. Fakultas, and K. Masyarakat, "Pengaruh Penggunaan Pelindung Telinga dan Earphone Terhadap Noise Induced Hearing Loss Dan Tinitus Pada. Pekerja Bengkel," *e-journal.unair.ac.id* PB Syah, S KemanJurnal Kesehat. Lingkungan, 2017•*e-journal.unair.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://e-journal.unair.ac.id/JKL/article/download/9148/5147>

- [8] "Susiyanti E, Imanto M. Efek Penggunaan Earphone sebagai... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Susiyanti+E%2C+Imanto+M.+Efek+Penggunaan+Earphone+sebagai+Faktor+Resiko+Kejadian+Noise+Induced+Hearing+Loss.+Majority.+2020%3B+9%282%29%3A+63-67.+&btnG= (accessed Oct. 07, 2025).
- [9] E. Yuliyani, D. Yudhanto, ... W. A.-I. J., and undefined 2022, "Peranan program acara stasiun TVRI NTB sebagai salah satu media edukasi kesehatan pendengaran," *Acad. Yuliyani, D Yudhanto, WS Aff. E Triani, F Libr. DAIS PrajnyaswarilNDRA J. Pengabdi. Kpd. Masyarakat, 2022*•*academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/93056133/125.pdf>
- [10] L. Moneta, L. Q.-D.-O.-H. and N. Surgery, and undefined 2017, "Embryology and anatomy of the ear," *Elsevier*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S1043181017300398>
- [11] R. Maynard and N. Downes, *Anatomy and histology of the laboratory rat in toxicology and biomedical research*. 2019. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=2kGHDwAAQBAJ&oi=fnd&pg=PP1&dq=Maynard+RL,+Downes+N.+The+Ear.+In:+Anatomy+and+Histology+of+the+Laboratory+Rat+in+Toxicology+and+Biomedical+Research+%5BInternet%5D.+Elsevier%3B+2019.+p.+293-302.+Available+from:+https://linkinghub.elsevier.com/retrieve/pii/B978012811837500023X&ots=-lqlh67SG&sig=eYuR9X9RkEvkTDRjKq-C7whl70>
- [12] A. Szymanski and Z. Geiger, "Anatomy, head and neck, ear," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/article/nbk/nbk470359>
- [13] A. Szymanski, J. Toth, M. Ogorevc, and Z. Geiger, "Anatomy, head and neck, ear tympanic membrane," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk448117>
- [14] "Sánchez López de Nava A, Lasrado S. Physiology, Ear. 2022 - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Sánchez+López+de+Nava+A%2C+Lasrado+S.+Physiology%2C+Ear.+2022&btnG= (accessed Oct. 07, 2025).
- [15] H. White, M. Helwany, ... A. B.-S., and undefined 2023, "Anatomy, head and neck, ear organ of corti," *ncbi.nlm.nih.gov/HJ White, M Helwany, AR Biknevicius, DC PetersonStatPearls [Internet], 2023*•*ncbi.nlm.nih.gov*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.ncbi.nlm.nih.gov/books/NBK538335/>
- [16] R. Sataloff and P. Roehm, *Hearing loss*. 2024. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=IG0iEQAAQBAJ&oi=fnd&pg=PP1&dq=Hearing+Loss&ots=Y1HvwBOzEA&sig=EjaY-3SDMk5upL0768pSMieSHE>
- [17] T. Sooriyamoorthy and O. De Jesus, "Conductive hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk563267>
- [18] R. Tanna, J. Lin, and O. De Jesus, "Sensorineural hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk565860>
- [19] S. C.-C. E. and H. Health and undefined 2019, "Noise-induced hearing loss," *account.cehh.press.lshrm.ac.ukS ChadhaCommunity Ear Hear. Heal. 2019*•*account.cehh.press.lshrm.ac.uk*, Accessed: Oct. 07, 2025. [Online]. Available: <https://account.cehh.press.lshrm.ac.uk/index.php/s-j-cehh/article/view/31>
- [20] "Chan Y, Goddard JC. Preface. In: KJ Lee's Essential... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Chan+Y%2C+Goddard+JC.+Preface.+In%3A+KJ+Lee's+Essential+Otolaryngology%3A+Head+and+Neck+Surgery%2C+12e+%5BInternet%5D.+New+York%2C+NY%3A+McGraw-Hill+Education%3B+2019.+Available+from%3A+accesssurgery.mhmedical.com%2Fcontent.aspx%3Faid%3D1172366487&btnG= (accessed Oct. 07, 2025).
- [21] L. Cunningham, D. T.-N. E. J. of Medicine, and undefined 2017, "Hearing loss in adults," *Mass Med. SocLL Cunningham, DL TucciNew Engl. J. Med. 2017*•*Mass Med. Soc*, vol. 377, no. 25, pp. 2465-2473, Dec. 2017, doi: 10.1056/NEJMRA1616601.
- [22] M. Erlanda, P. Negara, I. Triansyah, D. Hasni, and B. Y. Febrianto, "Hubungan Intensitas Penggunaan Earphone dengan Derajat Gangguan Pendengaran pada Mahasiswa Fakultas Kedokteran Universitas Baiturrahmah," *Journal.scientific.idMEP Negara, I Triansyah, D Hasni, BY FebriantoScientific Journal, 2022*•*journal.scientific.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://journal.scientific.id/index.php/sciena/article/view/49>
- [23] A. Velaro, D. Z.-O. R. L. Indonesiana, and undefined 2021, "The correlation between earphone usage patterns with the incidence rate and severity of tinnitus," *orli.or.idAJ Velaro, D Zahar. Rhino*

- Laryngol. Indones.* 2021•orli.or.id, Accessed: Oct. 07, 2025. [Online]. Available: <http://orli.or.id/index.php/orli/article/view/432>
- [24] M. Zhang, A. Jeske, S. Y. (yrs), and undefined 2017, "Changes in the frequency and intensity of Tinnitus using the Suppressive Noise Spectrum," *m.karmaimpex.com* M Zhang, A Jeske, S Youngage (yrs), 2017•*m.karmaimpex.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://m.karmaimpex.com/hjed/pdf/ated-aid1002.pdf>
- [25] B. Rosita, "Pemakaian Earphone dalam Mendengar Musik yang Menimbulkan Bising di Kalangan Milenial," 2019, Accessed: Oct. 07, 2025. [Online]. Available: https://osf.io/asmb_v1/
- [26] L. Setiani, N. Syakila, Y. Y.-J. K. N. Medika, and undefined 2018, "Hubungan Lama Paparan Penggunaan Earphone Musik Terhadap Terjadinya Gangguan Pendengaran Akibat Bising pada Mahasiswa Fakultas Kedokteran," *jknamed.com* L Setiani, N Syakila, Y YusniJurnal Kedokt. Nanggroe Med. 2018•*jknamed.com*, Accessed: Oct. 07, 2025. [Online]. Available: <http://www.jknamed.com/jknamed/article/view/51>
- [27] T. Wang, T. Chang, R. Tyler, Y. Lin, W. L.-B. sciences, and undefined 2020, "Noise induced hearing loss and tinnitus—new research developments and remaining gaps in disease assessment, treatment, and prevention," *mdpi.com* TC Wang, TY Chang. R Tyler, YJ Lin, WM Liang, YW Shau, WY Lin, YW Chen, CD LinBrain Sci. 2020•*mdpi.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.mdpi.com/2076-3425/10/10/732>
- [28] "Pratiwi Y. Pengaruh Dukungan Sosial Terhadap Kualitas... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Pratiwi+Y.+Pengaruh+Dukungan+Sosial+Terhadap+Kualitas+Hidup+Lanjut+Usia+di+Pusar+Santunan+Keluarga+%28PUSAKA%29+Kecamatan+Pancoran+Jakarta+Selatan+%5Bskripsi%5D.+Jakarta%3A+Fakultas+Dakwah+dan+Komunikasi+Universitas+Islam+Negeri+Syarif+Hidayatullah%3B+2017&btnG= (accessed Oct. 07, 2025).
- [29] "WHO. WHOQOL User Manual Programme On Mental Health... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=WHO.+WHOQOL+User+Manual+Programme+On+Mental+Health+Division+Of+Mental+Health+And+Prevention+Of+Substance+Abuse+World+Health+Organization.+2020&btnG= (accessed Oct. 07, 2025).
- [30] D. Maya et al., "Kualitas hidup lansia ditinjau dari sabar dan dukungan sosial," *ejournal.umm.ac.id* DMP Sari, CYD Lestari, EC Putra, F Nashorijurnal Ilm. Psikol. Ter. 2018•*ejournal.umm.ac.id*, vol. 06, no. 02, pp. 2540–8291, 2018, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.umm.ac.id/index.php/jipt/article/view/5341>
- [31] A. Nurfaizi, "MENINGKATKAN KEMAMPUAN DETEKSI DINI DAN PENATALAKSANAAN GANGGUAN PENDENGARAN DI KALIBARU KULON BANYUWANGI," *academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/108530358/14428.pdf>
- [32] "Hamzah NF. Pengaruh Perilaku Penggunaan Earphone... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Hamzah+NF.+Pengaruh+Perilaku+Penggunaan+Earphone+Terhadap+Gangguan+Telinga+Pada+Mahasiswa+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar+%5Bskripsi%5D.+Makassar%3A+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar%3B+2021&btnG= (accessed Oct. 07, 2025).
- [33] "Soepardi EA, Iskandar N, Bashiruddin J, Restuti RD.... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Soepardi+EA%2C+Iskandar+N%2C+Bashiruddin+J%2C+Restuti+RD.+Buku+Ajar+Ilmu+Kesehatan%3A+Telinga%2C+Hidung%2C+Tenggorok%2C+Kepala+%26+Leher.+7th+ed.+Jakarta%3A+BP+FKUI%3B+2012.+&btnG= (accessed Oct. 07, 2025).
- [34] N. Yudiasari, S. W.-P. J. K. Masyarakat, and undefined 2021, "Pengembangan Media Promosi Kesehatan Tentang Bahaya Penggunaan Earphone Mahasiswa Universitas Airlangga Di Surabaya," *103.245.72.38* N Yudiasari, S WidatiPreventif J. Kesehat. Masyarakat, 2021•*103.245.72.38*, vol. 12, pp. 57–86, 2021, Accessed: Oct. 07, 2025. [Online]. Available: <http://103.245.72.38/index.php/preventif/article/view/182>
- [35] S. Levey, B. Fligor, ... C. G.-C. I. in, and undefined 2012, "The effects of noise-induced hearing loss on children and young adults," *pubs.asha.org* S Levey, BJ Fligor, C Ginocchi, L KagimbiContemporary Issues Commun. Sci. Disord. 2012•*pubs.asha.org*, vol. 39, no. Fall, pp. 76–83, Oct. 2012, doi: 10.1044/CICSD_39_F_76.
- [36] J. Bashiruddin, W. Alviandi, B. Bramantyo, and R. Ismail, "Praktik Klinis: Hearing Handicap Inventory For Adults Adaptasi Bahasa Indonesia Sebagai Alat Ukur Kualitas Hidup Pasien Gangguan Pendengaran," 2019, Accessed: Oct. 07, 2025. [Online]. Available: <https://scholar.ui.ac.id/en/publications/buku-panduanpetunjuk-panduan-praktik-klinis-hearing-handicap-inve>

- [37] M. Voola, D. T.-V.-T. M. Journal, and undefined 2020, "Quality of life handicap measured in patients with profound unilateral or bilateral deafness," *tasmanmedicaljournal.com* M Voola, D Távora-Viera *Tasman Med. Journal*, 2020•*tasmanmedicaljournal.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://tasmanmedicaljournal.com/2020/12/quality-of-life-handicap-measured-in-patients-with-profound-unilateral-or-bilateral-deafness/>

Authors



Lina Marlina    is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. She is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery. She is a member of the Indonesian Medical Association (IDI). Her professional interests include ENT and ENT and clinical research. She is actively involved in academic research and public health programs. She can be contacted at lina.marlina@uki.ac.id.



Sena Pongtuluran    is a general practitioner who recently graduated from the Faculty of Medicine, Christian University of Indonesia, Jakarta. She is currently pursuing a Master's degree in Hospital Administration (MARS) at Pelita Harapan University (UPH). Her interests include clinical medicine, community health, and healthcare management. She is passionate about contributing to healthcare innovation and evidence-based medical practice. She can be contacted at senapongtuluran@gmail.com



Fransiskus Harf Poluan    is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. He is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery at Mayapada Hospital, Kuningan, Jakarta. He is actively involved in academic research and public health programs. He can be contacted at fransiskus.poluan@uki.ac.id

Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life

^{1,2}Lina Marlina*, ¹Sena Pongtuluran, ¹Fransiskus Harf Poluan

Corresponding Author: *lina.marlina@uki.ac.id

¹ Department of Ear Nose and Throat, Faculty of Medicine, Universitas Kristen Indonesia

² General Hospital Universitas Kristen Indonesia, Jakarta, Indonesia

ARTICLE INFO

Article history

Received XX July 2022

Revised XX August 2022

Accepted XX August 2022

ABSTRACT

In today's modern life, most people are constantly exposed to noise every day, potentially disrupting their hearing function. One such factor is the use of hearing devices, namely earphones. Approximately 1.1 billion young adults and adolescents are at risk of experiencing hearing loss in their lifetime. If someone is at risk of hearing loss, it can undoubtedly impact their quality of life. Participants in this study were students at the Christian University of Indonesia in 2019, 2020, and 2021. The purpose of this study was to determine whether risky earphone use patterns are associated with quality of life. Data for this study were collected using a cross-sectional observational analytical methodology with consecutive sampling technique. The study sample was assessed with a questionnaire created using Google Forms. The questionnaire, which is divided into two parts: the Earphone Use Pattern Questionnaire (PPE) and the Hearing Handicap Inventory for Adults (HHIA), to determine the patient's quality of life, was then analyzed using the Chi-Square and Spearman tests. Of the 225 respondents, there were 62.2% risky earphone usage patterns and 37.8% non-risky earphone users. According to the HHIA score findings, 74.2% of people had no disabilities, 9.8% had mild to moderate disabilities, and 16.0% had severe disabilities, with $p = 0.015$ ($p < 0.05$) obtained through the use of the Chi-Square statistical test. There is a relationship between earphone usage patterns and the risk of hearing loss which affects the quality of life of students of the Faculty of Medicine, Universitas Kristen Indonesia Class of 2019, 2020, and 2021.

Keywords

Earphone use patterns
Hearing loss
Hearing Handicap
Inventory for Adults
Quality of Life

This is an open-access article under the [CC-BY-SA](#) license.



Introduction

The ability to hear is at the heart of all human communication, both with each other and with the world around us [1]. Hearing loss is one of the most common disabilities worldwide, affecting more than 1.5 billion people, according to the World Health Organization (WHO) [2]. While common, hearing loss often goes unrecognized by patients and doctors, leading to inadequate treatment. The prevalence of clinically significant hearing loss has been found to double every decade. [3]. In Southeast Asia, the prevalence of hearing loss is estimated at 27%, which equates to approximately 156 million people in the total population. In contrast, the prevalence of hearing loss in Indonesia nationally is 2.6% of the total population, and this figure is expected to increase over time. [4]

In today's modern life, most people are constantly exposed to noise that can potentially disrupt hearing function [5]. The use of personal audio devices, specifically earphones, is a significant and growing risk factor for hearing loss in the context of this study. WHO estimates that 1.1 billion adolescents and young adults are at risk of hearing loss as a direct result of exposure to loud noises from electronic devices. This risk is primarily associated with the increased volume of sound produced by these devices [6]. The use of earphones is a significant indicator of increased risk, especially for noise-induced hearing loss (NIHL), as earphones can bring the noise source closer to the ear, resulting in the listener receiving significantly greater

exposure [7]. A study by Rumampuk in Indonesia found that 26.7% of students who use earphones experience some level of hearing loss, with 6.7% experiencing moderate hearing loss [8]. According to data from the WHO, approximately 430 million people, or more than 5% of the global population, experience hearing loss that affects their quality of life. The habit of listening to music at high volumes via earphones and other noisy environments for long periods of time is a primary cause [9]. Based on the above background, the researcher wants to know the relationship between earphone usage patterns (including volume and duration) and the risk of hearing loss which affects the quality of life of students of the Faculty of Medicine, Indonesian Christian University, Class of 2019-2021.

Literature Review

A. Ear Anatomy

Human ear development begins with the appearance of the otic placode and vestibulocochlear ganglia at three weeks of gestation. This is followed by the appearance of the external canal, which begins to develop from the first branchial cleft in the fourth week, then forms a hole through ectodermal proliferation. By the 28th week, the epithelial core has canalized from medial to lateral, resulting in a fully patent external canal [10].

The ear can be divided into three parts: the outer ear, the middle ear, and the inner ear [11]. The outer ear consists of the auricle (also known as the pinna) and the external auditory canal, as well as the lateral surface of the tympanic membrane, which is part of the outer ear [12]. The middle ear consists of the tympanic membrane and three auditory ossicles: the malleus, incus, and stapes. These three ossicles allow sound waves to travel from the outer ear to the inner ear by connecting the tympanic membrane to the inner ear [13]. The inner ear is a system of canals and a cavity within the petrous part of the temporal bone. The inner ear is formed by the bony labyrinth (osseous labyrinth), which contains the membranous labyrinth. The bony labyrinth consists of three components: the semicircular canals, the vestibule, and the bony cochlea. The membranous labyrinth is located within the bony labyrinth and is a system of interconnected channels lined with epithelium and containing endolymph. The membranous labyrinth consists of the membranous semicircular canals, the utricle, the saccule, the endolymphatic duct, the endolymphatic sac, and the reuniens duct. The cochlear duct contains the organ of Corti, which is the organ of hearing. There are sensory nerve endings in the ampullae of the semicircular canals (crista ampullaris) and in the utricle and saccule (macula sacculi and utriculi) which function as static and kinetic senses.




Figure 2.1 Ear Anatomy [3]

B. Physiology of Hearing

Hearing and balance are functions performed by the human ear. The transduction mechanism in the ear is responsible for detecting and analyzing sound. This mechanism is responsible for converting sound waves into electrochemical impulses. If the anatomy is abnormal, hearing will not function properly even if the nerves are intact. Hearing is the process by which sound vibrations from the surrounding environment are converted into action potentials in the cochlea and auditory nerve. Sound is produced when an object is made to vibrate, such as when a guitar string is plucked. The resulting vibrational pressure is called a sound wave. Therefore, the ear is designed to detect and categorize various sonic qualities, including pitch (the frequency of the sound wave) and loudness (the perceived intensity of the sound). Hertz is the standard system of frequency measurement (Hz, cycles per second). The human ear can generally detect sounds between 1,000 and 4,000 hertz, but younger ears can detect sounds between 20,000 and 200,000 hertz. The decibel (dB) is the unit used to measure the loudness of sound, and the audible range that humans can tolerate on the decibel scale is from 0 to 130 dB (where the sound can become painful). To enter the central nervous system, each of these physical properties must first undergo some kind of transformation. Air vibrations are first converted into vibrations of the tympanic membrane, which is the first step in the process. Once they reach these structures, the vibrations are captured by the middle ear and ossicles. Then, in the inner ear and cochlea, these vibrations are converted into vibrations of fluid, and these fluid vibrations stimulate areas known as the basilar membrane and the organ of Corti. Ultimately, these vibrations are converted into nerve impulses, which are then transmitted to the nervous system [14]

The primary function of the organ of Corti is auditory signal transduction. Sound waves enter the ear through the auditory canal and cause vibrations in the tympanic membrane.

Movement of the tympanic membrane causes subsequent vibrations in the auditory ossicles, the three middle ear bones, which transfer energy to the cochlea through the oval window. As the oval window moves, the waves travel through the perilymph fluid in the scala tympani and then the scala vestibuli of the cochlea. As fluid moves through these structures, the basilar membrane, located between the scala media and scala tympani, shifts against the tectorial membrane, respectively [15]

C. Hearing Loss

Hearing loss is a condition that occurs when the transmission of sound from the outer ear to the brain is disrupted. The disruption can occur at any stage, before, after, or after and before the cochlea. Hearing loss can be conductive, sensorineural, or mixed [16]. Conductive hearing loss is caused by impaired transmission of sound waves to the cochlea, which ultimately causes problems primarily in the outer ear and auditory canal. The problem may originate in the external canal due to obstruction from debris, wax, or foreign objects, or may result from a congenital abnormality such as aural atresia, which causes incomplete development of the outer ear. Perforation of the tympanic membrane can also cause conductive hearing loss. Perforation can be associated with trauma from using cotton swabs to clean the ears, barotrauma from diving, or as a sequela of otitis media [17]. Sensorineural hearing loss (SNHL) is the most common type and accounts for the majority of all hearing loss. Hearing loss resulting from neurological, otologic, or vascular disorders is referred to as SNHL. Sensorineural hearing loss can be caused by damage to the auditory nerve, the vestibulocochlear nerve, or the central auditory processing area of the brain [18].

Noise can damage the sensory hair cells of the inner ear through direct mechanical pressure and intense sound pressure as well as by activation of stress-induced molecular pathways, including the formation of reactive oxygen species and excess calcium. Assessed by the decibel level and duration of exposure this can cause noise-induced hearing loss which can also be known as Noise Induced Hearing Loss (NIHL) [19]. NIHL is hearing loss caused by exposure to sounds that are too strong so that they can damage hearing cells, this kind of hearing loss is often described as "noise-induced". NIHL can be classified into three, namely, Temporary Threshold Shift (TTS), Permanent Threshold Shift (PTS), and Acoustic Trauma. Four factors contribute to the effects of noise, namely, sound level (volume), audio spectrum, distribution of exposure time, cumulative noise exposure over days, weeks or years [20]. The term "transient threshold shift" is used to describe objective changes in hearing acuity that can be measured on audiometry immediately after an episode of loud noise exposure and which return to pre-exposure levels after a few days to two weeks. A temporary threshold shift is subjectively characterized by decreased hearing sensitivity, a feeling of fullness in the ear,

tinnitus (ringing), and the perception of hearing loss. Prolonged or repeated exposure to noise can lead to the death of sensory hair cells and result in permanent hearing loss, referred to as a “permanent threshold shift.” Hair cell death may be followed by a gradual loss of spiral ganglion neurons over months or years [21].

D. Earphones

To listen to audio, people usually need a device called earphones, which convert electrical energy into sound waves. Earphones are most often used for media consumption and digital communication via gadgets such as smartphones and computers. [22]. Consumers are increasingly turning to earphones due to advances in audiovisual and communication technology. Earphones are now commonly used to listen to music on mobile phones and other audio devices, thanks to these innovations. People, especially teenagers, have made listening to music through headphones or earphones a regular part of their lives. Most teenagers wear earphones or headphones, typically to listen to music for extended periods of time. They do so even while engaging in other activities, such as traveling, exercising, or even sleeping. Prolonged use can cause continuous noise, which can damage hearing [6].

Recent advances in audiovisual technology have had a significant impact on the creation of modern music players and the hearing aids used to listen to music on these devices. Earphones, headsets, headphones, and other types of hearing aids are just a few of the many types that have been produced, with varying capacities and sound output functions [23]. According to one study, 88.2% of 1,407 children used personal music players, also known as MP3 players, and 27.4% of these children listened to music at the highest possible volume for extended periods without interruption. Another study found that 94.3% of those participating used personal music players daily for between one and three hours over three years. Conversely, earphones are the most common form of amplification for the hearing impaired. This demonstrates the widespread use of earphones among today's youth [23].

E. The Impact of Earphone Use Patterns on Hearing Loss

While earphone use is currently more effective than loudspeakers, it's important to note that earphone use can impact hearing response and chronic noise. Even at the same volume, listening to music through earphones produces a higher overall noise intensity than listening to the same music without earphones. When someone listens to music with earphones, the sound source is physically closer to them than when listening without earphones. Earphones also don't fully distribute sound and don't completely block out external noise, which can contribute to hearing loss. Earphone use patterns can be divided into four components: frequency, duration, noise volume, and length of earphone use. These four components are the primary assessments for determining earphone use patterns, which are then classified into two categories: risky and non-risky earphone use patterns [24].

Hearing loss is one of the effects caused by earphone use, resulting from improper use. Continuous exposure is a risk factor for hearing loss. Approximately 5 to 10% of earphone users are at risk of hearing loss and even permanent hearing loss if they listen to music for more than one hour a day at high volume [23]

Due to the length of earphone use of more than three years and daily use, earphones require attention, as literature indicates that prolonged exposure to loud noise can cause noise-induced hearing loss (NIHL) [25]. Research by Setianiet (2018) found that 77.8% of those with a history of noise exposure from earphones experienced NIHL, and 22.2% experienced NIHL with more than five years of use. Therefore, using earphones for five years or more can increase the risk of NIHL [26]. Noise-induced hearing loss (NIHL) is a form of sensory deficit that can lead to sensorineural hearing loss due to continuous exposure to loud noise over a prolonged period [27].

F. The Impact of Hearing Loss on Quality of Life

According to the World Health Organization (WHOQOL), quality of life is defined as an individual's perception of their life in society within the context of their culture and existing value systems, related to goals, expectations, standards, and concerns. Based on this definition, quality of life can be defined as life satisfaction or the fulfillment of life's needs based on a person's perceived physical, psychological, and social conditions [28], [29]. Factors that can influence a person's quality of life include psychological conditions. For example, psychological conditions can arise from the impact of an individual's physical limitations, which can lead to anxiety disorders, which can reduce the sufferer's ability to work and socialize with others. This can indirectly impact a person's quality of life. Beyond psychological factors, there are social relationship factors such as social support, which is the presence of others who can be relied upon to provide assistance, encouragement, acceptance, and attention, thereby improving and influencing the well-being or quality of life for the individual concerned [30]

Hearing loss can lead to a potential handicap or disability, limiting the individual's ability to perform various tasks in life. This can have psychological impacts, such as communication barriers, leading to anxiety, depression, and even impaired social interaction. Psychological and social relationships are factors that can impact a person's quality of life [31]

Material And Methods

This study uses a cross-sectional observational analytical research design, the researcher wants to prove the extent of the closeness of the relationship between a variable and another variable, where in this study the researcher wants to know the relationship

between earphone use patterns and the risk of hearing loss which can affect the quality of life of students of the Faculty of Medicine, Indonesian Christian University Class of 2019-2021.

This research was conducted from December to January 2023 at the Faculty of Medicine, Christian University of Indonesia, on Jl. Mayjen Sutoyo No. 2, RT. 2/RW. 11, Cawang Village, Kramat Jati District in East Jakarta City. The study population consisted of students from the Faculty of Medicine at Universitas Kristen Indonesia who graduated during the 2019-2021 intake. The sample consisted of all students who met the inclusion and exclusion criteria as in Table 1, and had given their consent.

Table 1. Inclusion and Exclusion Criteria

Criteria	Inclusion Criteria	Exclusion Criteria
Participants	Students from the 2019–2021 intake of UKI from any faculty.	Students who have previously been diagnosed with a hearing disorder (e.g., presbycusis, otosclerosis, chronic otitis media).
	Willing to participate and sign the informed consent form.	Students who are currently taking medication known to be ototoxic (e.g., certain antibiotics or chemotherapy agents).
	Own and regularly use earphones/headphones for leisure or study purposes.	Students who did not complete the entire questionnaire.

The sampling technique used in this study was non-probability sampling, namely, consecutive sampling, which was used to determine the research sample based on predetermined criteria and objectives. The research instrument consisted of two questionnaires: the Earphone Usage Pattern Questionnaire (EUPQ), which was used to examine respondents' habits of using earphones, and the Hearing Handicap Inventory for Adults (HHIA) questionnaire, which was used to determine the quality of life of earphone user respondents.

Data collected through the questionnaire were processed using IBM SPSS (Statistics for Social Science) and Microsoft Office Excel. Categorical data were displayed as percentages, while numerical data were displayed as means and standard deviations. All data were presented in tabular form. An inferential analysis was then performed to analyze the relationship between EUPQ and HHIA using the Chi-square test. The Pearson correlation test was used to analyze the relationship between EUPQ scores and HHIA, provided the data were normally distributed; otherwise, the Spearman test was employed.

Results

A. Respondent Profile

This study used an online questionnaire created by Google Forms and administered to students at the Faculty of Medicine, Universitas Kristen Indonesia, between December 2022 and January 2023. The aim of this study was to determine the relationship between earphone use patterns and the risk of hearing loss, which impacts quality of life, as measured by the Hearing Handicap Inventory for Adults (HHIA) questionnaire, adapted into Indonesian.

The study population comprised 225 students from the 2019, 2020, and 2021 intakes at the Faculty of Medicine, Universitas Kristen Indonesia.

The study lasted three months and was divided into two parts. In the first phase, participants were asked to complete a questionnaire about their regular earphone use patterns, and then they were asked to complete the HHIA questionnaire. The data was then analyzed, and a thesis was prepared.

Table 1: EUPQ Distribution Overview

Assessment Criteria		<i>f</i>	%
Length of Use of Earphones	0-3 years	52	23,1
	>3 years	173	76,9
Duration of Earphone Use	<1 hour	97	43,1
	1-2 hours	66	29,3
	>2-4 hours	36	16,0
	>4-6 hours	14	6,2
	>6-8 hours	5	2,2
	>8 hours	7	3,1
Frequency of Earphone Use	1-2 days	115	51,1
	3-4 days	62	27,6
	5-6 days	17	7,6
	Every Days	31	13,8
Earphone Usage Volume	<60%	140	62,2
	60-80%	69	30,7
	>80%	16	7,1
Earphone Usage Patterns	No Risk	85	37,8
	Risk	140	62,2
Total		225	100,0




Figure 1. The Chart of EUPQ Distribution Overview

Based on Table 1, 52 respondents used earphones for 0-3 years, representing 23.1% of the total, while 173 respondents used earphones for more than 3 years, representing 76.9% of the total. Based on these findings, the majority of students using earphones at the Faculty of Medicine, Universitas Kristen Indonesia (UKI) have used them since their teenage years, with some even using them since junior high school. A total of 97 respondents (43.1%) used earphones for more than 1 hour, 66 respondents (29.3%) used earphones for 1-2 hours, 36 respondents (16.0%) used earphones for more than 2-4 hours, 14 respondents (6.2%) for more than 4-6 hours, 5 respondents (2.2%) for more than 6-8 hours, and 7 respondents (3.1%) for more than 8 hours. From this table, it can be seen that students of the Faculty of Medicine, Christian University of Indonesia, Class of 2019, 2020, and 2021 mostly use earphones for more than one hour per day.

In total, 115 respondents (51.1%) reported using earphones 1-2 days a week, while 62 (27.6%) used them 3-4 days a week, 17 (7.6%) used them 5-6 days a week, and 31 (13.8%) used them daily. Here, we see that first-, second-, and third-year medical students at the Christian University of Indonesia used earphones between two and three days a week. 140 (62.2%) respondents used earphones with a volume of <60% (≤ 9 clicks of 0 volume), 69 (30.7%) used earphones with a volume of 60-80% (10-12 clicks of 0 volume), and 16 (7.1%) used earphones with a volume of >80% (>12 clicks of 0 volume). In this case, students at the Faculty of Medicine, Indonesian Christian University, Class of 2019, 2020, and 2021 mostly used earphones with a volume of <60% (≤ 9 clicks from 0% volume). 85 respondents (37.8%) had earphone usage

patterns that did not pose a risk, while 140 (62.2%) had earphone usage patterns that did. From what we can tell from this study, the majority of medical students at the Christian University of Indonesia (UI) in the 2019, 2020, and 2021 intakes are at risk of hearing loss. Heavy, long-term earphone use (>3 years), daily use (>1 hour), weekly use (>4 days), and high volume use (>60%) all contribute to risky earphone use.

Table 2: HHIA Score Results Overview

Indicator		<i>f</i>	%
Hearing Handicap Inventory Adults Score Results	No Handicap	167	74,2
	Mild Moderate	22	9,8
	Handicap		
	Severe Handicap	36	16
Total		225	100,0

Table 2 shows the scores from the Hearing Handicap Inventory for Adults (HHIA) questionnaire distributed to groups at risk for hearing loss. Of the respondents who completed the questionnaire, 167 (74.2%) had no handicap, 22 (9.8%) had a mild-moderate handicap, and 36 (16%) had a severe handicap.

Table 3: The relationship between EUPQ and HHIA

Earphone Usage Patterns	Hearing Handicap Inventory Adults (HHIA)						p-value
	No Handicap		Mild Moderate Handicap		Severe Handicap		
	<i>f</i>	%	<i>f</i>	%	<i>f</i>	%	
No Risk	71	83,5	8	9,4	6	7,1	0,015
Risk	96	68,6	14	10,0	30	21,4	
Total	167	74,2	22	9,8	36	16	

Table 3 shows that 96 respondents (68.6%) had earphone use patterns that put them at risk for hearing loss, 14 (10.0%) had mild-moderate handicaps, and 30 (21.4%) had severe handicaps. Meanwhile, 71 respondents (83.5%) were not at risk for hearing loss, 8 (9.4%) had mild-moderate handicaps, and 6 (7.1%) had severe handicaps. Statistical tests revealed a statistically significant relationship between risky and non-risky earphone use and handicaps that can affect respondents' quality of life, with a p-value of 0.015 ($p < 0.05$). Therefore, it can be concluded that there is a significant relationship between earphone use patterns and the risk of hearing loss. This relationship affects a person's quality of life. To determine the significance of the correlation and the degree of correlation between the independent and dependent variables, an analysis was conducted based on numerical data from the earphone usage pattern questionnaire and HHIA scores using the Pearson correlation test if the data were normally distributed and alternatively using the Spearmann correlation test if the data

were not normally distributed. A normality test was conducted to determine the normality of the data distribution, using the Kolmogorov-Smirnov test because the sample size was >50.

Table 4: Normality Test Results

Variable	Mean	SD	Median	Min	Max	<i>p-value</i>
Earphone Usage Patterns	9,48	3,24	9,0	4,0	18,0	<0,001^a
HHIA	16,14	21,42	6,0	2,0	96.0	<0,001

Description:

^aKolmogorov-Smirnov

Table 4 shows the results of the Kolmogorov-Smirnov test on earphone usage patterns, with a p value of <0.05 and HHIA of <0.05.

Table 5: Spearman Test Results

Variable	r	P-value
EUPQ	0,202	0,002 ^b
HHIA		

Description:

^bSpearman's Test

A Spearman's correlation test followed this. Table 4.5 shows a weak, significant relationship between earphone use patterns and the risk of hearing loss, which impacts students' quality of life.

B. Discussion

1. Distribution of Respondents' Earphone Usage Patterns

From the results of the research conducted by researchers, a questionnaire was completed to assess earphone usage patterns, which included information on duration, frequency, duration, and volume. It was found that the majority of respondents (173 respondents) had used earphones for more than three years, totaling 76.9%. This indicates that students from the Faculty of Medicine, Christian University of Indonesia (2019-2021) have been listening to music through earphones since they were in school. This aligns with research conducted by Fatimah Hamzah and Nurul, whose respondents were students from the Faculty of Medicine, Muhammadiyah University of Makassar, with an average duration of earphone use of more than three years (95.7%) [32]. According to the ENT-KL textbook, hearing loss due to prolonged exposure to excessively loud noise can lead to sensorineural hearing loss [33].

According to the World Health Organization (WHO), the largest number of earphone users worldwide are adolescents aged 12 to 24, who constitute part of the urban population [34].

The use of earphones has become commonplace in modern culture, especially among young people, who are more likely to use them as part of their daily routine than older generations. Earphones can be used as a means of listening to music and communicating because they can produce clearer sound than listening through the earphone speakers themselves. Furthermore, earphones are easy to carry anywhere, the volume can be adjusted, and they can provide privacy for the user. Regarding the duration of earphone use, the respondents in this study mostly used earphones for less than 1 hour, amounting to 97 people (47.1%). This result is in line with research by Martanegara and Ilman, who found that most respondents in the study used earphones for <1 hour, accounting for 61.34%. Noise-induced hearing loss can occur in the general public, especially teenagers, if they are often exposed to the use of earphones at high volume and continue to use them for an average of one to three hours per day [26]. More frequent use of earphones can be caused by the attachment to using earphones at high intensity because the use of earphones can be used in all things, such as when driving, sitting relaxing, doing activities, this earphone is very helpful so that even while doing activities, the person can still communicate or listen to music [22].

This study also found that 51.5% of the sample, or 115 students, from the Faculty of Medicine, Universitas Kristen Indonesia, Class of 2019, 2020, and 2021, mostly used earphones once or twice a week. A low frequency of earphone use can be determined when respondents wear their earphones from 1-2 days per week. This finding is consistent with that obtained from a study conducted in the past by Fatimah Hamzah and Nurul on students of the Faculty of Medicine, Universitas Muhammadiyah Makassar. The researchers found that almost half of the students (47.8%) used earphones for 1-2 days per week.

Based on the results of this study, as many as 140 respondents (62.2% of the total) often use earphone volume with > 60% on earphones. This is different when compared to research by Velaro, Adrian on Medical Faculty Students, University of North Sumatra, most of the respondents in their study used earphones with a volume of 60-80%. 26 Respondents who use earphones with high volume usually occur because users are not satisfied listening to them at low volume, so they choose to increase the volume of the sound produced by the earphones even though they are aware of the dangers of using earphones with high volume. This occurs even though respondents are aware of the dangers of using earphones with high volume [35]. If the use of earphones with high volume has become a habit for its users, there will be a trauma effect on the sound receptors due to the use of earphones with high volume intensity and cause a decrease in hearing function.

2. HHIA Score Overview

Information about hearing loss, expectations, and motivation can be collected using the Hearing Handicap Inventory for Adults (HHIA). The World Health Organization (WHO) defines disability or handicap as "a disadvantage to a person due to an impairment or impairment that limits or prevents the fulfillment of a normal role," taking into account the person's age, gender, and social and cultural context. The HHIA is a questionnaire used to assess the extent of handicap in adults with hearing loss. The HHIA is just one of many self-report instruments available to measure the functional impact of hearing loss on a person's daily life. HHIA results can be used to address some of the psychological impacts of hearing loss. The higher the HHIA score, the greater the recognition that hearing loss is a significant disability for the respondent due to the greater limitations or obstacles experienced by the sufferer in fulfilling their daily roles, resulting in a diminished quality of life.

This tool can assist audiologists by providing additional information regarding: (1) helping to substantiate complaints of hearing loss that are not apparent on conventional audiometric testing results; (2) facilitating candidate decisions regarding the use of amplification (hearing aids); (3) assisting in the counseling process; (4) serving as a guide for determining client-centered rehabilitation programs; and (5) serving as a measurement criterion in rehabilitation evaluations, including the benefits of post-surgical hearing aids.⁴⁶ Hearing loss can impact quality of life and limit activities or participation in daily life. The impact of this hearing loss can be assessed using the HHIA questionnaire [36]. In this study, the researchers used the HHIA questionnaire, which had been adapted into Indonesian and had been tested for validity and reliability.

The HHIA questionnaire in this study was administered to groups of people at and not at risk of developing hearing loss due to earphone use. 225 people (100%) completed the HHIA questionnaire. The results showed that 74.2% had no handicap, 9.8% had mild-moderate handicap, and 16% had severe handicap. In a previous study by Marcus and Dayse, the majority of respondents (70.8%) had severe handicap, 22.9% had mild-moderate handicap, and 6.3% had no handicap [37]. The HHIA questionnaire varies widely and depends on life experiences, health expectations, and even the individual's ability to adapt. People with hearing loss will experience communication difficulties and experience social and emotional difficulties in their daily lives. This will lead to different perceptions of their quality of life [36].

3. Relationship between Earphone Use Patterns and HHIA Scores

This study found a statistically significant relationship between earphone use patterns and the risk of hearing loss, which impacts the quality of life of students at the Faculty of Medicine, Christian University of Indonesia, in the 2019, 2020, and 2021 intakes ($p = 0.015$) of the 2019, 2020, and 2021 intakes ($p < 0.05$). Furthermore, the Spearman correlation test

revealed a weakly significant relationship between earphone use patterns and the risk of hearing loss, which impacted students' quality of life ($p = 0.002b$). Similarly, research by Marcus and Dayse confirmed that hearing loss can significantly impact quality of life [37].

The correlation between the two variables supports this finding. Incorrect use of earphones can lead to a risk of hearing loss and negatively impact quality of life, as measured by the HHIA questionnaire. Earphone use patterns can increase the risk of hearing loss, which can lead to physical, social, and emotional changes in individuals. These changes can persist and disrupt daily activities, significantly impacting quality of life, especially for those at risk for hearing loss.

The results demonstrating a higher HHIA score in respondents with risky earphone use patterns can be explained through well-established biological mechanisms. Exposure to sounds above 85 dB for long durations causes damage to outer hair cells (OHCs) in the cochlea. These OHCs function as mechanical amplifiers, improving frequency selectivity and sound sensitivity. High sound pressure levels lead to excessive mechanical vibration, which triggers oxidative stress and mitochondrial dysfunction in cochlear cells. Over time, this results in apoptosis and irreversible loss of sensory cells, leading to permanent threshold shifts and progressive hearing impairment.

Furthermore, loud music especially when transmitted directly into the ear canal via in-ear earphones causes temporary threshold shifts that, with repeated exposure, accumulate into permanent hearing loss. Research indicates that high-volume listening can induce excessive glutamate release, resulting in excitotoxicity at the ribbon synapses between inner hair cells and auditory nerve fibers. This "hidden hearing loss" may develop even before pure-tone audiometry detects deficits, and can manifest as difficulties in speech.

Conclusion

This study shows that students with risky earphone use patterns—such as long listening duration, high volume, and frequent weekly use—are more likely to experience hearing difficulties based on their HHIA scores. Although most students reported no handicap, a meaningful number already showed mild to severe hearing problems, indicating early signs of noise-induced hearing loss. Unsafe listening habits can gradually damage the cochlea and affect communication, concentration, and overall quality of life. Because many students have used earphones since adolescence, early prevention is important. To reduce the risk of hearing loss, educational institutions and related agencies should take more active steps, such as: 1) Providing education about safe listening habits (e.g., 60–60 rule); 2) Offering regular hearing screenings for students; 3) Creating campus policies that encourage healthy gadget use; 4) Running digital campaigns to raise awareness of safe earphone practices; 5) Partnering with health professionals to offer counseling for at-risk students. In summary, unsafe earphone use is a growing concern among young adults, and schools and health agencies should work together to promote safer listening habits and protect students' long-term hearing health.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- [1] C. J. Plack, *The sense of hearing*. Taylor and Francis, 2023. doi: 10.4324/9781003303329/SENSE-HEARING-CHRISTOPHER-PLACK.
- [2] W. H. Organization, *World report on hearing*. 2021. Accessed: Oct. 07, 2025. [Online]. Available: https://books.google.com/books?hl=en&lr=&id=zMRqEAAAQBAJ&oi=fnd&pg=PR5&dq=WHO.+World+Report+On+Hearing.+2021&ots=chAtI_6aMx&sig=5ks-DjL_goiQWico-iRullwvc-8
- [3] C. L. Nieman and E. S. Oh, "Hearing loss," *acpjournals.org* CL Nieman, *ES OhAnnals Intern. Med.* 2020•*acpjournals.org*, vol. 173, no. 11, pp. ITC81–ITC96, Dec. 2020, doi: 10.7326/AITC202012010.
- [4] B. Putri, R. Halim, H. N.-J. Kesmas, and undefined 2021, "Studi kualitatif gangguan pendengaran akibat bising/noise induced hearing loss (NIHL) pada marshaller di Bandar Udara Sultan Thaha Kota Jambi tahun 2020," *Mail. Putri, R Halim, HS NasutionJurnal Kesmas Jambi, 2021*•*mail.online-journal.unja.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://mail.online-journal.unja.ac.id/jkmj/article/view/12400>
- [5] N. Sarah, ... F. L.-J. (Jurnal K., and undefined 2016, "Hubungan Penggunaan Earphone Dengan Gangguan Pendengaran Pada Siswa Sma Negeri 9 Manado," *ejournal.unsrat.ac.id* NA Sarah, *F Lintong, JF RumampukJJK (Jurnal Kedokt. Klin. 2016*•*ejournal.unsrat.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.unsrat.ac.id/index.php/jkk/article/view/14363>
- [6] A. Budiarmanto, A. Asnawati, H. Huldani, ... S. K.-, and undefined 2024, "GAMBARAN TINGKAT PENGETAHUAN MAHASISWA PSKPS FK ULM ANGKATAN 2020 TENTANG FAKTOR RISIKO NOISE-INDUCED HEARING LOSS (NIHL)," *ppjp.ulm.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://ppjp.ulm.ac.id/journals/index.php/hms/article/view/13178>
- [7] P. Berliana, S. Dan, S. Keman, D. Kesehatan, L. Fakultas, and K. Masyarakat, "Pengaruh Penggunaan Pelindung Telinga dan Earphone Terhadap Noise Induced Hearing Loss Dan Tinitus Pada. Pekerja Bengkel," *e-journal.unair.ac.id* PB Syah, *S KemanJurnal Kesehat. Lingkungan, 2017*•*e-journal.unair.ac.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://e-journal.unair.ac.id/JKL/article/download/9148/5147>

- [8] "Susiyanti E, Imanto M. Efek Penggunaan Earphone sebagai... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Susiyanti+E%2C+Imanto+M.+Ef+ek+Penggunaan+Earphone+sebagai+Faktor+Resiko+Kejadian+Noise+Induced+Hearing+Loss.+Majority.+2020%3B+9%282%29%3A+63-67.+&btnG= (accessed Oct. 07, 2025).
- [9] E. Yuliyani, D. Yudhanto, ... W. A.-I. J., and undefined 2022, "Peranan program acara stasiun TVRI NTB sebagai salah satu media edukasi kesehatan pendengaran," *Acad. Yuliyani, D Yudhanto, WS Aff. E Triani, F Libr. DAIS Prajnyaswari*INDRA J. Pengabdi. Kpd. Masyarakat, 2022•*academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/93056133/125.pdf>
- [10] L. Moneta, L. Q.-D.-O.-H. and N. Surgery, and undefined 2017, "Embryology and anatomy of the ear," *Elsevier*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S1043181017300398>
- [11] R. Maynard and N. Downes, *Anatomy and histology of the laboratory rat in toxicology and biomedical research*. 2019. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=2kGHDwAAQBAJ&oi=fnd&pg=PP1&dq=Maynard+RL,+Downes+N.+The+Ear.+In:+Anatomy+and+Histology+of+the+Laboratory+Rat+in+Toxicology+and+Biomedical+Research+%5BInternet%5D.+Elsevier%3B+2019.+p.+293-302.+Available+from:+https://linkinghub.elsevier.com/retrieve/pii/B978012811837500023X&ots=-Iqlh67SG&sig=eYuR9X9RkEvkTDRjfkG-C7whl70>
- [12] A. Szymanski and Z. Geiger, "Anatomy, head and neck, ear," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/article/nbk/nbk470359>
- [13] A. Szymanski, J. Toth, M. Ogorevc, and Z. Geiger, "Anatomy, head and neck, ear tympanic membrane," 2017, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk448117>
- [14] "Sánchez López de Nava A, Lasrado S. Physiology, Ear. 2022 - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Sánchez+López+de+Nava+A%2C+Lasrado+S.+Physiology%2C+Ear.+2022&btnG= (accessed Oct. 07, 2025).
- [15] H. White, M. Helwany, ... A. B.-S., and undefined 2023, "Anatomy, head and neck, ear organ of corti," *ncbi.nlm.nih.gov*HJ White, M Helwany, AR Biknevicius, DC PetersonStatPearls [Internet], 2023•*ncbi.nlm.nih.gov*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.ncbi.nlm.nih.gov/books/NBK538335/>
- [16] R. Sataloff and P. Roehm, *Hearing loss*. 2024. Accessed: Oct. 07, 2025. [Online]. Available: <https://books.google.com/books?hl=en&lr=&id=IG0iEQAAQBAJ&oi=fnd&pg=PP1&dq=Hearing+Loss&ots=Y1HvwBOzEA&sig=EjaY-3-SDMk5upL0768pSMieSHE>
- [17] T. Sooriyamoorthy and O. De Jesus, "Conductive hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk563267>
- [18] R. Tanna, J. Lin, and O. De Jesus, "Sensorineural hearing loss," 2020, Accessed: Oct. 07, 2025. [Online]. Available: <https://europepmc.org/books/nbk565860>
- [19] S. C.-C. E. and H. Health and undefined 2019, "Noise-induced hearing loss," *account.cehh.press.lshtm.ac.uk*S ChadhaCommunity Ear Hear. Heal. 2019•*account.cehh.press.lshtm.ac.uk*, Accessed: Oct. 07, 2025. [Online]. Available: <https://account.cehh.press.lshtm.ac.uk/index.php/s-j-cehh/article/view/31>
- [20] "Chan Y, Goddard JC. Preface. In: KJ Lee's Essential... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Chan+Y%2C+Goddard+JC.+Pref+ace.+In%3A+KJ+Lee's+Essential+Otolaryngology%3A+Head+and+Neck+Surgery%2C+12e+%5BInternet%5D.+New+York%2C+NY%3A+McGraw-Hill+Education%3B+2019.+Available+from%3A+accesssurgery.mhmedical.com%2Fcontent.aspx%3Faid%3D1172366487&btnG= (accessed Oct. 07, 2025).
- [21] L. Cunningham, D. T.-N. E. J. of Medicine, and undefined 2017, "Hearing loss in adults," *Mass Med. SocLL Cunningham, DL TucciNew Engl. J. Med. 2017•Mass Med. Soc*, vol. 377, no. 25, pp. 2465-2473, Dec. 2017, doi: 10.1056/NEJMRA1616601.
- [22] M. Erlanda, P. Negara, I. Triansyah, D. Hasni, and B. Y. Febrianto, "Hubungan Intensitas Penggunaan Earphone dengan Derajat Gangguan Pendengaran pada Mahasiswa Fakultas Kedokteran Universitas Baiturrahmah," *journal.scientic.id*MEP Negara, I Triansyah, D Hasni, BY FebriantoScientific Journal, 2022•*journal.scientic.id*, Accessed: Oct. 07, 2025. [Online]. Available: <https://journal.scientic.id/index.php/sciena/article/view/49>
- [23] A. Velaro, D. Z.-O. R. L. Indonesiana, and undefined 2021, "The correlation between earphone usage patterns with the incidence rate and severity of tinnitus," *orli.or.id*AJ Velaro, D Zahar. Rhino

- Laryngol. Indones.* 2021•orli.or.id, Accessed: Oct. 07, 2025. [Online]. Available: <http://orli.or.id/index.php/orli/article/view/432>
- [24] M. Zhang, A. Jeske, S. Y. (yrs), and undefined 2017, "Changes in the frequency and intensity of Tinnitus using the Suppressive Noise Spectrum," *m.karmaimpex.com* M Zhang, A Jeske, S Youngage (yrs), 2017•*m.karmaimpex.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://m.karmaimpex.com/hjed/pdf/ated-aid1002.pdf>
- [25] B. Rosita, "Pemakaian Earphone dalam Mendengar Musik yang Menimbulkan Bising di Kalangan Milenial," 2019, Accessed: Oct. 07, 2025. [Online]. Available: https://osf.io/ascomb_v1/
- [26] L. Setiani, N. Syakila, Y. Y.-J. K. N. Medika, and undefined 2018, "Hubungan Lama Paparan Penggunaan Earphone Musik Terhadap Terjadinya Gangguan Pendengaran Akibat Bising pada Mahasiswa Fakultas Kedokteran," *jknamed.com* L Setiani, N Syakila, Y Yusni *Jurnal Kedokt. Nanggroe Med.* 2018•*jknamed.com*, Accessed: Oct. 07, 2025. [Online]. Available: <http://www.jknamed.com/jknamed/article/view/51>
- [27] T. Wang, T. Chang, R. Tyler, Y. Lin, W. L.-B. sciences, and undefined 2020, "Noise induced hearing loss and tinnitus—new research developments and remaining gaps in disease assessment, treatment, and prevention," *mdpi.com* TC Wang, TY Chang. R Tyler, YJ Lin, WM Liang, YW Shau, WY Lin, YW Chen, CD Lin *Brain Sci.* 2020•*mdpi.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.mdpi.com/2076-3425/10/10/732>
- [28] "Pratiwi Y. Pengaruh Dukungan Sosial Terhadap Kualitas... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Pratiwi+Y.+Pengaruh+Dukungan+Sosial+Terhadap+Kualitas+Hidup+Lanjut+Usia+di+Pusar+Santunan+Keluarga+%28PUSAKA%29+Kecamatan+Pancoran+Jakarta+Selatan+%5Bskripsi%5D.+Jakarta%3A+Fakultas+Dakwah+dan+Komunikasi+Universitas+Islam+Negeri+Syarif+Hidayatullah%3B+2017&btnG= (accessed Oct. 07, 2025).
- [29] "WHO. WHOQOL User Manual Programme On Mental Health... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=WHO.+WHOQOL+User+Manual+Programme+On+Mental+Health+Division+Of+Mental+Health+And+Prevention+Of+Substance+Abuse+World+Health+Organization.+2020&btnG= (accessed Oct. 07, 2025).
- [30] D. Maya *et al.*, "Kualitas hidup lansia ditinjau dari sabar dan dukungan sosial," *ejournal.umm.ac.id* DMP Sari, CYD Lestari, EC Putra, F Nashori *Jurnal Ilm. Psikol. Ter.* 2018•*ejournal.umm.ac.id*, vol. 06, no. 02, pp. 2540–8291, 2018, Accessed: Oct. 07, 2025. [Online]. Available: <https://ejournal.umm.ac.id/index.php/jipt/article/view/5341>
- [31] A. Nurfaizi, "MENINGKATKAN KEMAMPUAN DETEKSI DINI DAN PENATALAKSANAAN GANGGUAN PENDENGARAN DI KALIBARU KULON BANYUWANGI," *academia.edu*, Accessed: Oct. 07, 2025. [Online]. Available: <https://www.academia.edu/download/108530358/14428.pdf>
- [32] "Hamzah NF. Pengaruh Perilaku Penggunaan Earphone... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=+Hamzah+NF.+Pengaruh+Perilaku+Penggunaan+Earphone+Terhadap+Gangguan+Telinga+Pada+Mahasiswa+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar+%5Bskripsi%5D.+Makassar%3A+Fakultas+Kedokteran+Universitas+Muhammadiyah+Makassar%3B+2021&btnG= (accessed Oct. 07, 2025).
- [33] "Soepardi EA, Iskandar N, Bashiruddin J, Restuti RD.... - Google Scholar." https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Soepardi+EA%2C+Iskandar+N%2C+Bashiruddin+J%2C+Restuti+RD.+Buku+Ajar+Ilmu+Kesehatan%3A+Telinga%2C+Hidung%2C+Tenggorok%2C+Kepala+%26+Leher.+7th+ed.+Jakarta%3A+BP+FKUI%3B+2012.+&btnG= (accessed Oct. 07, 2025).
- [34] N. Yudiasari, S. W.-P. J. K. Masyarakat, and undefined 2021, "Pengembangan Media Promosi Kesehatan Tentang Bahaya Penggunaan Earphone Mahasiswa Universitas Airlangga Di Surabaya," *103.245.72.38* N Yudiasari, S Widati *Preventif J. Kesehat. Masyarakat*, 2021•*103.245.72.38*, vol. 12, pp. 57–86, 2021, Accessed: Oct. 07, 2025. [Online]. Available: <http://103.245.72.38/index.php/preventif/article/view/182>
- [35] S. Levey, B. Fligor, ... C. G.-C. I. in, and undefined 2012, "The effects of noise-induced hearing loss on children and young adults," *pubs.asha.org* S Levey, BJ Fligor, C Ginocchi, L Kagimbi *Contemporary Issues Commun. Sci. Disord.* 2012•*pubs.asha.org*, vol. 39, no. Fall, pp. 76–83, Oct. 2012, doi: 10.1044/CICSD_39_F_76.
- [36] J. Bashiruddin, W. Alviandi, B. Bramantyo, and R. Ismail, "Praktik Klinis: Hearing Handicap Inventory For Adults Adaptasi Bahasa Indonesia Sebagai Alat Ukur Kualitas Hidup Pasien Gangguan Pendengaran," 2019, Accessed: Oct. 07, 2025. [Online]. Available: <https://scholar.ui.ac.id/en/publications/buku-panduanpetunjuk-panduan-praktik-klinis-hearing-handicap-inve>

- [37] M. Voola, D. T.-V.-T. M. Journal, and undefined 2020, "Quality of life handicap measured in patients with profound unilateral or bilateral deafness," *tasmanmedicaljournal.com* M Voola, D Távora-Viera *Tasman Med. Journal*, 2020 • *tasmanmedicaljournal.com*, Accessed: Oct. 07, 2025. [Online]. Available: <https://tasmanmedicaljournal.com/2020/12/quality-of-life-handicap-measured-in-patients-with-profound-unilateral-or-bilateral-deafness/>

Authors



Lina Marlina     is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. She is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery. She is a member of the Indonesian Medical Association (IDI). Her professional interests include ENT and ENT and clinical research. She is actively involved in academic research and public health programs. She can be contacted at lina.marlina@uki.ac.id.



Sena Pongtuluran     is a general practitioner who recently graduated from the Faculty of Medicine, Christian University of Indonesia, Jakarta. She is currently pursuing a Master's degree in Hospital Administration (MARS) at Pelita Harapan University (UPH). Her interests include clinical medicine, community health, and healthcare management. She is passionate about contributing to healthcare innovation and evidence-based medical practice. She can be contacted at senapongtuluran@gmail.com



Fransiskus Harf Poluan     is a lecturer in the Department of Otolaryngology and ENT, Faculty of Medicine, Christian University of Indonesia, Jakarta. He is also a specialist practicing in the field of ear, nose, throat, and head and neck surgery at Mayapada Hospital, Kuningan, Jakarta. He is actively involved in academic research and public health programs. He can be contacted at fransiskus.poluan@uki.ac.id

[← Back to Submissions](#)

Revisions Required



Participants

- Vikas Gupta (vikas_gupta)
- Lina Marlina (lina25)

Messages

Note	From
Dear Authors, Please also attach point-by-point responses addressing each reviewer's comments (refer to the template here). We are looking forward to receiving your revised manuscript.	vikas_gupta 2025-11-25 12:44 PM UTC
▶ Dear Editor, Below we sent the file of the response to the reviewer. Thank you 📄 IISTR Response to Reviewer_LIna Marlina.doc	lina25 2025-12-16 07:03 AM UTC

Original Manuscript ID: XXXX

Original Article Title: “Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life”

To: IISTR Editorial Office

Re: Response to reviewers

Dear Editor,

Thank you for allowing a resubmission of our manuscript, with an opportunity to address the reviewers' comments.

We are uploading (a) our point-by-point response to the comments (below) (response to reviewers), (b) an updated manuscript with yellow highlighting indicating changes (pdf), and (c) a clean updated manuscript without highlights (*.docx main document/LaTeX files).

Best regards,

Lina Marlina, et al.

Reviewer#1, Concern # 1: for Introduction: While informative, the introduction could be more focused by reducing some overly technical information about ear anatomy to more directly address the research objectives.

Author response:

Author action: We updated the manuscript by revision on the pages 2 & 3

Reviewer#1, Concern # 2: The use of earphones could be clarified as a primary risk factor in the context of this study to make it more relevant to the research focus.

Author response:

Author action: We updated the manuscript by revision on the pages 2 & 4

Reviewer#1, Concern # 3: The method used was a cross-sectional observational study design with consecutive sampling. The authors collected data using two questionnaires: the Earphone Usage Pattern Questionnaire (EUPQ) and the Hearing Handicap Inventory for Adults (HHIA). Data analysis was performed using the Chi-square test and Spearman's correlation test.

Strengths:

1. The study design used was appropriate for identifying the relationship between earphone usage patterns and hearing loss.
2. The selection of two relevant questionnaire instruments (EUPQ and HHIA) supported the study's objective of measuring earphone usage patterns and their impact on quality of life.

Recommendations for Revision:

1. The sample used was students from the Faculty of Medicine, which could be biased because it is not representative of the general population. The authors recommend adding more sample variation to increase the generalizability of the study results.
2. Further explanation of the inclusion and exclusion criteria would improve the transparency of the research method.

Author response:

Author action: For the inclusion and exclusion criteria, we have revised and updated them as stated in the method section on page 8. However, we apologize, but it seems we cannot yet meet the requirements for increasing sample variation. We hope this will be a concern for our future research.

Reviewer#1, Concern # 4: The results showed that 62.2% of respondents used earphones in a risky manner, while 37.8% did not. Based on the results of the HHIA questionnaire, the majority of respondents (74.2%) did not experience hearing loss, but 16% of respondents experienced moderate to severe hearing loss. Statistical tests showed a significant relationship between risky earphone use patterns and hearing loss that impacted quality of life ($p = 0.015$).

Strengths:

1. The study results are presented clearly through tables and statistics that support the conclusion that risky earphone use patterns are associated with hearing loss that impacts quality of life.
2. The presentation of data through tables helps readers easily understand the distribution of respondents and the questionnaire results.

Revision Suggestions:

1. Although the data are presented clearly, the presentation of the results could be supplemented with graphs that more visually illustrate the relationship between variables.
2. The author could add more detailed information regarding respondent characteristics, for example, age or gender, which may have influenced the study results.

Author response:

Author action: We updated the manuscript by revision on page 10

Reviewer#1, Concern # 5: This article's discussion links the research findings to previous studies, highlighting the importance of earphone usage patterns in increasing the risk of hearing loss. The author also explains how factors such as the duration and volume of earphone use contribute to the risk of hearing loss. This discussion emphasizes the importance of awareness of the negative impacts of unwise earphone use.

Strengths:

1. A fairly robust discussion, linking the research findings to existing literature, provides readers with a better understanding of the topic.
2. The author also cites several other studies that support the findings of this study, enriching the discussion.

Revision Suggestions:

1. The discussion could be more in-depth by further exploring the biological mechanisms involved in listening to music at high volume through earphones and how this affects hearing in the long term.
2. A discussion of other factors influencing quality of life, such as social support or mental health, which may interact with the results of this study could also be added.

Author response:

Author action: We updated the manuscript by revision on page 15

Reviewer#1, Concern # 6: The conclusion could be stronger by adding further recommendations regarding policies or interventions that can be implemented by educational institutions or related agencies to reduce the risk of hearing loss due to unsafe earphone use.

Author response:

Author action: We updated the manuscript by revision on page 16

Note: *References suggested by reviewers should only be added if it is relevant to the article and makes it more complete. Excessive cases of recommending non-relevant articles should be reported to editor@journal.iistr.org*

Missing Revision and Response to Reviewer 2



Participants

Vikas Gupta (vikas_gupta)

Lina Marlina (lina25)

Messages

Note	From
Dear Authors,	vikas_gupta
Thank you for submitting the revised version of your manuscript.	2025-12-26 12:12 PM UTC
After carefully checking the resubmission, we note that the revision does not address the comments provided by Reviewer 2 , and we could not find a point-by-point response to Reviewer 2's comments in the response-to-reviewers document.	
We kindly ask you to revise the manuscript accordingly and resubmit it together with a complete response to Reviewer 2.	
If you have any questions regarding the reviewer comments or the revision process, please feel free to contact us.	
Best regards	

[Add Message](#)

[← Back to Submissions](#)1440 / **Marliana et al.** / Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life[Library](#)

Workflow

Publication

Submission

Review

Copyediting

Production

Copyediting Discussions

[Add discussion](#)

Name	From	Last Reply	Replies	Closed
▶ Proof Payment Article Code 1440_Lina Marliana	lina25 2026-03-05 07:22 AM UTC	-	0	☐
[JPHS] Post-Acceptance Requirements for Your Manuscript	editor3 2026-03-11 06:02 AM UTC	-	0	☐
▶ Turnitin Check Results Document, Ethics Review and Inform Concern	lina25 2026-03-25 04:04 AM UTC	-	0	☐
[JPHS] Revisi Perbaikan Reference	editor3 2026-04-06 05:52 AM UTC	lina25 2026-04-06 11:50 AM UTC	1	☐
▶ File Revisi Referensi	lina25 2026-04-06 11:53 AM UTC	-	0	☐

Copyedited

[Q Search](#)



13476

JPHS_5_2_5.docx

May
23,
2026

Article Text



13521

1440-Article Text-10972-13476-9-20260523.docx

May
25,
2026

Article Text



← Back to Submissions

1440 / **Marliana et al.** / Earphone Use Patterns and Hearing Loss Risk Affecting Quality of Life

Library

Workflow

Publication

Submission

Review

Copyediting

Production

Production Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
[JPHS] Congratulations - Your article is now available online!	vikas_gupta 2026-05-23 01:25 PM UTC	-	0	☐