

# The Effect of Sperm DNA Fragmentation on Fertilization and Embryogenesis Rates

*by* Batara Imanuel Sirait

---

**Submission date:** 06-Oct-2020 01:20PM (UTC+0700)

**Submission ID:** 1406801196

**File name:** 69\_2.\_SST\_Batara\_Imanuel\_Sirait.pdf (1.49M)

**Word count:** 5428

**Character count:** 30541

## The Effect of Sperm DNA Fragmentation on Fertilization and Embryogenesis Rates

Batara Imanuel Sirait

Department of Obstetrics and Gynecology, Medical Faculty, Universitas Kristen Indonesia  
Corresponding Email: batara.sirait@uki.ac.id

### Abstract:

This research is the effects of sperm DNA fragmentation on fertilization and embryogenesis rates. The purpose of doing this research is to find out the effect of sperm DNA fragmentation on fertilization and embryogenesis rates. It is a quantitative study with cross-sectional design and carried out by taking male semen samples from infertile couples who participated in the IVF-ICSI program, examined WHO sperm analysis, and assessed the sperm DNA fragmentation index. For the analysis of the mean difference of 2 categories using independent sample t-tests while those of more than 2 categories use Anova. Analysis of categorical variable types uses the chi-square test. The result of this research shows that In the 10-month study period, the number of study subjects who met the inclusion criteria was 29 infertile couples with 29 ejaculate sperm samples and 200 oocytes. From the analysis of sperm DNA fragmentation index with fertilization rates, it appears that good fertilization (Z1 and Z2) is more common in groups with sperm DNA fragmentation index  $\leq 30\%$ , and statistically significantly different with  $p < 0.05$ , and from analysis of sperm DNA fragmentation index with embryo scoring seen better embryo scoring (grades A and B) in the sperm DNA fragmentation index group  $\leq 30\%$ , which was not statistically significant. It is concluded that the sperm DNA fragmentation rates can show a negative relationship with fertilization rates but can not show a statistically significant relationship with embryogenesis rates. A statistically significant relationship was seen between the World Health Organization (WHO) sperm analysis parameters and DNA fragmentation index examined using the SCD method.

**Keywords:** Sperm DNA fragmentation index, Fertilization, Embryogenesis.

## 1. INTRODUCTION

Infertility can be caused by both female and male or both factors. It is a complex situation, where often a combination of several factors plays an important role. Traditionally genital infections, endocrine disorders, and immunological factors are considered as the most common causes of male subfertility. However, at this time often genetic/molecular causes are found as contributing factors, such as chromatin damage which is considered as defragmentation of sperm DNA. However, 60-75% of cases of decreased semen quality cannot be explained and are therefore diagnosed as idiopathic infertility. The gold standard of diagnosis of male infertility or subfertility is sperm analysis which includes concentration, motility, and morphology according to standards determined by WHO. Over the past decade several sperm function tests have been recommended, including vital staining, hemizona assay, biochemical analysis of semen, anti-sperm antibody test, hypoosmotic swelling test, sperm penetration assay, reactive oxygen species (ROS)

tests, and computer-assisted sperm analysis (CASA) [1;2;3].

Several examination techniques have been proposed to study the disorder. Some [33] are still used today are the techniques of TUNEL, Comet, orange acridine, and sperm chromatin structure assay (SCSA). Furthermore, there is also an examination of Sperm Chromatin Dispersion (SCD) which turned out to be a fair [17] accurate examination to assess sperm DNA fragmentation. This examination is based on the presence or absence of halo by giving acidic liquid to sperm which is then followed by lysis buffer, halo DNA dispersion can be seen in the sperm nucleus with unfragmented DNA after the loss of core proteins and on the req [11] with fragmented DNA, this halo looks small or not formed at all [4;5;6;7]. Our understanding of the female reproductive function and the importance [11] of male-factors in infertility have increased significantly over the past decade. In the past, female partners were the main focus of attention and the husband factor was considered a relatively unusual [24] cause of infertility. It is now known that abnormalities in men can be the sole cause of infertility in approximately 20% of infertile couples and play an important role in 20–40% of couples with reproductive disorders [8; 9; 10;11].

The effects [38] of sperm DNA fragmentation on infertility have been the subject of several studies. Previously 10-20% sperm DNA fragmentation has been reported in ejaculate sperm. Infertile men with poor motility and morphology have been suspected to have increased sperm DNA fragmentation when compared to men with normal sperm parameters. Men with normal sperm parameters can also have high sperm [5] DNA fragmentation, which can be an unexplained reason for infertility [12;13;14]. Aberrant chromatin packaging during spermatogenesis, defective apoptosis [3] before ejaculation, or overproduction of reactive oxygen species (ROS) causes sperm DNA fragmentation, but the [46] mechanism underlying the situation is unclear. There is still controversy over the effects of sperm DNA damage on reproductive output. Some examiners have found that clinical pregnancy is inversely affected by damage to sperm DNA in ICSI cases. Moreover, fertilization obtained through sperm which has DNA fragmentation can cause poor embryonic development, reduced implantation rates, and pregnancy. Another examine [6] found that sperm DNA damage caused ineffectiveness in fertilization, decreased embryo quality and pregnancy rates in IVF and ICSI [15; 16; 17; 18].

[1] Many studies have shown that [1] paternal effects can cause the repeated failure of assisted reproductive techniques. Many authors have shown that the paternal effect can be traced to the anomalous sperm chromatin organization. Sperm from [45] infertile men are characterized by being susceptible to acid-induced denaturation, reduced chromatin condensation, chromosomal anomalies, and/or increased DNA fragmentation. Many sperm seen with changes in the structure of chromatin [1] harms the outcome of the TRB procedure. These studies focus on examining possible correlations between changes in paternal chromatin and fertilization rates, embryo division, blastocyst development, and clinical pregnancy in both IVF and ICSI [19;20]. These findings suggest that paternal genomic changes can jeopardize fertilization, embryonic quality as well as embryonic viability [1] and continuity of pregnancy, leading to spontaneous abortion. At present, several studies in men and animals have underlined the importance of paternal factors, including male age or exposure to toxic material in cases of spontaneous abortion, but the relationship between early embryonic development after implantation in couples undergoing

TRB and examination of sperm DNA integrity is still a necessary further explanation. The effect of changes in sperm chromatin integrity on the early development of embryos after implantation is still debated [21; 22].

Mammalian fertility and subsequent embryonic development depend in part on the integrity of sperm DNA. It even seems that there is a threshold level of damage to sperm DNA (DNA fragmentation, abnormal chromatin, and protamine deficiency) wherein above that the rate of embryonic development and pregnancy can be disrupted. Sperm integrity testing has been developed and clinically applied. Sperm DNA fragmentation is increasingly understood as an important cause of infertility. Recent clinical research states that levels of sperm DNA fragmentation above 30% as measured by the SCSA method are incompatible with the initiation and occurrence of term pregnancy. Recent data shows that the value of the SCSA examination called the DNA Fragmentation Index correlates significantly with pregnancy rates both invivo and invitro. All pregnancies occur when the DNA Fragmentation Index number is less than 30% [23]. SCD (sperm chromatin dispersion) was not created as a substitute for SCSA. However, like SCSA, SCD can also distinguish sperm DNA fragmentation. The results of examining sperm DNA fragmentation by the SCD method are also consistent when compared with the results of examinations using the SCSA method. This shows that even with much simpler equipment and procedures an SCD examination is fast, relatively inexpensive, accurate, and can be repeated with an accuracy comparable to an SCSA examination. Therefore, SCD examination has the potential to be used as a routine screening test for sperm DNA fragmentation [24;25].

This result was confirmed by DNA breakage detection-fluorescence in situ hybridization (DBD-FISH), a procedure that uses restricted single-stranded DNA motifs resulting from DNA damage that can be detected and calculated. So DNA fragmentation that is reflected by the halo size can be determined accurately using SCD examination which is a simple, accurate, well-repeated, and inexpensive technique. In the SCD protocol, sperm nucleoids can be seen using fluorosens microscope after staining with DNA specific fluorochrome (6-diamino-2-phenylindole [DAPI]) or with a light microscope after Diff-Quik staining. The ease of use and interpretation of SCD, as well as sensitive and reliable techniques, has the potential to be carried out in a variety of basic studies in clinical laboratories, in contrast to the more complex SCSA [26;27].

## 2. METHOD

This research is observational research, by taking semen samples of infertile men who participated in the IVF-ICSI program, examined WHO sperm analysis and assessment of sperm DNA fragmentation index. After ICSI, fertilization and embryogenesis were assessed. Then analyze by calculating the correlation. The study design was cross-sectional to see the correlation between the sperm DNA fragmentation index with the rate of fertilization and embryogenesis. Inclusion criteria were: Sperm of patients who participated in IVF using the ICSI method at Permata Hati Clinic Dr. Sardjito Yogyakarta who is willing to take part in the research. Exclusion Criteria: Couples who participated in IVF with the ICSI method at Permata Hati Clinic Dr. Sardjito Yogyakarta with her woman suffering from polycystic ovaries and or endometriosis. The sample size is Proportion

Sample, the patient who came to take the IVF program with the ICSI method at Permata Hati Clinic Dr. Sardjito Yogyakarta from July 20<sup>20</sup> to May 2013. Researchers have obtained information on ethical eligibility (Ethical Clearance) from the Biomedical Research Ethics Commission in Humans, Faculty of Medicine, Gadjah Mada University Yogyakarta. Intact spermatozoa are dissolved in agarose matrix on the slide, given an acid solution for denaturation of DNA that contains damage, then lysis buffer is given to remove the membrane and protein. Agarose matrices make it possible to work without fixed sperm on an object-glass in an environment such as a suspension. The removal of the nucleus protein produces nuclei with central nuclei and peripheral halo from the dispersed DNA strands. By using fluorosens staining, the obtained sperm nucleus with increased DNA fragmentation produces a very small halo or absence of halo from DNA dispersion, where sperm with a low fragmentation level releases DNA strands that will form large halos.




Figure 1. Flow Check of Sperm Chromatin Dispersion

The assessment of sperm DNA fragmentation examination<sup>22</sup> results using the sperm chromatin dispersion method is done by looking at the size of the halo that forms around the nucleus of sperm cells.






Figure 2. Assessment of Sperm DNA Fragmentation by SCD Method

Scott et al use the Z-scoring system to divide zygotes into four categories based on an assessment of the size and alignment of the nucleus and also the number and distribution of nucleoli [28].



**Figure 3. Z-scoring system [28]**

Baczkowski T et al used a scoring system as below to divide embryos into four groups in which embryos included in groups A and B were considered good embryos, while those in groups C and D considered embryos that were considered to be unfavourable with the risk of developing congenital defects if an embryo transfer is done.



**Figure 4. Morphological assessment of third-day embryos [28]**

### 3. RESULT AND DISCUSSION

In our 10-month study period, the number of study subjects after excluding patients with a diagnosis of endometriosis and PCOS according to exclusion criteria was 29 infertile couples with 29 ejaculate sperm samples and 200 oocytes. The statistical test used is the parametric test. For the analysis of the mean difference of 2 categories using an independent sample t-test while those of more than 2 categories use ANOVA. Analysis of categorical variable types uses the chi-square test.

<sup>22</sup>  
**Table 1. Results of Analysis of Sperm DNA Fragmentation Index by examination of WHO Sperm parameters**

Variable	Fragmentation Index Sperm DNA	
	≤30% N (%)	>30% N (%)
<b>WHO Sperm Analysis</b>		
Normozoospermia	28 (23.1%)	0
Oligozoospermia	5 (4.1%)	11 (14.1%)
Asthenozoospermia	34 (28.1%)	20 (25.6%)
Oligoasthenozoospermia	3 (2.5%)	0
Teratozoospermia	20 (16.5%)	14 (17.9%)
Oligoasthenoteratozoospermia	31 (25.6%)	33 (42.3%)

**P<0.00**

From Sperm DNA Fragmentation Index Analysis with WHO Sperm Analysis seen in sperm DNA fragmentation index group indeks30% is dominated by WHO sperm analysis results which are *asthenozoospermia*, *oligoasthenoteratozoospermia*, *normozoospermia*, and *teratozoospermia*. In groups>30%, it is dominated by *oligoasthenoteratozoospermia*, *asthenozoospermia*, *teratozoospermia*, and <sup>36</sup> followed by *normozoospermia*. These results were statistically seen that there were significant differences  $p < 0.05$ .

<sup>7</sup>  
**Table 2. Results of Sperm DNA Fragmentation Index Analysis with Fertilization Rates**

Variable	Fertilization Figures				
	Negative	Z1	Z2	Z3	Z4
Sperm DNA Fragmentation Index					
≤30%	59 (60.2%)	9 (81.8%)	48 (60.8%)	1 (14.3%)	4 (100.0%)
>30%	39 (39.8%)	2 (18.2%)	31 (39.2%)	6 (85.7%)	0

**P=0.027**

<sup>48</sup>  
From the Analysis of Sperm DNA Fragmentation In <sup>43</sup> with Fertilization Figures, it appears that good fertilization (Z1 and Z2) is more in the group with sperm DNA fragmentation index ≤30%, but there are conflict <sup>42</sup> results where negative fertilization was also found more in the group with sperm DNA fragmentation index ≤30%. The results of the analysis of the table above between the DNA fragmentation index <sup>41</sup> with the fertilization rate obtained statistically significantly different results  $p < 0.05$ .

**Table 3. Results of Analysis of Sperm DNA Fragmentation Index by Embryo Scoring**

Variable	Skoring Embriogenesis				
	Negative	A	B	C	D
Sperm DNA Fragmentation Index					
≤30%	65 (59.1%)	7 (87.5%)	24 (55.8%)	24 (66.7%)	1 (50.0%)
>30%	45 (40.9%)	1 (12.5%)	19 (44.2%)	12 (33.3%)	1 (50.0%)

**P=0.464**

Analysis of Sperm DNA Fragmentation Index with Embryo Scoring shows better embryo scoring (Grade A and B) in the sperm DNA fragmentation index group ≤30%, but there are conflicting results where negative embryogenesis is also found more in groups with index sperm DNA fragmentation ≤30%. The results of the above table analysis between DNA fragmentation index and Embryo Scoring obtained results that were not statistically significant  $p > 0.05$ .

**Table 4. Analysis results of examining WHO Sperm parameters with Fertilization Rates**

Variable	Fertilization				
	Negative	1	2	3	4
WHO Sperm Analysis					
Normozoospermia	8 (8.2%)	(36.4%)	6 (20.3%)	0	0
Oligozoospermia	5 (5.1%)	0	1 (13.9%)	0	0
Asthenozoospermia	27 (27.6%)	0	0 (25.3%)	(42.9%)	(100.0%)
Oligoasthenozoospermia	1 (1.0%)	(18.2%)	0	0	0
Teratozoospermia	19 (19.4%)	(18.2%)	3 (16.5%)	0	0
Oligoasthenoteratozoospermia	38 (38.8%)	(27.3%)	9 (24.1%)	(57.1%)	0

**P<0.001**

From the Relationship between WHO Sperm Analysis with Fertilization Figures, there is better fertilization (Z1) found in the WHO sperm analysis group which is normozoospermia when compared with each of the other groups, followed by oligoasthenoteratozoospermia, then subsequently oligoasthenozoospermia and teratozoospermia with the respective numbers each the same. As for Z2, there are more in the asthenozoospermia group followed by oligoasthenoteratozoospermia, normozoospermia, teratozoospermia, and oligozoospermia. While the results of the most negative fertilization in the oligoasthenoteratozoospermia group. Results from the analysis of the above table between World Health Organization sperm analysis with fertilization rates obtained statistically significant results  $p < 0.05$ .

**Table 5. Analysis results of examining WHO sperm parameters by embryogenesis scoring**

Variable	Scoring Embryogenesis				
	Negative	A	B	C	D
Analysis Sperma WHO					
Normozoospermia	1 (10.0%)	(62.5%)	(18.6%)	(11.1%)	0
Oligozoospermia	(4.5%)	0	4 (9.3%)	7 (19.4%)	0
Asthenozoospermia	8 (25.5%)	3 (37.5%)	1 (25.6%)	11 (30.6%)	1 (50.0%)
Oligoasthenozoospermia	1 (0.9%)	0	0	2 (5.6%)	0
Teratozoospermia	3 (20.9%)	0	8 (18.6%)	3 (8.3%)	0
Oligoasthenoteratozoospermia	2 (38.2%)	0	12 (27.9%)	9 (25.0%)	1 (50.0%)

**P=0.006**

From the Relationship between WHO Sperm Analysis with Fertilization Figures, there is better fertilization (Z1) found in the WHO sperm analysis group which is normozoospermia when compared with each other group, followed by oligoasthenoteratozoospermia, then subsequently oligoasthenozoospermia and teratozoospermia with numbers from the other groups. WHO Sperm Analysis with Embryogenesis Scoring shows better embryo scoring (Grade A) in WHO normozoospermia sperm analysis when compared with each other group and also seen worse embryo scoring (Grades C and D) in the abnormal group namely asthenozoospermia and oligoasthenoteratozoospermia. Most negative embryogenesis scoring in the oligoasthenoteratozoospermia group was followed by asthenozoospermia and teratozoospermia. Results from the analysis of the above table Relationship between WHO Sperm Analysis with Embryogenesis Scoring obtained statistically significant results  $p < 0.05$ .

In this study, we tried to exclude factors that could affect the quality of oocytes such as PCOS and Endometriosis, so that the study subjects included in this study only consisted of cases of infertility caused by male factors, myoma uteri, both tubal non-patents, hydrosalpinges, and unexplained infertility. We did this to get more reliable results, although we therefore only got 29 infertile couples and 200 oocytes during the study period.

From the analysis of sperm DNA fragmentation index by examining WHO sperm parameters, it appears that the sperm DNA fragmentation index group  $\leq 30\%$  is dominated by an abnormal group but in the sperm DNA fragmentation index group  $> 30\%$  we did not get normozoospermia whereas in the group sperm DNA fragmentation index  $\leq 30\%$  we get 23.1%. This is consistent with the study of Lin et al in his study with 86 cases of ICSI found there was an influence between sperm DNA fragmentation on sperm motility and also Irvine et al who stated that there was a very significant negative relationship between sperm concentration and sperm DNA damage [29;30].

From the analysis of sperm DNA fragmentation index with fertilization rates, it is seen that better fertilization rates are more numerous in the sperm DNA fragmentation index group  $\leq 30\%$ , according to the research of Muriel et al and Benchaib et al who stated a negative correlation between the level of sperm DNA

fragmentation with fertilization rates. [31;5;32]. However, the negative fertilization group was also more prevalent in the sperm DNA fragmentation index group  $\leq 30\%$ .

From the analysis of sperm DNA fragmentation index by the embryo, scoring can be obtained that better embryogenesis occurs in the sperm DNA fragmentation index group  $\leq 30\%$ , by the results obtained by Bungum et al in his 2004 study in which biochemical pregnancy rates in the fragmentation group Sperm DNA  $< 27\%$  (51.2%) is higher than the sperm DNA fragmentation group  $> 27\%$  (47.1%) and also Virro et al who stated that men with sperm DNA fragmentation  $> 30\%$  had a higher risk for low blastocyst numbers ( $< 30\%$ ) and also the cessation of pregnancy [33;34;35]. However, we also found negative embryogenesis which was also higher in the sperm DNA fragmentation index group  $\leq 30\%$ .

From the results of an analysis of WHO sperm parameters examination with fertilization rates, it appears that the normozoospermia group produces the highest quality fertilization (Z1), but in the Z2 group it is dominated by abnormal groups, namely asthenozoospermia and oligoasthenoteratozoospermia. The most negative fertilization comes from the oligoasthenoteratozoospermia group. From the results of an analysis of WHO sperm parameters examination by embryogenesis scoring obtained the most Grade A embryogenesis score scoring in the normozoospermia WHO sperm analysis.

These variations in output from sperm DNA fragmentation may be explained as follows; First, sperm DNA fragmentation may not show similar damage in terms of quantity or quality. Second, the outcome may be a balance between sperm DNA damage and the ability to repair from oocyte [36;37;38;39]. Li Z et al in a systematic review and meta-analysis found that sperm DNA damage did not have a significant impact on the chance of getting a clinical pregnancy in patients undergoing IVF or ICSI [40;41;42;43].

There are conflicting results regarding the effect of sperm DNA fragmentation on semen parameters and also regarding the need for examination of sperm chromatin assays as routine laboratory examinations. Some studies report no relationship between sperm DNA fragmentation and semen parameters, while others find a negative relationship on some or all parameters.

Our data show significant results in the relationship between obtaining fertilization but not statistically significant in the relationship between sperm DNA fragmentation with embryogenesis even though there are significant differences between sperm DNA fragmentation groups  $\leq 30\%$  and  $> 30\%$ . This might be due to the small number of research subjects in our study, but this is also made possible by the complexity of the genomic integrity of spermatozoa as a determining factor in TRB.

It is also well known that oocytes can repair fragmented sperm DNA, which is also likely to have a major influence on the outcome of the research. Besides sperm, DNA integrity cannot be considered an absolute picture of all the paternal effects that control early embryonic activity and development after ICSI/IVF [44].

#### 4. CONCLUSION

In this study, it was seen that the sperm DNA fragmentation rate could show a negative relationship with the rate of fertilization but could not show a statistically significant relationship with the rate of embryogenesis. A statistically significant relationship was seen between WHO sperm analysis parameters and DNA fragmentation index examined using the SCD method. Based on this study it can be suggested that examination of sperm DNA fragmentation be included as an integrated part of routine sperm analysis examinations in infertile patients.

#### REFERENCES

- [1] Jungwirth, A., Diemer, T., Dohle, G. R., Giwercman, A., Kopa, Z., Tournaye, H., & Krausz, C. (2004). Guidelines for the investigation and treatment of male infertility. *Eur Urol*, 46(5), 555-8.
- [2] Fritz, M. A. (2011). Assisted reproductive technologies. U: Fritz MA, Speroff L.: Clinical gynecologic endocrinology and infertility.
- [3] Larson, K. L., DeJonge, C. J., Barnes, A. M., Jost, L. K., & Evenson, D. P. (2000). Sperm chromatin structure assay parameters as predictors of failed pregnancy following assisted reproductive techniques. *Human Reproduction*, 15(8), 1717-1722.
- [4] Evenson, D. P., Jost, L. K., Marshall, D., Zinman, M. J., Clegg, E., Purvis, K., ... & Claussen, O. P. (1999). The utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Human Reproduction*, 14(4), 1039-1049.
- [5] Shaman, J. A., Yamauchi, Y., & Ward, W. S. (2007). Sperm DNA fragmentation: awakening the sleeping genome.
- [6] Benchaib, M., Braun, V., Lornage, J., Hadj, S., Salle, B., Lejeune, H., & Guérin, J. F. (2003). Sperm DNA fragmentation decreases the pregnancy rate in an assisted reproductive technique. *Human Reproduction*, 18(5), 1023-1028.
- [7] Fernández, J. L., Muriel, L., Rivero, M. T., Goyanes, V., Vazquez, R., & Alvarez, J. G. (2003). The sperm chromatin dispersion test: a simple method for the determination of sperm DNA fragmentation. *Journal of andrology*, 24(1), 59-66.
- [8] Tamburrino, L., Marchiani, S., Montoya, M., Marino, F. E., Natali, I., Cambi, M., ... & Muratori, M. (2012). Mechanisms and clinical correlates of sperm DNA damage. *Asian journal of andrology*, 14(1), 24.
- [9] Gil-Villa, A. M., Cardona-Maya, W., Agarwal, A., Sharma, R., & Cadavid, Á. (2009). Role of the male factor in early recurrent embryo loss: do antioxidants have any effect?. *Fertility and sterility*, 92(2), 565-571.
- [10] Simon, L., Lutton, D., McManus, J., & Lewis, S. E. (2011). Sperm DNA damage measured by the alkaline Comet assay as an independent predictor of male infertility and in vitro fertilization success. *Fertility and sterility*, 95(2), 652-657.
- [11] Bungum, M. (2012). Sperm DNA integrity assessment: a new tool in the diagnosis and treatment of fertility. *Obstetrics and gynaecology international*, 2012.
- [12] Cohen-Bacrie, P., Belloc, S., Ménézo, Y. J., Clement, P., Hamidi, J., & Benkhalifa, M. (2009). Correlation between DNA damage and sperm

- parameters: a prospective study of 1,633 patients. *Fertility and sterility*, 91(5), 1801-1805.
- [13] De Iuliis, G. N., Thomson, L. K., Mitchell, L. A., Finnie, J. M., Koppers, A. J., Hedges, A., ... & Aitken, R. J. (2009). DNA damage in human spermatozoa is highly correlated with the efficiency of chromatin remodelling and the formation of 8-hydroxy-2'-deoxyguanosine, a marker of oxidative stress. *Biology of reproduction*, 81(3), 517-524.
  - [14] Sakkas, D., & Alvarez, J. G. (2010). Sperm DNA fragmentation: mechanisms of origin, impact on reproductive outcome, and analysis. *Fertility and sterility*, 93(4), 1027-1036.
  - [15] Brazil, C. (2010). Practical semen analysis: from A to Z. *Asian journal of andrology*, 12(1), 14.
  - [16] Khalili, M. A., Aghaie-Maybodi, F., Anvari, M., & Talebi, A. R. (2009). Sperm nuclear DNA in ejaculates of fertile and infertile men: correlation with semen parameters. *Urology Journal*, 3(3), 154-159.
  - [17] Moustafa, M. H., Sharma, R. K., Thornton, J., Mascha, E., Abdel-Hafez, M. A., Thomas, A. J., & Agarwal, A. (2004). Relationship between ROS production, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. *Human Reproduction*, 19(1), 129-138.
  - [18] Berger, D. S., AbdelHafez, F., Russell, H., Goldfarb, J., & Desai, N. (2011). Severe teratozoospermia and its influence on pronuclear morphology, embryonic cleavage and compaction. *Reproductive Biology and Endocrinology*, 9(1), 37.
  - [19] Esteves, S. C., & Agarwal, A. (2011). Novel concepts in male infertility. *International Braz j Urol*, 37(1), 5-15.
  - [20] Tesarik, J., Greco, E., & Mendoza, C. (2004). Late, but not early, the paternal effect on human embryo development is related to sperm DNA fragmentation. *Human Reproduction*, 19(3), 611-615.
  - [21] Knez, K., Zorn, B., Tomazevic, T., Vrtacnik-Bokal, E., & Virant-Klun, I. (2011). The IMSI procedure improves poor embryo development in the same infertile couples with poor semen quality: a comparative prospective randomized study. *Reproductive Biology and Endocrinology*, 9(1), 123.
  - [22] Cohen-Bacrie, P., Belloc, S., Ménézo, Y. J., Clement, P., Hamidi, J., & Benkhalifa, M. (2009). Correlation between DNA damage and sperm parameters: a prospective study of 1,633 patients. *Fertility and sterility*, 91(5), 1801-1805.
  - [23] Hourcade, J. D., Pérez-Crespo, M., Fernández-González, R., Pintado, B., & Gutiérrez-Adán, A. (2010). Selection against spermatozoa with fragmented DNA after postovulatory mating depends on the type of damage. *Reproductive Biology and Endocrinology*, 8(1), 1-11.
  - [24] Nadeak, B., Deliviana, E., Sormin, E., Naibaho, L., & Juwita, C. P. (2019). Coaching Marriage Endurance and Family Harmony With a Theme "The Family Relationship and Intimacy. *Journal ComunitÁ Servizio*, 1(2), 179-185
  - [25] Agarwal, A., & Allamaneni, S. S. (2004). The effect of sperm DNA damage on assisted reproduction outcomes. *Minerva Ginecol*, 56(3), 235-245.
  - [26] Nadeak, B., Deliviana, E., Sormin, E., Naibaho, L., & Juwita, C. P. (2019). Coaching Marriage Endurance and Family Harmony With a Theme "The Family Relationship and Intimacy. *Journal ComunitÁ Servizio*, 1(2), 179-185
  - [27] Scott, L., Alvero, R., Leondires, M., & Miller, B. (2000). The morphology of human pronuclear embryos is positively related to blastocyst development

- and implantation. *Human Reproduction*, 15(11), 2394-2403.
- [28] Lin, M. H., Lee, R. K. K., Li, S. H., Lu, C. H., Sun, F. J., & Hwu, Y. M. (2008). Sperm chromatin structure assay parameters are not related to fertilization rates, embryo quality, and pregnancy rates in vitro fertilization and intracytoplasmic sperm injection, but might be related to spontaneous abortion rates. *Fertility and sterility*, 90(2), 352-359.
  - [29] Nadeak, B., Iriani, U. E., Naibaho, L., Sormin, E., & Juwita, C. P. (2019). Building Employees' Mental Health: The Correlation between Transactional Leadership and Training Program with Employees' Work Motivation at XWJ Factory. *Indian Journal of Public Health Research & Development*, 10(6), 1373-1379.
  - [30] Muriel, L., Garrido, N., Fernández, J. L., Remohí, J., Pellicer, A., de los Santos, M. J., & Meseguer, M. (2006). Value of the sperm deoxyribonucleic acid fragmentation level, as measured by the sperm chromatin dispersion test, in the outcome of in vitro fertilization and intracytoplasmic sperm injection. *Fertility and Sterility*, 85(2), 371-383.
  - [31] Nadeak, B., Naibaho, L., Sormin, E., & Juwita, C. P. (2019). Healthy Work Culture Stimulate Performance. *Indian Journal of Public Health Research & Development*, 10(6), 1385-1389.
  - [32] Bungum, M., Humaidan, P., Spano, M., Jepson, K., Bungum, L., & Giwercman, A. (2004). The predictive value of sperm chromatin structure assay (SCSA) parameters for the outcome of intrauterine insemination, IVF and ICSI. *Human Reproduction*, 19(6), 1401-1408.
  - [33] Nadeak, B., Simanjuntak, D. R., Naibaho, L., Sormin, E., Juwita, C. P., & Pardede, S. O. (2019). Analysis of Nursing Quality Services. *Indian Journal of Public Health Research & Development*, 10(6), 1380-1384.
  - [34] Virro, M. R., Larson-Cook, K. L., & Evenson, D. P. (2004). Sperm chromatin structure assay (SCSA®) parameters are related to fertilization, blastocyst development, and ongoing pregnancy in vitro fertilization and intracytoplasmic sperm injection cycles. *Fertility and sterility*, 81(5), 1289-1295.
  - [35] Karydis, S., Asimakopoulos, B., Papadopoulos, N., Vakalopoulos, I., Al-Hasani, S. A. F. A. A., & Nikoletos, N. (2005). ICSI outcome is not associated with the incidence of spermatozoa with abnormal chromatin condensation. *in vivo*, 19(5), 921-925.
  - [36] Nadeak, B., & Naibaho, L. (2019, November). Investigating the effect of learning multimedia and thinking style preference on learning achievement on anatomy at Universitas Kristen Indonesia. In *Journal of Physics: Conference Series* (Vol. 1387, No. 1, p. 012116). IOP Publishing.
  - [37] Tamburrino, L., Marchiani, S., Montoya, M., Marino, F. E., Natali, I., Cambi, M., ... & Muratori, M. (2012). Mechanisms and clinical correlates of sperm DNA damage. *Asian journal of andrology*, 14(1), 24.
  - [38] Lewis, S. E., Aitken, R. J., Conner, S. J., De Iuliis, G., Evenson, D. P., Henkel, R., ... & Gharagozloo, P. (2013). The impact of sperm DNA damage in assisted conception and beyond: recent advances in diagnosis and treatment. *Reproductive biomedicine online*, 27(4), 325-337.
  - [39] Li, Z., Wang, L., Cai, J., & Huang, H. (2006). Correlation of sperm DNA damage with IVF and ICSI outcomes: a systematic review and meta-analysis. *Journal of assisted reproduction and genetics*, 23(9-10), 367-376.
  - [40] Bach, P. V., & Schlegel, P. N. (2016). Sperm DNA damage and its role in

- IVF and ICSI. *Basic and clinical andrology*, 26(1), 1-10.
- [41] Wright, C., Milne, S., & Leeson, H. (2014). Sperm DNA damage caused by oxidative stress: modifiable clinical, lifestyle and nutritional factors in male infertility. *Reproductive biomedicine online*, 28(6), 684-703.
- [42] Ni, K., Spiess, A. N., Schuppe, H. C., & Steger, K. (2016). The impact of sperm protamine deficiency and sperm DNA damage on human male fertility: a systematic review and meta-analysis. *Andrology*, 4(5), 789-799.
- [43] Gavrilouk, D., & Aitken, R. J. (2015). Damage to sperm DNA mediated by reactive oxygen species: its impact on human reproduction and the health trajectory of offspring. *The male role in pregnancy loss and embryo implantation failure* (pp. 23-47). Springer, Cham.
- [44] Sheikh, N., Amiri, I., Farimani, M., Najafi, R., & Hadeie, J. (2008). Correlation between sperm parameters and sperm DNA fragmentation in fertile and infertile men. *International Journal of Reproductive BioMedicine*, 6(2), 13-0.

# The Effect of Sperm DNA Fragmentation on Fertilization and Embryogenesis Rates

## ORIGINALITY REPORT

26%

SIMILARITY INDEX

14%

INTERNET SOURCES

24%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1

G. Coticchio. "Sperm DNA fragmentation: paternal effect on early post-implantation embryo development in ART", Human Reproduction, 09/01/2006

Publication

2%

2

"A Clinician's Guide to Sperm DNA and Chromatin Damage", Springer Science and Business Media LLC, 2018

Publication

2%

3

[downloads.hindawi.com](https://downloads.hindawi.com)

Internet Source

1%

4

Fernandez, J.L.. "Simple determination of human sperm DNA fragmentation with an improved sperm chromatin dispersion test", Fertility and Sterility, 200510

Publication

1%

5

"In Vitro Fertilization", Springer Science and Business Media LLC, 2019

Publication

1%

6	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1 %
7	"Abstracts of the 35th Annual Meeting of the European Society of Human Reproduction and Embryology", Human Reproduction, 2019 Publication	1 %
8	<a href="http://www.andrologyjournal.org">www.andrologyjournal.org</a> Internet Source	1 %
9	Khadem, N., A. Poorhoseyni, M. Jalali, A. Akbary, and S. T. Heydari. "Sperm DNA fragmentation in couples with unexplained recurrent spontaneous abortions", Andrologia, 2012. Publication	1 %
10	<a href="http://journals.sagepub.com">journals.sagepub.com</a> Internet Source	1 %
11	Submitted to National postgraduate Medical College of Nigeria Student Paper	1 %
12	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	1 %
13	K.-C. Lan. "The predictive value of using a combined Z-score and day 3 embryo morphology score in the assessment of embryo survival on day 5", Human Reproduction,	1 %

06/01/2003

Publication

14

A. Zini. "Potential adverse effect of sperm DNA damage on embryo quality after ICSI", Human Reproduction, 07/29/2005

Publication

1 %

15

"Abstracts of the 34rd Annual Meeting of the European Society of Human Reproduction and Embryology", Human Reproduction, 2018

Publication

1 %

16

Sperm Chromatin, 2011.

Publication

<1 %

17

"Endocrinology of the Testis and Male Reproduction", Springer Science and Business Media LLC, 2017

Publication

<1 %

18

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

19

[www.idosi.org](http://www.idosi.org)

Internet Source

<1 %

20

[www.ijphrd.com](http://www.ijphrd.com)

Internet Source

<1 %

21

Muriel, L.. "Value of the sperm deoxyribonucleic acid fragmentation level, as measured by the sperm chromatin dispersion test, in the outcome of in vitro fertilization and intracytoplasmic

<1 %

22

"Male Infertility", Springer Science and Business Media LLC, 2020

Publication

<1 %

23

H KIM. "InfertilityThe Male", Principles of Gender-Specific Medicine, 2010

Publication

<1 %

24

Submitted to University of Technology, Sydney

Student Paper

<1 %

25

Isabel Pons, Rosa Cercas, Celia Villas, Cristina Braña, Sylvia Fernández-Shaw. "One abstinence day decreases sperm DNA fragmentation in 90 % of selected patients", Journal of Assisted Reproduction and Genetics, 2013

Publication

<1 %

26

[www.rbej.com](http://www.rbej.com)

Internet Source

<1 %

27

[www.gynemed.de](http://www.gynemed.de)

Internet Source

<1 %

28

María Hebles, Monica Dorado, Miguel Gallardo, Mercedes González-Martínez, Pacual Sánchez-Martín. "Seminal quality in the first fraction of ejaculate", Systems Biology in Reproductive Medicine, 2014

<1 %

29

[www.researchgate.net](http://www.researchgate.net)

Internet Source

<1 %

30

Hewedy, El-Sayed S., Desouky E. Abou Ammo, Tarek E. Amin, Naeim M. Dawoud, Arwa M. Hassan, Mohamed Attaia, Ola El-Shora, and Mohamed M. Fawzy. "Sperm DNA damage in varicocele patients :", Human Andrology, 2012.

Publication

<1 %

31

[tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr)

Internet Source

<1 %

32

Abdelshakour A. Al-Mohammady, Ahmed F. El-Sherbiny, Amira B. Mehaney, Yasser A. Ghobara. "Varicocele repair in patients prepared for intracytoplasmic sperm injection: To do or not to do?", Andrologia, 2019

Publication

<1 %

33

Hassan W. Bakos. "Sperm DNA damage is associated with assisted reproductive technology pregnancy", International Journal of Andrology, 9/5/2007

Publication

<1 %

34

Chun-Chia Huang, David Pei-Cheng Lin, Hui-Mei Tsao, Tzu-Chun Cheng, Chung-Hsien Liu, Maw-Sheng Lee. "Sperm DNA fragmentation negatively correlates with velocity and

<1 %

fertilization rates but might not affect pregnancy rates", Fertility and Sterility, 2005

Publication

35

M. B. Herrero, M. F. Lusignan, W.-Y. Son, M. Sabbah, W. Buckett, P. Chan. " outcomes using testicular spermatozoa in non-azoospermic couples with recurrent failure and no previous live births ", Andrology, 2019

Publication

<1 %

36

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

37

[molehr.oxfordjournals.org](http://molehr.oxfordjournals.org)

Internet Source

<1 %

38

K. R. Chohan. "Comparison of Chromatin Assays for DNA Fragmentation Evaluation in Human Sperm", Journal of Andrology, 2006

Publication

<1 %

39

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

<1 %

40

[www.bioone.org](http://www.bioone.org)

Internet Source

<1 %

41

[file.scirp.org](http://file.scirp.org)

Internet Source

<1 %

42

M. Sergerie. "Sperm DNA fragmentation: threshold value in male fertility", Human

<1 %

43

[irhrc.sbmu.ac.ir](http://irhrc.sbmu.ac.ir)

Internet Source

<1 %

44

Dul, E. C., C. M. A. van Ravenswaaij-Arts, H. Groen, J. van Echten-Arends, J. A. Land, Y. Tyulenev, V. Naumenko, L. Kurilo, L. Shileiko, A. Segal, R. Klimova, A. Kushch, J. Ribas-Maynou, A. Garcia-Peiro, C. Abad, M. J. Amengual, J. Benet, J. Navarro, A. Co. "POSTER VIEWING SESSION - ANDROLOGY", Human Reproduction, 2011.

Publication

<1 %

45

[hdl.handle.net](http://hdl.handle.net)

Internet Source

<1 %

46

[buleria.unileon.es](http://buleria.unileon.es)

Internet Source

<1 %

47

Lin-Tao Xue, Rui-Xue Wang, Bing He, Wei-Ying Mo et al. "Effect of sperm DNA fragmentation on clinical outcomes for Chinese couples undergoing in vitro fertilization or intracytoplasmic sperm injection", Journal of International Medical Research, 2016

Publication

<1 %

48

Sonia Brahem, Hatem Elghezal, Houda Ghédir, Hanène Landolsi et al. "Cytogenetic and

<1 %

# Molecular Aspects of Absolute Teratozoospermia: Comparison Between Polymorphic and Monomorphic Forms", Urology, 2011

Publication

---

---

Exclude quotes      On

Exclude bibliography      On

Exclude matches      Off