



Source details

Journal of Physics: Conference Series

Scopus coverage years: from 2005 to Present

Publisher: Institute of Physics Publishing

ISSN: 1742-6588 E-ISSN: 1742-6596

Subject area: Physics and Astronomy: General Physics and Astronomy

Source type: Conference Proceeding

CiteScore 2020

0.7



SJR 2020

0.210



SNIP 2020

0.464



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

$$0.7 = \frac{52,411 \text{ Citations } 2017 - 2020}{72,842 \text{ Documents } 2017 - 2020}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021

$$0.6 = \frac{52,959 \text{ Citations to date}}{81,810 \text{ Documents to date}}$$

Last updated on 04 August, 2021 • Updated monthly

CiteScore rank 2020

Category	Rank	Percentile
Physics and Astronomy		
General Physics and Astronomy	#191/233	18th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

- [What is Scopus](#)
- [Content coverage](#)
- [Scopus blog](#)
- [Scopus API](#)
- [Privacy matters](#)

Language

- [日本語に切り替える](#)
- [切换到简体中文](#)
- [切换到繁體中文](#)
- [Русский язык](#)

Customer Service

- [Help](#)
- [Contact us](#)



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name



Home

Journal Rankings

Country Rankings

Viz Tools


Help

About Us

Journal of Physics: Conference Series

COUNTRY

[United Kingdom](#)



SUBJECT AREA AND CATEGORY

[Physics and Astronomy](#)
└ [Physics and Astronomy \(miscellaneous\)](#)

PUBLISHER

[IOP Publishing Ltd.](#)

Ad closed by Google

PUBLICATION TYPE

Journals

ISSN

17426588, 17426596

COVERAGE

2005-2020




Ad closed by Google

SCOPE


Join the conversation about this journal






Ad closed by Google

Quartiles



Citations per document





Journal of Physics: Conference Series

Q4 Physics and Astronomy (miscellaneous) best quartile

SJR 2020 0.21

powered by scimagojr.com

Show this widget in your own website


Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="629 379 779 392">
```

SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it



Scopus Indexed Journal

Call for Papers August Issue

Fast Track Peer Reviewed Publication.

tojqi.net

OPEN

Metrics based on Scopus® data as of April 2021



Arcadius Benawa 2 weeks ago

How about publication in ScitePress? Is it indexed by Scopus? My paper which I presented in Ebic 2019 has been published in ScitePress publication in 2021 and so my paper which I have

Table of contents

Volume 1783

2021

◀ Previous issue Next issue ▶

Annual Conference on Science and Technology Research (ACOSTER) 2020, 20-21 June 2020, Medan, Indonesia

Accepted papers received: 20 January 2021

Published online: 12 February 2021

Open all abstracts

Preface

OPEN ACCESS 011001

Preface Annual Conference on Science and Technology Research (ACOSTER) 2020

+ Open abstract  View article  PDF

OPEN ACCESS 011002

Peer review declaration

+ Open abstract  View article  PDF

Papers

OPEN ACCESS 012001

Effect of Different Solvents On The Phytochemical Compounds of *Sargassum* sp. From Yogyakarta and East Nusa Tenggara

Dheasy Herawati and Pratiwi Pudjiastuti

+ Open abstract  View article  PDF

OPEN ACCESS 012002


Radical catcher activities free of methanol fruit leather and pomegranate seeds (*punica granatum. L*)

Vina N Van Harling

+ Open abstract  View article  PDF

OPEN ACCESS 012003

Concentration Effect of Ethanol Extract *Pinus merkusii* Leaves Litter on *Zea mays* L.

Seed Germination. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy. 

DNA Amplification of Methanogenic Bacteria from The Sewage of Coconut Water by Specific Primer 16S rRNA and Specific Methanogenic Gene

Ardi Kapahang, Maria Bintang and Dedy D Solihin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012107

Analysis of Intensity, Duration, and Frequency Rain Daily of Java Island Using Mononobe Method

Faradiba

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012108

Problem Based Learning using Open Educational Resources to enhance Higher Order Thinking Skills in Physics Learning

Dwi Sulisworo, Fakhrunisyah and Khusnul Basriyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012109

The role of information technology on the online learning success based on self-regulated learning data

Dwi Sulisworo, Ika Maryani and Dian Artha Kusumaningtyas

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012110

Effectiveness of E-Learning Approach to Contextual Teaching and Learning in Improving Students 'Ability

Zulfi Azhar, Suparno, Kasman Rukun, Jalius Jama, Hansi Effendi and Mukhlidi Muskhir

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012111

Gamification in Learning using Quizizz Application as Assessment Tools

Wiwini Handoko, Eva Mizkat, Auliana Nasution, Hambali and Juna Eska

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012112

Face to Face Learning vs Blended Learning vs Online Learning (Student Perception of Learning)

Awal Kurnia Putra Nasution, Andika Hariyanto Surbakti, Rahmanita Zakaria, Sungkawati Kardi Wahyuningsih and Leni Agustina Daulay

[+ Open abstract](#) [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy](#) and [Cookies](#) policy.

012113

PAPER • OPEN ACCESS

Analysis of Intensity, Duration, and Frequency Rain Daily of Java Island Using Mononobe Method

To cite this article: Faradiba 2021 *J. Phys.: Conf. Ser.* **1783** 012107

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Analysis of Intensity, Duration, and Frequency Rain Daily of Java Island Using Mononobe Method

Faradiba

Department of Physics Education, Universitas Kristen Indonesia

*faradiba@uki.ac.id

Abstract. The importance of analyzing rainfall is expected to help in handling disaster-prone areas, so that the consequences can be minimized. This study aims to determine the intensity of rain in the short term. The data used in this study is the annual rainfall data in Java for 2005 - 2019 for each province. The method used in this study is the Mononobe Method to see the intensity of daily rain and its daily return period. The results of this study indicate that the intensity of rain during the duration of one hour to three hours decreases quite significantly, but at the duration of four to eight hours the decrease in rain intensity is not as big as the decrease in the duration of one hour to three hours. The highest rainfall intensity is in the one hour duration in West Java Province. However, the intensity for the five to eight hour duration tends to be the same for 6 provinces in Java. The rainfall threshold in Java is included in the category of heavy rain. High rainfall intensity lasts for a short duration, on the other hand, rainfall with low intensity tends to occur over a relatively long duration.

Keywords: Daily rain intensity, Rainfall, Java Island, Mononobe Method

1. Introduction

The climate that has occurred in Indonesia in recent years has experienced a shift, both for the rainy season and the dry season. Changes in the rainy season which are quite extreme can result in drought or floods [1], [2]. The climate phenomenon that occurs is quite difficult to predict precisely, so it can have a negative impact on various sectors, particularly the agricultural sector. The worst impact that appears in the agricultural sector is crop failure [3], [4].

Indonesia has about 270 million inhabitants and has been named an agricultural country. This condition is reflected in the livelihoods of most of the population, who mostly work as farmers. Agricultural production in Indonesia is dominated by Java Island, so that Java Island has an important role in meeting national food needs. Geographically, Java Island is bordered by the Java Sea on the northern side. The south is bordered by the Indian Ocean. To the east by the Bali Strait, and to the west by the Sunda Strait [5]. Based on information submitted by the Meteorology, Climatology and Geophysics Agency (BMKG), regarding early warning of extreme weather, where heavy rain and strong winds will occur on April 15-16, 2020. This is due to low pressure in the Indian Ocean (southwest Sumatra), which gives the effect of increasing wind speed in low layers, up to more than 25 knots [6]. Meanwhile, Java Island, which is located in the south, will feel relatively hot temperatures, this is due to the position of the sun in the southern hemisphere of Indonesia.

Almost all parts of Indonesia have a tropical climate, as does Java Island. The annual average temperature in Java can reach 22-29 degrees Celsius, while the average humidity is around 75 percent



[7]. The temperature in Java varies widely, where during the day and during the dry season the temperature around the coast is higher, reaching 34 degrees Celsius.

The position of the coastal plains also greatly affects temperature variations [8], [9], where areas on the north coast have temperatures that tend to be higher (hotter) than areas on the south coast [10], [11]. Meanwhile, in mountainous areas, the temperature is lower or cooler than lowland areas or coastal areas [12], [13].

Java Island is located in the equatorial rain belt. This results in a typical tropical climate characterized by high levels of rainfall, temperature and humidity [14]. Java Island experiences 2 types of seasons, namely the rainy season which normally occurs from December to March (the following year), and the dry season from May to October.

The temperature in the rainy season ranges from 21 °C (70 °F) to 33 °C (90 °F), except in high elevation areas (mountains), where the temperature is much cooler. The lowest temperature ever recorded on the island is 18 °C (65 °F). Frequent seasonal changes in wind direction (monsoons) from the Indian Ocean and Pacific Ocean, resulting in more varied climatic conditions on the island [15], [16].

Temperature in an area tends to correlate with rainfall intensity [17]. In the dry season with temperatures higher than average, tend to have relatively low rainfall intensity, and vice versa. So that through the measurement of rainfall intensity, it can be seen that the start of the dry or rainy season [18], [19].

Rain intensity is the amount of rain expressed in the height of rain or the volume of rain per unit time. The amount of intensity varies, depending on the length of rainfall and the frequency of its occurrence [20]. Rain intensity is obtained by analyzing rainfall data, both statistically and empirically.

Rain intensity is associated with short-term rain duration, for example 5 minutes, 30 minutes, 60 minutes and hours. This short-term rainfall data can only be obtained by using an automatic rain recorder. In Indonesia, there are not many of these tools, so in their measurements more common rain recording devices are used, which measure 24-hour rain or called daily rain [21].

In diversifying rain into streams, there are several characteristics of rain that are important to note, including rain intensity (I), length of time of rain (t), rain depth (d), frequency (f), and area of influence of rain (A). Analysis of the relationship between two important rain parameters in the form of intensity and duration can be statistically related to a frequency of occurrence. The graphical presentation of this relationship is in the form of an Intensity-Duration-Frequency (IDF) curve. [22]. This curve can be used for the calculation of runoff and peak discharge at concentration time. Through this curve, it will also be known the probability of rainfall intensity occurring for the duration of indiscriminate rainfall.

Several studies on rainfall intensity analysis and IDF Curve have been done before [23], [24]. IDF curve depiction based on the results of rain intensity analysis using the Van Breen and Hasper der Weduwen method with the Talbot, Sherman and Ishiguro equation approach has been carried out. The IDF curve of daily rainfall is depicted based on rainfall intensity using the Mononobe formula. IDF curves were also created for the Mount Merapi slopes using several intensity formulas, such as the Sherman, Kimijima, Haspers and Mononobe formulas. [25]. Apart from that, another study also analyzed the IDF curve for the Surakarta City [26]. IDF curves are created using several approach formulas, such as Talbot, Sherman, Ishiguro and SDR-IDF.

The question is, what if we only have the accumulated daily (monthly) rain data? Of course this is not an obstacle for us to calculate the intensity of rain for a short time duration (minutes or hours), because the rain intensity for a short duration can be estimated using the Mononobe formula. The choice of this formula is due to the available daily rainfall data. Generally the IDF curve is formed using short duration rain data. However, if short period rainfall data is not available, then the IDF curve can still be created using daily rainfall data using the approach formula, namely the Mononobe and Haspers Formulas. So that the purpose of this study is to determine and analyze the intensity of rainfall for various periods in Java.

2. Data and Methodology

In this study, the object of research was the island of Java, which consists of six provinces: DKI Jakarta, West Java, Central Java, DI Yogyakarta, East Java, and Banten. The variables used in this study are rainfall data from 2005 - 2019 which were taken from the Indonesian Statistical Publication published by the Central Statistics Agency.

The data collection and processing stages include (i) data collection by year for each province; (ii) Data entry and tabulation; (iii) Calculating the required statistical parameters. The required statistical parameters include: average, standard deviation, coefficient of variation, inclination coefficient, kurtosis coefficient and type of data distribution; (iv) Determination of the coefficient of Reduce Mean (Y_n), Reduce variation (Y_t) and Reduce standard deviation (S_n); (v) Calculation of daily rainfall (24 hours) for each province [27].

The calculation of rainfall for a 24 hour period uses the Gumbel formula [28].

$$R_{24} = \bar{X} + \frac{S_x}{S_n}(Y_t - Y_n) \quad \dots\dots\dots [1]$$

where:

- R_{24} = The amount of daily rainfall maximum 24 hours (mm / 24 hours)
- \bar{X} = Average rainfall (mm)
- S_x = Standard deviation
- Y_n = Reduced mean
- S_n = Reduced standard deviation
- Y_t = Reduced variation as the return period

Data analysis using the Mononobe method [29].

$$I = \frac{R_{24}}{24} \left(\frac{24}{t} \right)^{\frac{2}{3}} \quad \dots\dots\dots [2]$$

where:

- I = Rainfall intensity in mm / hour.
- R_{24} = Maximum rainfall in 24 hours in mm
- T = The amount of rain is in hours

3. Result and Discussion

Rainfall data from 2005 - 2019 were obtained from certain measurement stations, where each province was selected so that it could represent the condition of the province as a whole. On average, West Java Province has high rainfall compared to other provinces in Java Island. This is supported by land elevation conditions. Meanwhile, Banten Province is the province with the lowest rainfall, this is because many of these provinces are located in coastal areas.

Table 1. Java Island Annual Rainfall, 2005 - 2019

Year	Jakarta	West Java	Central Java	SR Yogyakarta	East Java	Banten
Station	Kemayoran	Bandung	Semarang	Yogyakarta	Juanda	Serang
2005	1,433.50	1,796.30	1,132.10	1,886.00	2,260.80	1,192.00
2006	1,527.10	1,368.00	1,142.00	1,824.50	1,446.20	1,563.00
2007	2,353.90	2,219.00	1,868.00	1,919.00	1,515.20	1,735.00
2008	1,779.10	2,001.00	2,690.30	1,870.80	1,502.90	1,476.00
2009	1,973.00	2,098.00	2,807.00	1,584.00	1,987.00	1,414.00
2010	2,405.00	3,868.40	3,228.00	1,390.00	2,895.00	2,136.00
2011	1,274.00	1,789.00	1,879.00	2,285.00	1,790.00	1,141.00
2012	1,570.00	2,510.00	2,248.00	2,014.00	1,389.00	1,197.00

2013	2,524.60	2,682.00	2,628.00	2,309.00	2,270.00	3,573.00
2014	2,908.00	2,388.00	2,628.00	2,025.90	1,980.20	1,521.00
2015	2,169.50	2,199.30	1,620.70	2,045.50	2,024.70	1,310.10
2016	2,711.50	3,549.10	2,590.00	3,030.10	2,976.80	1,807.70
2017	2,152.10	2,297.30	2,657.00	2,596.00	2,123.80	1,762.30
2018	1,501.60	2,982.80	1,995.60	2,488.30	1,772.50	1,634.70
2019	1,560.00	3,555.90	1,249.40	2,121.40	1,884.80	1,606.90

The results of calculating the rainfall for a 24 hour period using the Gumbel Distribution formula, obtained a coefficient Y_n , S_n , and Y_t . From Table 2, information is obtained that the average for a longer period will produce a greater value, using data from 6 provinces over a 15 years period.

Table 2. 24 hour Rainfall Calculation

	Y_n	S_n	Y_t			
			5 Years	10 Years	25 Years	50 Years
N = 15	0.5128	1.0206	1.4999	2.2562	3.1985	3.9019

Table 3 is the result of the analysis of the frequency of daily rain which is processed through Table 2. The calculation results show that the largest increase occurred in the period 10 to 25 years.

Table 3. Results of Daily Rain Frequency Analysis

Province	Daily Rainfall Return Period (mm)			
	5 Years	10 Years	25 Years	50 Years
Jakarta	2,562.38	3,103.45	3,777.58	4,280.80
West Java	3,183.79	3,717.70	4,382.91	4,879.48
Central Java	2,797.12	3,287.16	3,897.72	4,353.48
SR Yogyakarta	2,517.31	2,893.78	3,362.83	3,712.96
East Java	2,510.62	2,962.17	3,524.78	3,944.76
Banten	2,216.99	2,788.31	3,500.14	4,031.50

Then, the rain intensity obtained from the results of the approach with the Mononobe formula is shown in Table 4 and in Table Appendix 1 to Table Appendix 6. Furthermore, analysis of rain intensity is carried out in different time durations or less than 24 hours (daily).

Table 4. Daily Rain Intensity For Several Return Rain Periods

Rainy Return Period	Rain Intensity (mm / 24 hours)					
	Jakarta	West Java	Central Java	SR Yogyakarta	East Java	Banten
5 Years	106.77	132.66	116.55	104.89	104.61	92.37
10 Years	129.31	154.90	136.97	120.57	123.42	116.18
25 Years	157.40	182.62	162.40	140.12	146.87	145.84
50 Years	178.37	203.31	181.39	154.71	164.36	167.98

The daily rainfall threshold (24 hours) based on the Meteorology, Climatology and Geophysics Agency (BMKG) is divided into several categories, namely:


Table 5. Category of Daily Rainfall Intensity

Rainfall Range	Category
0 mm/day	Cloudy
0,5 – 20 mm/day	Light rain
20 – 50 mm/day	Moderate rain
50 – 100 mm/day	Heavy rain
100 – 150 mm/day	Very Heavy Rain
>150 mm/day	Extreme Rain

The daily rainfall intensity for six provinces is in the range of around 100-200 mm / day. Based on the daily rainfall threshold according to the Meteorology, Climatology and Geophysics Agency (BMKG), the intensity of rain in Java is in the category of very heavy rain and extreme rain. The return period of rain in Java for 5 years and 10 years is included in the very heavy rain category, and for the return period of rain in Java for 25 years and 50 years, it is included in the extreme rain category.

The decrease in rain intensity can be seen using the IDF curve analysis. The IDF curve is a curve that presents the relationship between the frequency, intensity and duration of rain which is expressed in terms of curved rain intensity with a certain return period. IDF curve creation can be done from the results of the automatic rain data frequency analysis (duration minutes and hours). If automatic data is not available, IDF can be derived based on daily data analysis using an approximate formula.

The duration and results of rainfall intensity analysis for each return period use the Monobebe formula approach, which is then depicted in Cartesian coordinates. The abscissa axis (X) represents duration, while the ordinate axis (Y) represents intensity. The results of the depiction of the data, hereinafter referred to as the IDF Curve.

**Figure 1.** Intensity-Duration-Frequency (IDF) Curve

Rain and high rain intensity generally lasts for a short duration and covers a small area. Rain covering a large area, rarely with high intensity, but can last quite a long duration. The combination of high rainfall intensity with long duration is rare. The amount of rainfall intensity is not the same for each province. It can be seen from the IDF curve that West Java Province has a fairly high rain intensity for one hour

duration compared to the other 5 provinces. Rain intensity with a duration of five to eight hours from six provinces in Java Island tends to have almost the same rain intensity. The difference in rainfall intensity in the six provinces is influenced by the topography, duration and frequency of the place or location concerned.

In this study, the formula for the best empirical approach has not been determined to calculate the intensity of rainfall in each province in Java. This is due to the unavailability of hourly rain data to validate the calculated rainfall intensity.

4. Conclusion

Rain intensity and IDF curve depiction in Java with the Mononobe method have been successfully analyzed in this study. Analysis was carried out on the intensity, duration and frequency of rain events from 2005-2019. The results of this study indicate that high intensity rain occurs in a short duration. There are differences in the value of rain intensity between provinces in Java Island. However, this study has not been able to decide whether this method is suitable for use in Java. This is due to the limited available rain data. Further research can validate the results of rainfall intensity calculations with other approaches, through hourly rainfall record data, so that we can determine whether the rainfall intensity approach formula used for an area is appropriate.

References

- [1] P. Aryastana, T. Tanaka, and M. S. Mahendra, "Characteristic of rainfall pattern before flood occur in Indonesia based on rainfall data from GSMaP," *Ecotrophic J. Environ. Sci.*, vol. 7, no. 2, pp. 100–110, 2012.
- [2] M. S. Lisboa, R. L. Schneider, P. J. Sullivan, and M. T. Walter, "Drought and post-drought rain effect on stream phosphorus and other nutrient losses in the Northeastern USA," *J. Hydrol. Reg. Stud.*, vol. 28, p. 100672, 2020.
- [3] F. Faradiba, "Peramalan curah hujan dan luas serangan organisme pengganggu tanaman di Kabupaten Bogor," *Pro-Life*, vol. 5, no. 3, pp. 688–699, 2018.
- [4] T. W. Hertel and C. Z. de Lima, "Climate impacts on agriculture: Searching for keys under the streetlight," *Food Policy*, vol. 95, p. 101954, 2020.
- [5] W. B. Setyawan and A. Pamungkas, "Perbandingan karakteristik oseanografi pesisir utara dan selatan Pulau Jawa: pasang-surut, arus, dan gelombang," in *Prosiding Seminar Nasional Kelautan dan Perikanan III*, 2017, pp. 191–202.
- [6] BMKG, "Probabilistik curah hujan," Jakarta, 2018.
- [7] R. Gernowo and D. N. Sugianto, "Model evaluasi iklim maritim tropis berbasis sistem inferensi fuzzy jaringan saraf adaptif," *ILMU Kelaut. Indones. J. Mar. Sci.*, vol. 9, no. 2, pp. 115–119, 2012.
- [8] M. Hernando, D. E. Varela, G. Malanga, G. O. Almandoz, and I. R. Schloss, "Effects of climate-induced changes in temperature and salinity on phytoplankton physiology and stress responses in coastal Antarctica," *J. Exp. Mar. Bio. Ecol.*, vol. 530, p. 151400, 2020.
- [9] X. Li, J. Sardans, L. Hou, M. Liu, C. Xu, and J. Peñuelas, "Climatic temperature controls the geographical patterns of coastal marshes greenhouse gases emissions over China," *J. Hydrol.*, vol. 590, p. 125378, 2020.
- [10] B. Munandar, P. Purwanto, and K. Kunarso, "Kaitan monsun terhadap variabilitas suhu permukaan laut dan klorofil-a untuk prediksi potensi fishing ground di Perairan Karimunjawa," *J. Oceanogr.*, vol. 5, no. 4, pp. 505–511, 2016.
- [11] W. Soon and D. R. Legates, "Solar irradiance modulation of Equator-to-Pole (Arctic) temperature gradients: Empirical evidence for climate variation on multi-decadal timescales," *J. Atmos. Solar-Terrestrial Phys.*, vol. 93, pp. 45–56, 2013.
- [12] S. K. K. Reddy, H. Gupta, B. Upendra, D. V. Reddy, R. M. Kurakalva, and D. Kumar, "Export of particulate organic carbon by the mountainous tropical rivers of Western Ghats, India:

- Variations and controls,” *Sci. Total Environ.*, p. 142115, 2020.
- [13] A. Konishi, T. Fujiike, O. Okano, H. Chiba, and A. Ueda, “Geochemical and isotopic imprints of groundwater evolution in mountainous areas of Maniwa City, Okayama Prefecture, Japan,” *Groundw. Sustain. Dev.*, p. 100412, 2020.
- [14] J. Susanto, X. Zheng, Y. Liu, and C. Wang, “The Impacts of Climate Variables and Climate-related Extreme Events on Island Country’s Tourism: Evidence from Indonesia,” *J. Clean. Prod.*, p. 124204, 2020.
- [15] C. Jung and D. Schindler, “Changing wind speed distributions under future global climate,” *Energy Convers. Manag.*, vol. 198, p. 111841, 2019.
- [16] G. Roshan, M. Moghbel, and S. Attia, “Evaluating the wind cooling potential on outdoor thermal comfort in selected Iranian climate types,” *J. Therm. Biol.*, vol. 92, p. 102660, 2020.
- [17] Y. Iwata, M. Nemoto, S. Hasegawa, Y. Yanai, K. Kuwao, and T. Hirota, “Influence of rain, air temperature, and snow cover on subsequent spring-snowmelt infiltration into thin frozen soil layer in northern Japan,” *J. Hydrol.*, vol. 401, no. 3–4, pp. 165–176, 2011.
- [18] N. Wu *et al.*, “Contrasting frontal and warm-sector heavy rainfalls over South China during the early-summer rainy season,” *Atmos. Res.*, vol. 235, p. 104693, 2020.
- [19] K. Schollaen *et al.*, “Multiple tree-ring chronologies (ring width, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) reveal dry and rainy season signals in Indonesia,” *EGUGA*, p. 13529, 2013.
- [20] P. D. Roy *et al.*, “Depositional histories of vegetation and rainfall intensity in Sierra Madre Oriental Mountains (northeast Mexico) since the late Last Glacial,” *Glob. Planet. Change*, vol. 187, p. 103136, 2020.
- [21] D. A. Sofia and N. Nursila, “Analisis intensitas, durasi, dan frekuensi kejadian hujan di wilayah Sukabumi,” vol. 4, no. 1, pp. 85–92, 2019, doi: 10.31544/jtera.v4.i1.2019.85-92.
- [22] E. Yulius, “Analisa curah hujan dalam membuat kurva intensity duration frequency (IDF) pada DAS Bekasi,” *Bentang*, vol. 2, no. 1, 2014.
- [23] V. Agilan and N. V. Umamahesh, “Is the covariate based non-stationary rainfall IDF curve capable of encompassing future rainfall changes?,” *J. Hydrol.*, vol. 541, pp. 1441–1455, 2016.
- [24] C. H. R. Lima, H.-H. Kwon, and Y.-T. Kim, “A local-regional scaling-invariant Bayesian GEV model for estimating rainfall IDF curves in a future climate,” *J. Hydrol.*, vol. 566, pp. 73–88, 2018.
- [25] P. J. Laksana, “Rainfall characteristic on the slopes of Mount Merapi Region (Empirical formula, duration, distribution, and critical line Woro River),” in *Journal of the civil engineering forum*, 2015, vol. 1, no. 2, pp. 43–50.
- [26] S. Fauziyah, S. Sobriyah, and S. Susilowati, “Analisis karakteristik dan intensitas hujan Kota Surakarta,” *Matriks Tek. Sipil*, vol. 1, no. 2, 2013.
- [27] Wesli, “Drainase Perkotaan,” *Yogyakarta Graha Ilmu*, 2008.
- [28] R. I. Harris, “Gumbel re-visited—a new look at extreme value statistics applied to wind speeds,” *J. Wind Eng. Ind. Aerodyn.*, vol. 59, no. 1, pp. 1–22, 1996.
- [29] Y.-S. Fang and T.-J. Chen, “Modification of Mononobe–Okabe theory,” *Geotechnique*, vol. 45, no. 1, pp. 165–167, 1995.