

DAFTAR PUSTAKA

1. Rusdi NK, Trisna Y, Soemiati A. Pola pengobatan flour albus di Rumah Sakit Umum Pusat Nasional DR Cipto Mangunkusumo serta faktor-faktor yang mempengaruhinya (analisis data rekam medik tahun 2006-2007). *Pharma Scien Research*. 2008;V(2):91-100.
2. Emilia Y, Suling PL, Pandeleg HEJM. Profil duh tubuh vagina di Poliklinik Kulit dan Kelamin RSUP Prof. DR. R.D Kandou Manado Tahun 2009-2011.J *eClinic unsrat*.2013;1(1):670-5.
3. Himalaya D. Pengaruh pemberian ekstrak biji manjakani (*quercus infectoria gall*) terhadap bakteri vaginosis dan candida penyebab keputihan (leukorhea).J *Midwifery*.2017;5(1):38-44.
4. Andalas M, Syukri A, Azhar A. Gambaran penderita flour albus di Poliklinik Gynekologi RSUD Dr. Zainoel Abidin Banda Aceh periode Agustus 2011.J *Kedokteran Syiah Kuala*. 2011;11(3):143-50.
5. Achdiat PA, Dwiwana RF, Feriza V, Rowawi R, Effendi RA, Suwarsa O, et al. Prevalence of trchomoniasis in asymptomatic pregnant women populationin Bandung, West Java. *Indo J Tropic Infect Dis*. 2019;7(4):138-42.
6. Susanto I, Ismid IS, Sjarifuddin PK, Sungkar S. Mastigopora flagelata traktus digestif dan traktus urogenital: *Trichomonas vaginalis*. in : *Buku Ajar Parasitologi Kedokteran*. 4th ed. Jakarta: Badan Penerbit FK UI; 2016.h;138-42.
7. Olowe OA, Makanjuola OB, Olowe R, Adekanle DA. Prevalence of vulvovaginal candidiasis, trichomonas and bacterial vaginosis among pregnant women receiving antenatal care In Southwestern Nigeria. *Eur J Microbiol Immunol*. 2014;4(4):193-7.
8. Goo Y, Shin W, Yang H, Joo S, Song S, Ryu J, et al. Prevalence of *Trichomonas vaginalis* in women visiting 2 obstetrics and gynecology clinics in Daegu, South Korea. *Korean J Parasitol*. 2016;1:75-80.

9. Crucitti T, Jespers V, Mulenga C, Khondowe S, Vandepitte J, Buve A. *Trichomonas vaginalis* is highly prevalent in adolescent girls, pregnant women, and commercial sex workers in Ndola, Zambia. *J Amer Sex Transm Dis*. 2010;37(4):223-7.
10. Nourian A, Shabani N, Fazaeli A, Mousavinasab SN. Prevalence of *Trichomonas vaginalis* in pregnant women in Zanjan, Northwest of Iran. *Jundishapur J Microbiol*. 2013;6(8):58-75.
11. Miranda AE, Pinto VM, Gaydos CA. *Trichomonas vaginalis* infection among young pregnant women in Brazil. *Braz J Infect Dis*. 2014;18(6):669-71.
12. Deivam S, Rajalakshmi R, Priyadarshini S, Seethalaksmi RS, Balasubramanian N, Brinda T, et al. Prevalence of *Trichomonas vaginalis* infection among patients that presented to rural tertiary care Hospital in Tiruchirapalli, India in 2011 and 2013. *Internas J Pharma Research Health Scien*.2013;2(3):255-9.
13. Coleman JS, Gaydos CA, Witter F. *Trichomonas vaginalis* vaginitis in obstetrics and gynecology practice: New concepts and controversies. *Obstet Gynecol Surv*. 2013;68(1):43-8.
14. Astuti R. Gambaran status gizi dan asupan zat gizi pada ibu hamil di kota Semarang. *J Nutr Health*. 2019,7(1): 40-5.
15. Donbraye E, Donbraye EOOB, Okonko IO, Okedeji IO, Alli JA, Nwanze JC. Detection and prevalence of *Trichomonas vaginalis* among pregnant women in Ibadan, Southwestern Nigeria. *World Appl Scien J*. 2010;11(12):1512-17.
16. Allsworth JE, Ratner JA, Peipert JF. Trichomoniasis and other sexually transmitted infections: results from the 2001-2004 National Health and Nutrition Examination Surveys. *J Sex Transm Dis*.2009;36(12):738-44.
17. Etuketu IM, Mogaji HO, Alabi OM, Adeniran AA, Oluwole AS, Ekpo UM. Prevalence and risk factors of *Trichomonas vaginalis* infection among pregnant women receiving antenatal care in Abeokuta, Nigeria. *J Infect Dis*. 2015;9(2):51-55.
18. Sastroasmoro S, Ismael S. Perkiraan besar sampel. in: *Dasar-dasar Metodologi Penelitian Klinis*. 5th ed. Jakarta : Sagung Seto;2011.h;348-52.

19. Nouraddin AS, Alsakee HM. Prevalence of *Trichomonas Vaginalis* infection among women in Erbil, Governorate, Northern Iraq: An Epidemiological Approach. Eur Scient J. 2015;11(24): 243-55.
20. Romoren M, Velauthapillai M, Rahman M, Sundby J, Klouman E, Hjortdahl P. Trichomoniasis and bacterial vaginosis in pregnancy: inadequately managed with the syndromic approach. Bull World Health Organ. 2007;85(4):297-304.