

# Ethnobotanical Study of Oke Sou: Traditional Herbal Drink from Lako Akediri Village in West Halmahera, Indonesia

*by* Marina Silalahi

---

**Submission date:** 26-May-2020 03:11PM (UTC+0700)

**Submission ID:** 1332035135

**File name:** ou\_Traditional\_Herbal\_Drink\_from\_West\_Halmahera\_2C\_Indonesia.pdf (756.05K)

**Word count:** 3794

**Character count:** 21575

## Ethnobotanical Study of *Oke Sou*: Traditional Herbal Drink from Lako Akediri Village in West Halmahera, Indonesia

Anisatu Z. Wakhidah<sup>1\*</sup>, Marina Silalahi<sup>2</sup>, Nisyawati<sup>3</sup>

<sup>1</sup>Student & researcher, Department Biology, Faculty of Mathematics and Science, Universitas Indonesia

<sup>2</sup>Lecture & researcher, Faculty of Education & Teacher Training, Department of Biology Education, Universitas Jember Indonesia,

<sup>3</sup>Senior Lecture & researcher, Department Biology, Faculty of Mathematics and Science, Universitas Indonesia

### Abstract

*Oke sou* is a herbal drink from Lako Akediri village West Halmahera, to maintain health of women reproductive function. This drink is consumed when women get their first menstruation. This is the first study of *Oke sou* to document all plant species used in preparing the herbal drink. It describes phytochemical content of the most cited plant based on the study of literature. Ethnobotanical data were collected using semi-structured interviews individually on indigenous medical practitioner and a local woman (30 respondents). Plant specimens were collected from the habitat, made into herbarium voucher, and then identified. Recorded as many as 66 plant species from 59 genera belonging to 37 families are used making the *oke sou*. The most frequently mentioned plants (>5 respondents) are (number of respondents; part used), *Cananga odorata* (Lam.) Hook.f. & Thomson (10; bark), *Surcuma longa* L. (8; rhizome), *Cymbopogon citratus* (DC.) Stapf. (7; stem), *Kaempferia galanga* L. (7; rhizome), *Myristica fragrans* Houtt. (7; fruit and seeds), *Syzygium aromaticum* (L.) Merr. & L.M. Perry (7; leaf & flower), *Cynometra cauliflora* L. (6; bark), and *Tamarindus indica* L. (6; bark). These plants are already well studied regarding phytochemical content in maintaining women's reproductive health. Therefore, the results of this study can be used as a reference for the development of medical products based on local knowledge.

**Keywords:** ethnobotany, oke sou, herbal drink, women, west halmahera

### Introduction

Traditional herbal drinks in Indonesia is still used till today. These drinks have become a part of living culture to maintaining body health or beauty care, such as *jamu*. *Jamu* is traditional herbal drink from Java that have been used for a long time. This herbal drink can consist of a single or mixture some medical plants<sup>(17)</sup>. It is used to treat some diseases and to maintain good health.

Beside in Java, traditional herbal drink also found in another region, for example Bali. Based on Sujarwo dkk. (2015), the Bali community, especially in ancient villages, still produce and consume *lolo* to prevent and treat different ailments. *Lolo* are the most common herbal drinks in Bali which generally prepared as decoctions of some medical plants.

Traditional herbal drinks are also found in the eastern region of Indonesia, precisely at Lako Akediri village, District West Halmahera, North Moluccas. This herbal drink is made from various kind of plants at Lako Akediri village. It is believed efficacious to maintain health of girl reproductive function and to eliminate body odor of girl. The community at Lako Akediri call that herbal drink by the name *oke sou*.

In Indonesia, traditional medicine knowledge passed down orally<sup>(4)</sup>. This is also occurred on inheritance of knowledge composition plants used in *oke sou* herbal drink. Inheritance knowledge by oral is highly vulnerable to losing traditional medicine knowledge because of no one documentation can be inherited<sup>(22)</sup>. Moreover, research on *oke sou* herbal drink has not ever been implemented. Therefore, an inventory about its diversity plant species is quite important to be conducted.

There are two objectives of this research. First, to inventory all plant species used in preparing *oke sou* herbal drink. Second, to describe and to explain phytochemical content of the most frequently mentioned plants used in *oke sou* herbal drink at Lako Akediri village, West Halmahera, North Moluccas - Indonesia.

Contact Author:

Anisatu Z. Wakhidah

Bachelor of Science, Department Biology, Faculty of Mathematics and Science, Universitas Indonesia

Depok 16424, Indonesia

Tel: +62-897-6113-393 Fax: -

Email: khistia.nisa@gmail.com

## Material and Methods

### Study Area

The study was conducted at Lako Akediri Village (Fig. 1), on May - June 2014 and October 2014. Lako Akediri Village (Fig. 2) is geographically located on coastal area  $27^{\circ}22'17.323''$  – E  $127^{\circ}37'5.214''$  and N  $0^{\circ}58'13.505''$  – N  $1^{\circ}8'5.332''$ . Total area of Lako Akediri Village is 10 hectares, which located at an altitude 31 meters above sea level with average rainfall 15 mm/month<sup>(16)</sup>. The population in 2014 was 344 people; 175 males and



Fig 1. Study site at Lako Akediri Village Sub District Sahu, District West Halmahera, North Moluccas - Indonesia



Fig 2. The Lako Akediri Village on Sub District Sahu

169 females, with number of households were 85. As many as 98% community at Lako Akediri Village come from Sahu tribe, while the rest are ethnic immigrants, such as Buton, Bugis, and Sasak. The agricultural plants at Lako Akediri Village are tubers, corn, coconuts, clove, and nutmeg<sup>(16)</sup>.

### Ethnobotanical Data Collection

The ethnobotanical data in this research consists of interview result and list of used plant species. Interview data were collected using semi-structured interviews method that conducted individually on key respondent and general respondent. Key respondent is person who considered having more knowledge about *oke sou* herbal drink, such as indigenous medical practitioner. General respondent is local woman who ever participated in producing *oke sou* herbal drink with age range of 12 - 40 years old (30 respondents).

Plant specimens were collected from the habitat together with key respondent. Then plant specimens were made into herbarium voucher. After that specimens were identified at Laboratory of Plant Taxonomy in Department Biology Universitas Indonesia.

### Data Analysis

Data were analyzed using quantitative and qualitative approaches. Quantitative analysis was conducted to obtain total plants species and families, also to know the most frequently mentioned plant and part used in *oke sou* herbal drink. Qualitative analysis aimed to determine phytochemical content in most frequently mentioned plants used in *oke sou* herbal drink at Lako Akediri village.

## Result

### Plant species used in “oke sou” herbal drink

The investigation recorded as many as 66 plant species from 59 genera used for preparation *oke sou* herbal drink (Table 1). These plants belong to 37 families which are Acanthaceae, Fabaceae, and Lamiaceae being the most represented family (6 plant species each family). There are eight plant species that most frequently mentioned by respondents (plant species; part used), *Cananga odorata* (Lam.) Hook.f. & Thomson (bark), *Curcuma longa* L. (rhizome), *Cymbopogon citratus* (DC.) Stapf. (stem), *Kaempferia galanga* L. (rhizome), *Myristica fragrans* Houtt. (fruit and seeds), *Syzygium aromaticum* (L.) Merr. & L.M. Perry (leaf & flower), *Cynometra cauliflora* L. (bark), and *Tamarindus indica* L. (bark).

2

The plant parts, which are harvested to prepare *oke sou* herbal drink, are bark, leaves, stems, rhizomes, flowers, fruits, seeds (Fig. 3). Bark is being the most frequently used part in preparing *oke sou* herbal drink (31 species). Meanwhile, the least frequently used part is rhizomes. The data showed that aerial parts (79%) are preferred than underground parts (21%). This may be because of the easier accessibility in picking plant source and the greater quantity of aerial parts than underground parts <sup>(1)(7)</sup>.




Fig 3. Number of species and plant part used in preparing *oke sou* herbal drink

### Preparation of “oke sou” herbal drink

The preparation of *oke sou* herbal drink is started by classifying the same part of used plants, such as leaves with leaves, bark with bark, or root with root. Then, each group is crushed separately. The collisions were given water and then squeezed in a clean cloth - same as the technique in making juice-. After that, the juice of each part of used plants is all mixed, then boiled until boiling. When boiled, *oke sou* herbal drink is mixed with herb spices to improve the acceptability of this herbal drink. Usually the herb spices that used are *Coriandrum sativum* L., *Piper nigrum* L., *Curcuma longa* L., *Zingiber officinale* Roscoe., *Cymbopogon citratus* (DC.) Stapf., *Kaempferia galanga* L., *Myristica fragrans* Houtt., and *Syzygium aromaticum* (L.) Merr. & L.M.Perry. The *oke sou* herbal drink is ready to be consumed when its color becoming as brown as the color of strong tea.

*Oke sou* herbal drink is only taken by girl when getting her first menstruation in tradition ceremony welcoming maturity girl. The tradition is held for 3, 7, or 9 days depending on length of menstruation period and decision of the girl's family. During that time, the girl takes *oke sou* herbal drink 3 times for a day. Usually the girl drinks as much as 8.1 liters of *oke sou* herbal drink as her lifetime.

Table 1. Plant used in preparing *oke sou* herbal drink at Lako Akediri Village Sub District Sahu, District West Halmahera, North Moluccas - Indonesia. The life form, vernacular name, part used, and number informant also provided.

Plant families and species	Life form	Vernacular Name	Part Used	Number Informant
<b>Family: Acanthaceae</b>			16	
<i>Graptophyllum pictum</i> (L.) Griff	clump	kabi-kabi merah	leaf	1
<i>Graptophyllum pictum</i> 'roseum variegatum'	clump	kabi-kabi putih	leaf	1
<i>Hemigraphis alternata</i> (Burm. F) T. Anderson	herb	lire buntal (♀)	leaf	4
<i>Hemigraphis rependa</i> (L.) Hall. F	herb	lire panjang (♂)	leaf	1
<i>Justicia gendarussa</i> Burm. F.w	herb	gandarusa	leaf	3
<i>Ruellia simplex</i> C. Wright.	herb	Puli	leaf, stem	2
<b>Family: Anacardiaceae</b>				
<i>Mangifera</i> sp.	tree	mangga dodol	bark	5
<b>Family: Annonaceae</b>				
<i>Annona muricata</i> L.	tree	nangka belanda	bark	5
<i>Cananga odorata</i> (Lam.) Hook.f. & Thomson	tree	kenanga	bark	10
<b>Family: Apiaceae</b>				
<i>Coriandrum sativum</i> L.	herb	surai	leaf, seed	2
<b>Family: Apocynaceae</b>				
<i>Alstonia scholaris</i> R. Br.	tree	hange	bark	4
<b>Family: Asteraceae</b>				
<i>Blumea balsamifera</i> (L.) DC.	herb	madikapu	leaf	1
<i>Wollastonia biflora</i> (L.) DC.	herb	cinga-cinga	leaf	1

Table 1. (continued)

Plant families and species	Life form	Vernacular Name	Part Used	Number Informant
<b>Family: Bombacaceae</b>	7		8	
<i>Durio zibethinus</i> L.	tree	durian	bark	3
<b>Family: Burseraceae</b>				
<i>Canarium umbonense</i> Hoch.	tree	kenari	bark	1
<b>Family: Clusiaceae</b>				
<i>Garcinia mangostana</i> L.	tree	manggis	bark	5
<b>Family: Combretaceae</b>				
<i>Terminalia catappa</i> L.	tree	ngusu	bark	2
<b>Family: Commelinaceae</b>				
<i>Tradescantia spathacea</i> Sw.	herb	bia-bia	leaf, flower	3
<b>Family: Convolvulaceae</b>				
<i>Merremia peltata</i> (L.) Merr.	herb	koge	bark	1
<b>Family: Cyperaceae</b>				
<i>Scleria</i> sp.	shrub	cakagole	bark	1
<b>Family: Euphorbiaceae</b>				
<i>Homalanthus novoguineensis</i> (Warb.) K. Schum.	tree	gidilule	bark	1
<i>Jatropha curcas</i> L.	clump	balacai putih	leaf	5
<i>Macaranga tanarius</i> (L.) Müll.Arg.	tree	same	bark	1
<i>Mallotus apelta</i> (Lour.) Müll.Arg.	tree	lufiti	leaf	4
<b>Family: Fabaceae</b>				
<i>Albizia saponaria</i> (Lour.) Miq	tree	fau-fau	bark	1
<i>Cynometra cauliflora</i> L.	tree	mano mano	bark	6
<i>Pongamia pinnata</i> (L.) Pierre	tree	hatehira	bark	4
<i>Pterocarpus indicus</i> Wild.	tree	ligua	bark	2
<i>Sesbania grandiflora</i> Pers.	tree	Turi	bark, leaf	2
<i>Tamarindus indica</i> L.	tree	asam jawa	bark, leaf	6
<b>Family: Lamiaceae</b>				
<i>Callicarpa rubella</i> Lindl.	herb	ngaai madudera	bark	4
<i>Coleus scutellarioides</i> Bth.	herb	mayana	leaf	5
<i>Leucas zeylanica</i> (L.) R.Br.	herb	gofu hairani	leaf	2
<i>Orthosiphon grandiflorus</i> Bold.	shrub	kumis kucing	leaf	4
<i>Premna serratifolia</i> (Blanco) Benth.	tree	gumira	bark	3
<i>Vitex pinnata</i> L.	tree	gofasa	bark	2
<b>Family: Lauraceae</b>				
<i>Cassytha</i> f. filiformis	climber	tali kuning	stem	2
<b>Family: Lygodiaceae</b>				
<i>Lygodium</i> sp.	herb	gumoho	leaf	1
<b>Family: Magnoliaceae</b>				
<i>Michelia champaca</i> L.	tree	cempaka	bark	4
<b>Family: Malvaceae</b>				
<i>Kleinhovia hospita</i> L.	tree	liwui	bark	2
<b>Family: Meliaceae</b>				
<i>Xylocarpus moluccensis</i> (Lam.) M. Roem	tree	lolesou	bark	3
<b>Family: Moraceae</b>				
<i>Ficus</i> cf. <i>ribes</i>	tree	senang	leaf	1
<i>Ficus fistulosa</i> Reinw. Ex Blume	tree	Coro	bark	4



Table 1. (continued)

Plant families and species	Life form	Vernacular Name	Part Used	Number Informant
<i>Ficus hispida</i> Linn.	tree	tagalolo	bark	3
<b>Family: Myristicaceae</b>				
<i>Myristica fragrans</i> Houtt.	tree	Pala	fruit, seed	7
<b>Family: Myrtaceae</b>				
<i>Psidium guajava</i> L.	tree	giawas	leaf	6
<i>Syzygium aqueum</i> (Burm.f.) Alston.	tree	gora	leaf	3
<i>Syzygium aromaticum</i> (L.) Merr. & L.M. Perry	tree	cengkeh	leaf, flower	7
<b>Family: Oxalidaceae</b>				
<i>Averrhoa bilimbi</i> L.	shrub	belimbing wuluh	bark, fruit	2
<b>Family: Phyllanthaceae</b>				
<i>Breynia cernua</i> (Poir.) Müll.Arg.	tree	gagilamo	bark	3
<i>Phyllanthus</i> sp.	herb	balakama seed	leaf	3
<b>Family: Piperaceae</b>				
<i>Piper nigrum</i> L.	climber	rica jawa	fruit	2
<i>Piper sarmentosum</i> Roxb.	herb	tofure	leaf	1
<b>Family: Poaceae</b>				
<i>Cymbopogon citratus</i> (DC.) Stapf.	herb	gramakusu	stem	7
<b>Family: Ranunculaceae</b>				
<i>Nigella sativa</i> Linn.	herb	jinta hitam	fruit	5
<b>Family: Rhamnaceae</b>				
<i>Alphitonia moluccana</i> Teijsm. & Binn. Ex Brais.	tree	raurika	bark	2
<b>Family: Rubiaceae</b>				
<i>Morinda citrifolia</i> L.	tree	kome	bark	1
<b>Family: Rutaceae</b>				
<i>Melicope latifolia</i> (DC.) T.G. Hartley	shrub	sawuyo	leaf	2
<b>Family: Selaginellaceae</b>				
<i>Selaginella</i> sp.	herb	rutu-rutu	leaf	4
<b>Family: Solanaceae</b>				
<i>Physallis peruviana</i> L.	herb	dagameme	leaf	4
<b>Family: Sonneratiaceae</b>				
<i>Sonneratia alba</i> Sm.	tree	posi-posi / soki bulat	bark	3
<b>Family: Zingiberaceae</b>				
<i>Boesenbergia rotunda</i> (L.) Mansf.	herb	tumbukunci	rhizome	1
<i>Curcuma longa</i> L.	herb	kuning	rhizome	8
<i>Curcuma zanthorrhiza</i> Roxb.	herb	tumbulawak	rhizome	1
<i>Kaempferia galanga</i> L.	herb	bataka	rhizome	7
<i>Zingiber officinale</i> Roscoe.	herb	guraka	rhizome	4

### Phytochemical profile

All the most frequently mentioned plants are well studied and have documented phytochemical profile also pharmacological activities (Table 2.). The common pharmacological activities related to efficacy of *oke sou* herbal drink are antimicrobial, antifungal, aromatherapy, antioxidant, and anti-cancer. *Canangan odorata* (Lam.) Hook.f. & Thomson has 65 different chemical compound have

been isolated with more than 13 pharmacological activities. This plant is effective to maintain cleanness vagina area due to its antimicrobial activity that contains essential oil, ethyl acetate ethanolic, methanolic, cyclohexane, and clorofrom<sup>(25)</sup>. The other plants that also contain antimicrobial activity are *Kaempferia galanga* L., *Syzygium*

*aromaticum* (L.) Merr. & L.M. Perry. and *Tamarindus indica* L.<sup>(2)(6)(16)(27)</sup>.

Some plants have essential oil which efficacious to reduce body odor, such as camphene (*Cananga odorata* (Lam.) Hook.f. & Thomson)<sup>(26)</sup>, geraniol (*Cymbopogon citratus* (DC.) Stapf.)<sup>(28)</sup>, myristicin (*Myristica fragrans* Houtt.)<sup>(10)</sup>, and

eugenol (*Syzygium aromaticum* (L.) Merr. & L.M. Perry.)<sup>(10)</sup>. These chemical compound have pharmacological activities as aromatherapy and become basic material in perfume producing. Based on the data (Table 2) there are antifungal activity in some plants used in *oke sou* herbal drink which effective against *Candida albicans* activity, vaginal discharge agent<sup>(26)</sup>.

20

**Table 2.** The most frequently mentioned plant species (>5 respondents) to prepare *oke sou* herbal drink and their phytochemical profile and pharmacological activities at Lako Akeriri Village Sub District Sahu, District West Halmahera, North Moluccas - Indonesia.

Plant Species	Phytochemical profile	Pharmacological activities
12 <i>Cananga odorata</i> (Lam.) Hook.f. & Thomson	bornyl acetate (leaves); camphene (leaves, flowers); geraniol (leaves, flowers); geranyl acetate (flowers); limonene (leaves, flowers, fruits); (E,Z)-farnesal (leaves) 1-epi-cubenol (flowers); caryophyllene epoxide (leaves); spathulenol (leaves); <i>r</i> -cadinol (leaves); $\alpha$ -amorphene (leaves, flowers); $\alpha$ -ylangene (leaves, flowers); methyl anthranilate (flowers) <sup>(26)</sup> ; lirioidenine, sampangine (bark) <sup>(19)</sup> ; methylisoeugenol, benzyl benzoate (flower) <sup>(17)</sup>	aromatherapy, anti-microbial, anti-inflammatory, antivector <sup>(26)</sup>
<i>Curcuma longa</i> L.	curcumin; dimethoxy curcumin; bisdemethoxy curcumin; sodium curcumin (rhizomes) <sup>(19)(9)</sup>	antifungal, anti-mycobacterial, antimalarial <sup>(28)(13)</sup>
<i>Cymbopogon citratus</i> (DC.) Stapf.	d-Limonene, geraniol (leaves) <sup>(28)</sup> ; $\alpha$ -citral, $\beta$ -neral, m <sup>19</sup> ene (leaves) <sup>(13)</sup>	anti-carcinogenic <sup>(8)</sup> anti-bacteria, anti-HIV, antioxidant, anti-inflammatory, anti-tumor <sup>(9)</sup>
<i>Kaempferia galanga</i> L.	$\alpha$ -pinene, camphene, carvone, benzene, eucalypto, borneol, methyl cinnamate, ethyl- <i>p</i> -methoxycinnate (rhizomes) <sup>(26)</sup> ; $\beta$ -phyllandrene, $\alpha$ -terpineol, ethylcinnate, dihydro $\beta$ -sesquiphylandrene (rhizomes) <sup>(21)</sup>	anticancer, antimicrobial activity, antioxidant <sup>(27)</sup>
9 <i>Myristica fragrans</i> Houtt.	macelignan (fruits) <sup>(5)</sup> ; ethanolic (seeds) <sup>(26)</sup> ; myristicin (fruits) <sup>(10)</sup> ; malabaricone B, malabaricone C (fruits) <sup>(14)</sup>	anti-bacterial <sup>(5)</sup> ; aphrodisiac <sup>(25)</sup> ; antifungal <sup>(14)</sup> ; aromatherapy <sup>(10)</sup>
<i>Syzygium aromaticum</i> (L.) Merr. & L.M. Perry.	eugenol, eugenyl acetate, benzyl alcohol (leaves) <sup>(12)(15)</sup> , ethanolic (seeds) <sup>(25)</sup>	antioxidant <sup>(12)</sup> ; antimicrobial, antifungal <sup>(15)</sup> ; aphrodisiac <sup>(24)</sup>
<i>Cynometra cauliflora</i> L.	methanolic (fruits) <sup>(26)</sup> ; tannin, saponin, flavonoid (leaves, stems, barks); terpenoid (leaves, stems) <sup>(5)</sup>	anti-cancer <sup>(26)</sup> ; antioxidant <sup>(4)</sup>
<i>Tamarindus indica</i> L.	acetone, methanol (seeds) <sup>(11)</sup> ; alkaloids, flavonoids, saponins, tannins (fruits) <sup>(6)</sup> ; glycosides, cardiac glycosides (seeds) <sup>(2)</sup>	antibacterial <sup>(11)</sup> ; antimicrobial <sup>(6)(2)</sup> ; antifungal hypoglycaemic, cytotoxic effects, cholesterolemic <sup>(2)</sup>

## Discussion

The knowledge of diversity plants that used in preparing *oke sou* herbal drink is obtained by oral from older indigenous medical practitioner to younger, who is her daughter or her niece. Indigenous medical practitioners has dominant role in keeping the information about composition *oke sou* herbal drink. They have prohibition to bequeath that information to people except to her maternal ancestry. Therefore, not all community at Lako Akeriri village know composition the herbal drink.

*Oke sou* herbal drink believed by community at Lako Akeriri efficacious to maintain the health of women's reproductive function. Based on science investigation *oke sou* herbal drink maintains the reproductive health by keeping the cleanness of organ reproduction<sup>(2)(6)(16)(27)(26)</sup>; reducing bad odor on vagina area and girl's body<sup>(10)(26)(28)</sup>; protecting the organ reproduction from potency of cancer<sup>(8)(29)(26)</sup>; and free radicals<sup>(9)(27)(4)</sup>. Those efficacies are obtained from diversity of plants that used in preparing *oke sou* herbal drink. For example, plants that are useful to keep the cleanness of organ

reproduction (vagina), *Cananga odorata* (Lam.) Hook.f. & Thomson<sup>(26)(28)(13)</sup>, *Curcuma longa* L.<sup>(9)</sup>, *Cymbopogon citratus* (DC.) Stapf.<sup>(12)</sup>, and *Kaempferia galanga* L.<sup>(27)</sup>. Those plants have pharmacological activities such as, anti-microbial, anti-fungal, and antibacterial.

Futhermore, the preparation *oke sou* herbal drink has a boiling stage that aims to extract the phytochemical content in part of plant used<sup>(23)</sup>. The efficacy of *oke sou* herbal drink is better when phytochemicals content in plants used can be completely soluble in water. Meanwhile classification of plant part and crushing separately in order to facilitate the process of squeezing of the juice plant.

## Conclusion

Recorded as many as 66 plant species from 59 genera used for preparation *oke sou* herbal drink with *Cananga odorata* (Lam.) Hook.f. & Thomson being the most frequently mentioned plant by community at Lako Akediri village. The phytochemical content in used plants have various compound, but the pharmacological activities can be summarized in common are antimicrobial, antifungal, aromatherapy, antioxidant, and anti-cancer. The results of this study can be used as new reference for the development medical herbal product based science, especially for maintaining the healthy women's reproduction.

## References

- Amiri, M. S., P. Jabarzadeh, & M. Akhondi. (2012). An ethnobotanical survey of medicinal plants used by indigenous people in Zangelanlo district, Northeast Iran. *Journal of Medicinal Plants Research* 6(5): 749--753
- Ara, N. & M.D.M. Islam. (2009). Phytochemical screening and in vitro antibacterial activity of *Tamarindus indica* seeds ethanolic extract. *Pakista Journal of Pharmacology*. 26(1): pp. 19--23
- Aziz, A., A. Farina, M. Iqbal. (2013). Antioxidant activity and phytochemical composition of *Cynometra cauliflora*. *Journal of Experimental & Integrative Medicine* 3 (4): 337--341
- Batoro, J., D. Setiadi, T. Chikmawati, & Y. Purwanto. (2013). Pengetahuan tentang tumbuhan masyarakat Tengger di Bromo Tengger Semeru Jawa Timur. Jurusan Biologi. FMIPA Universitas Brawijaya: pp. 1--10.
- Chung, J.Y., J.H. Choo, M.H. Lee, J.K. Hwang (2006). Anticariogenic activity of macelignan isolated from *Myristica fragrans* (nutmeg) against *Streptococcus mutans*. *Journal Phyromedicine - Elsevier* 13 (4): 261--266
- Daniyan, S.Y. & H.B. Muhammad. (2008). Evaluation of the microbial activities and phytochemical properties of extracts of *Tamarindus indica* against some diseases causing bacteria. *African Journal of Biotechnology* 7(14): pp. 2451--2453
- Gazzaneo, L.R.S., R.F.P. De Lucena, & U.P. De Albuquerque. (2005). Knowledge and use of medicinal plants by local specialist in an region of Atlantic forest in the state of Pernambuco (Northeastern Brazil). *Journal of Ethnobiology and Ethnomedicine* 10: 1--9
- Gupta, M.P., P.N. Solis, A.I. Calderon, F. Guionneau-Sinclairi, C. Correa, C. Galdames, C. Guerra, A. Espinosa, G.I. Alvenda, G. Robles, & R. Ocampo. 2005. Medical ethnobotany of the Teribes of Bocas del Toro, Panama. *Journal of Ethnopharmacology* 96: 389--401
- Janerio, R.D. (2001). Biological Activities of *Curcuma longa* L. *Mem Inst Oswaldo Cruz* 96(5): 723--728
- Kardinan, A. 2005. *Tanaman penghasil minyak atsiri - komoditas wangi penuh potensi - kiat mengatasi permasalahan praktis*. PT Agromedia Pustaka. Jakarta: vi + pp. 74
- Kothari, V. & S. Seshadri. (2010). In vitro antibacterial activity in seed extracts of *Manilkara zapota*, *Annona squamosa*, & *Tamarindus indica*. *Biol Res* 43: 165--168
- Lee, K.G., T. Shibamoto. (2001). Antioxidant property of aroma extract isolated from clove buds *Syzygium aromaticum* (L.) Merr. Et Perry. *Journal of Food Chemistry - Elsevier* 74(4): 443--448
- Onawunmi, G.O., W.A. Yisak, E.O. Ogunlana. 2002. Antibacterial constituents in the essential oil of *Cymbopogon citratus* (DC.) Stapf. *Journal of Ethnopharmacology* 12(3): 279--2861
- Orabi, K.Y., J. S. Mossa, El-Feraly, & Farouk S. 1991. Isolation and characterization of two antimicrobial agents from mace (*Myristica Fragrans*). *Journal of Natural Products* 54(3): 856--859
- Pinto, E., L.V. Silva, C. Cavaleiro. L. Salgueiro. (2009). Antifungal activity of the clove essential oil from *Syzygium aromaticum* on *Candida*, *Aspergillus*, and dermatophyte species. *Journal of Medical Microbiology* 58(2009): 1454--1462
- Profil desa dan kelurahan. 2011. *Desa Lako Akediri Kec. Sahu Kab. Halmahera Barat*. Direktorat Jenderal Pemberdayaan Masyarakat dan Desa. Kementrian Dalam Negeri: pp. 91
- Rahman M.M., S. S. Lopa, & G. Sadik et al. (2005). Antibacterial and cytotoxic compounds from the bark of *Cananga odorata*. *Fitoterapia*. 76(7-8): pp.758--761
- Riswan, S. & H.S. Roemantyo. (2002). Jamu as Traditional Medicine in Java, Indonesia. *South Pacific Study*. 23 (1) 1-10
- Sacchetti G., S. Maietti, & M. Muzzoli et al. (2005). Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chemistry*. 91(4): pp.621--632
- Sharma R.A, Gescher AJ, Steward WP. (2005) Curcumin: the story so far. *European Journal of Cancer*; 41:1955--68
- Sudibyo, R.S. (2000). The contents of volatile oil isolated from *Kaempferia galanga* rhizomes. Mass spectroscopic approach. *Majalah Farmasi Indonesia*. 11(3): 142--149
- Sujarwo, W., A.P. Keim, V. Savo, P.M. Guarrera, & G. Caneva. (2015). Ethnobotanical study of *loloh*: traditional herbal drinks from Bali (Indonesia). *Journal of Ethnopharmacology* 169: 34--48
- Supardi, S. & M. Notosiswoyo. (2005). Pengobatan Sendiri Sakit Kepala, Demam, Batuk dan Pilek pada Masyarakat Di Desa Ciwalen, Kecamatan Warungkondang, Kabupaten



Cianjur, Jawa Barat. *Majalah Ilmu Kefarmasian*, **2** (3): 134--144

- 24) Suryadarma, I.G.P. (2010). Keanekaragaman tumbuhan bahan kebugaran dalam naskah lontar *rukmini tatwa* masyarakat Bali. *Biota*. **15**(2): 294--305.
- 25) Tajuddin, S. Ahmad, A. Latif, & I.A. Qasmi. (2003). Aphrodisiac activity of 50% ethanolic extracts of *Myristica fragrans* Houtt. (nutmeg) & *Syzygium aromaticum* (L.) Merr. & Perry. (clove) in male mice: a comparative study. *BMC Complementary & Alternative Medicine* **3**(6): 1--5
- 26) Tan, L.T.H., L. H.Lee, W. F.Yin, C.K.Chan., H.A. Kadir, K.G. Chan, & B.H. Goh. (2015). Traditional uses, phytochemistry, and bioactivities of *Cananga odorata* (Ylang-Ylang). *Evidence-Based Complementary and Alternative Medicine*. **2015**: pp 1--30
- 27) Tewtrakul, S., S. Yuenyongsawad, S. Kummee, L. Atsawajaruwan. (2005). Chemical components and biological activities of volatile oil of *Kaempferia galanga* Linn. *Songklanakarin Journal Science and Technology* **27**(2): 503--507
- 28) Zheng G.Q., P.M. Kenney, LKT Lam. (1993) Potential anti-carcinogenic natural products isolated from lemon grass oil and galangal root oil. *Journal of Agricultural and Food Chemistry* **41**:153--6

# Ethnobotanical Study of Oke Sou: Traditional Herbal Drink from Lako Akediri Village in West Halmahera, Indonesia

## ORIGINALITY REPORT

31%

SIMILARITY INDEX

26%

INTERNET SOURCES

12%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

1

[biodiversitas.mipa.uns.ac.id](http://biodiversitas.mipa.uns.ac.id)

Internet Source

18%

2

Wawan Sujarwo, Ary Prihardhyanto Keim, Valentina Savo, Paolo Maria Guarrera, Giulia Caneva. "Ethnobotanical study of Loloh: Traditional herbal drinks from Bali (Indonesia)", *Journal of Ethnopharmacology*, 2015

Publication

1%

3

Teng, Woon-Chien, Ho Kiat, Rossarin Suwanarusk, and Hwee-Ling Koh. "Research on Medicinal Plants for Malaria", *Traditional Herbal Medicines for Modern Times*, 2016.

Publication

1%

4

Wawan Sujarwo, Ary Prihardhyanto Keim. "Ethnobotanical Study of Traditional Building Materials from the Island of Bali, Indonesia", *Economic Botany*, 2017

Publication

1%

5

[medplants.blogspot.com](http://medplants.blogspot.com)

Internet Source

1 %

6

[www.aisf.or.jp](http://www.aisf.or.jp)

Internet Source

1 %

7

S. Augustino. "Medicinal Plants in Urban Districts of Tanzania: Plants, Gender Roles and Sustainable Use", International Forestry Review, 03/2005

Publication

1 %

8

Mary C. Setzer. "The Medicinal Value of Tropical Rainforest Plants from Paluma, North Queensland, Australia", Pharmaceutical Biology (Formerly International Journal of Pharmacognosy), 2/1/2001

Publication

1 %

9

Submitted to Mahidol University

Student Paper

1 %

10

[worldwidescience.org](http://worldwidescience.org)

Internet Source

1 %

11

[www.wrmp.org](http://www.wrmp.org)

Internet Source

1 %

12

[es.scribd.com](http://es.scribd.com)

Internet Source

1 %

13

Submitted to Universitas Terbuka

Student Paper

1 %

14 Eliana Rodrigues. "Plants and animals utilized as medicines in the Jaú National Park (JNP), Brazilian Amazon", *Phytotherapy Research*, 05/2006  
Publication

---

15 Submitted to Unika Soegijapranata  
Student Paper

---

16 Submitted to Georgia Gwinnett College  
Student Paper

---

17 [academic.oup.com](http://academic.oup.com)  
Internet Source

---

18 [www.pharmazeutische-biologie.de](http://www.pharmazeutische-biologie.de)  
Internet Source

---

19 [journalofethnicfoods.biomedcentral.com](http://journalofethnicfoods.biomedcentral.com)  
Internet Source

---

20 Yao Fu, Jin-chao Yang, Anthony B. Cunningham, Alexandra Maria Towns, Yu Zhang, Hua-ying Yang, Jian-wen Li, Xue-fei Yang. "A billion cups: The diversity, traditional uses, safety issues and potential of Chinese herbal teas", *Journal of Ethnopharmacology*, 2018  
Publication

---

21 Submitted to Higher Education Commission Pakistan  
Student Paper

---



---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On