

Contents lists available at ScienceDirect

Journal of Ethnic Foods

journal homepage: <http://journalofethnicfoods.net>

Original Article

Gastronomic ethnobiology of “*terites*”—a traditional Batak Karo medicinal food: A ruminant's stomach content as a human food resourceEndang C. Purba^{a,*}, Marina Silalahi^b, Nisyawati^a^a Universitas Indonesia, Departemen Biologi, Depok, West Java, Indonesia^b Universitas Kristen Indonesia, Fakultas Keguruan dan Ilmu Pendidikan, Departemen Biologi, Jakarta, Indonesia

ARTICLE INFO

Article history:

Received 26 April 2018

Received in revised form

14 June 2018

Accepted 15 June 2018

Available online 21 June 2018

Keywords:

Biocultural heritage

Ethnobotany

Stomach content

Traditional food

Wild edible plants

ABSTRACT

Background: *Terites* is a traditional food of Batak Karo ethnic group, which is cooked with a juice of partly digested food (chyme) of slaughtered cattle. The stomach juice serves as a soup base, cooked together with certain wild and cultivated vegetables, aromatic herbs and possibly also meat. The objectives of this ethnobiological study were to describe *terites* preparation, document plant species used and to discuss possible implications for the human nutrition.

Methods: The data were gathered through individual interviews and group discussions with informants from 6 villages in Karo regency of North Sumatra. The plant specimens were collected in the field and identified taxonomically.

Results: A total of 29 plant species belonging to 17 families were used to prepare *terites*. The main rationale behind consuming this indigenous food was its perceived medicinal value, particularly for the treatment of digestive disorders. Karo people use several lesser-known wild food plants for preparation of this local specialty. To best of our knowledge, consumption of chyme in tropical Asia is so far unique solely to the Batak Karo people. The present ethnographic record of consuming chyme as a medicinal food is also discussed in the context of paleodietary reconstructions.

Conclusion: This extraordinary food heritage of Karo indigenous gastronomy, based on traditional knowledge, indicates rich foodscapes and bio-cultural diversity of the Batak Karo ethnic group.

© 2018 Korea Food Research Institute. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Most of modern medicines were discovered because it was known that certain plants were very effective to treat for particular ailments. Ethnobotanical study can serve as a platform for studying specific relationships between indigenous cultures and plants, thus identifying new chemical compounds used as drugs [1–4]. The traditional food and consumption of healthy and edible plants play an important role in supporting ethnobotany. Unfortunately, many of the traditional foods are approaching extinction even in developing countries such as Indonesia.

Local knowledge in Indonesia is quite extensive because of the considerable diversity of local cultures. Batak Karo is one of the ethnic groups who inhabit in *Taneh Karo* (Karo Land) in North

Sumatra. According to the study by Harsono and Restuati [5], Batak Karo ethnic group has traditional food, *terites*, which is unique and only can be found in this area. The primary meaning of the word *terites* is the liquid of the partially digested grass from the rumen of ruminants. They add some different plants, spices, and meat into the *terites*. Nowadays, the young generations of Batak Karo prefer not to consume this local cuisine.

The Batak Karo people in North Sumatra are endowed with local knowledge about use of plants. They have many culinary specialties and use lot of plants and spices. Furthermore, they believe that all diseases can be cured with healthy food prepared from natural products that have medicinal properties. Unfortunately, most of them have not been documented very well. To initiate for not losing it, the ethnobotanical study was conducted in the Batak Karo community. The aims of this study were to provide information about the process of *terites* preparation, document the plant species used, and discuss possible implications for the human nutrition and further research.

* Corresponding author.

E-mail address: endang.christine@yahoo.com (E.C. Purba).

2. Materials and methods

2.1. Study area

The study was conducted in Semangat Gunung, Jaranguda, Merdeka, Cinta Rayat, Doulu and Lingga villages of Karo Regency in North Sumatra, Indonesia (Fig. 1). Batak Karo people who live in Semangat Gunung, Jaranguda, Merdeka, Cinta Rayat, and Doulu generally maintain a traditional medicine and have access to forests areas near to Mt. Sibayak. Meanwhile, Lingga is a cultural village that preserves some local knowledge of Batak Karo, e.g., food, medicinal plants, and traditional buildings. Geographically, these villages are situated about 1,162–1,453 m above sea level.

The Karo is one of the six Batak tribes with traditional homelands in the periequatorial highlands of North Sumatra, Indonesia [6] (Fig. 2). The majority of Batak Karo people are farmers, primarily cash crop and subsistence agriculturalists. They use traditional farming implements and water buffalo or cow for traction.

Batak Karo ethnic group is known as the group of people who have local knowledge about the use of plants. They believe that food and medicinal plants are of equal importance to prevent and treat disease. They use plants as pharmaceuticals and food and during ceremonials [6,7]. Various studies have been conducted to

assess local knowledge about the use of plants by those ethnic groups such as the use of medicinal plants in the village Tongkoh, Karo [6] and in Kaban Tua, Karo [8].

2.2. Interviews and data collection

Information for this study was collected during April–June 2014 and August 2016 through questionnaires and semistructured and general conversations from the herbalists, village elders, and farmers. The interview was carried out with the help of 29 informants; Semangat Gunung (7), Jaranguda (3), Merdeka (2), Cinta Rayat (4), Doulu (4), and Lingga (9). The head of the village was used as a key informant who could provide information on how many people know about *terites*. The informants were chosen based on traditional ecological knowledge, professional activity, and age. Each informant was interviewed individually in Batak Karo language.

Informants were asked to provide a list of ingredients used to prepare *terites*. The local names of all the plants used, use of plants, plant parts used, and mode of preparation of *terites* were then collected under the guidance of whoever gave the information about the preparation of *terites*. The plants used included not only wild species but also cultivated plants that had been taken from the




Fig. 1. The location of Karo Regency in North Sumatra, Indonesia. Karo's geography with mountainous area provides cool weather.



Fig. 2. Batak Karo people. Traditional food is an important part of Batak Karo culture. They usually treat others with meals to make new friends and enhance relationships.

forest and planted in gardens or agricultural fields. The specimens were identified and deposited at the herbarium of the Universitas Indonesia, Depok, Indonesia. Scientific names of the plants species were verified using The Plant List online source.

3. Results

3.1. Plant species used in terites soup

A total of 29 plant species were used to prepare *terites* which belong to 17 families and 26 genera (Table 1). Among them, 18 (62 %) were herbs, six (21 %) were shrubs, and five (17 %) were trees. The family Zingiberaceae had the highest proportion of herbs represented by five species, followed by Liliaceae with four, and Euphorbiaceae and Solanaceae, each with 3 (Fig. 3). Leaves are the most frequently used to prepare *terites* (Fig. 4). This adds up to about 34% of wild edible plant species. *Bischofia javanica* Blume was cited by 29 informants as an ingredient of *terites*, *Aleurites moluccanus* by 27 informants, and *Begonia laruei* by 22 informants in the same way. These and other most widely used wild edible plants are listed and described in Table 2.

3.2. Preparations and prescription

Terites actually refers to the juice of partially digested grass from the rumen of ruminants; the juice is then filtered using a clean cloth or sieve (Fig. 5). Batak Karo people use the filtrate as a stock to make traditional soup that is cooked with spices, herbs, and meat and served during cultural occasions. *Pagit-pagit* is another name for *terites*. “Pagit” means “bitter,” which refers to the taste of *terites* soup.

Cooking *terites* involves pounding and grinding ingredients. Briefly, *terites* soup was prepared by grinding and filtering *Bischofia javanica* (bark); pounding *Etlingera elatior* (fruits), *Alpinia galanga* (rhizome), and *Cymbopogon citratus* (whole plant); chopping *Solanum lycopersicum* (fruits), *Begonia laruei* (leaves and stems), and *Medinilla speciosa* (leaves and stems); and grinding spices such as *Allium fistulosum* (leaves), *A. cepa* (bulbs), *A. sativum*

(bulbs), *Capsicum annuum* (fruits), *Aleurites moluccanus* (fruit), *Curcuma domestica* (rhizome), *Zingiber* sp. (rhizome), and *Zingiber officinale* (rhizome). The stock that contains the filtrate of the rumen and water was cooked with beef or pork. The stock was cooked until the meat becomes tender. The juice of *Bischofia javanica* and pounded *Alpinia galanga* and *Cymbopogon citratus* were added to the stock until its color changed to light green. Then, *Citrus hystrix*, the fruit of *Etlingera elatior*, and whole spices are added and cooked until the stock gets spicy and piquant flavor. Furthermore, ground *Aleurites moluccanus*, *Begonia laruei*, flowers of *Etlingera elatior*, and *Manihot esculenta* were added and cooked until the leaves of *Manihot esculenta* were cooked well. At the end, *Cocos nucifera*, *Solanum lycopersicum*, *Allium fistulosum*, and salt were added and cooked well (Fig. 6). The fruit of *Cocos nucifera* provides coconut milk that is commonly called as *santan* in Indonesian language. It is generally used to get a savory taste in Minang cuisines such as *ketupat bareh* and *katan kapau* [9]. *Cocos nucifera* has a high cultural importance and high frequency of use and is irreplaceable with other species among the Sasak tribe, Lombok Island [10].

In some cases, the Batak Karo people add *Artabotrys suaveolens*, *Apium graveolens*, *Enhydra fluctuans*, *Allium tuberosum*, *Medinilla speciosa*, *Polygonum chinense*, *Zanthoxylum acanthopodium*, *Solanum torvum*, *Debregeasia longifolia*, and *Musa acuminata* to improve the acceptability of *terites* soup.

4. Discussions

Terites is a unique cuisine and belongs only to the Batak Karo culture (Fig. 7). *Terites* is served during cultural occasions such as weddings and harvest festivals. In special occasions, Batak Karo people have a cultural ceremony of slaughtering cows. It determines how big the occasions is; the more the cows, the bigger the occasion. Now, many Karo restaurants in Medan, Berastagi, and Kabanjahe serve *terites* soup in their menu. Batak Karo people believe that *terites* soup is good for humans, especially for elders, because it can treat gastritis and cold, act as an appetite enhancer, and lower cholesterol levels. According to a study carried out by Harsono and Restuati [5], *terites* was commonly used to treat gastritis, to support the availability of energy reserves in the human body, and to keep the body warm in low temperatures. Besides the Batak Karo people, in high latitude environments such as Arctic, where there is little edible vegetation for humans, Inuit people consume the chyme from seal, narwhal, walrus, and reindeer [11]. This potentially explains the reason why Batak Karo people, living in an area mainly consisting of mountains (Sinabung and Sibayak) and highlands, are familiar with chyme consumption [6,12].

The chyme consumption for humans has played a particular role in the diet of the young and the old [11]. The chyme from an herbivore with a specialized digestion system and microbiota capable of digesting chitin and cellulose, which humans cannot, could be the potential energy source because this provides additional benefits in enabling access to otherwise inedible plant species [11]. Ruminants evolved to digest grass; hence, the four stomachs were dedicated to the digestion of cellulose. Bacterial fermentation creates a high level of conjugated linoleic acid (fatty acid) [13–15].

Rumen fluid is an organic material that has been digested by rumen microbes; the cattle rumen fluid contains protein (10.56%), vitamin, ash (18.38%), Ca (1.34%), and P (0.32%), and the buffalo rumen fluid contains protein (7.97%), vitamin, ash (24.40%), Ca (0.67%), and P (0.63%) [16–18]. *Terites* soup contains protein (19.9%), fat (15.4%), and crude fiber (1.68%) [19]. The fat was considered to have come from the meat that was added to the soup. The *terites* stock that is obtained from the rumen fluid contains high

Table 1
Plant species used in the preparation of *terites*.

Scientific name	Plant life form	Habitat	Local name	Parts used	Number of informants	Villages
Annonaceae						
<i>Artabotrys suaveolens</i> (Blume) Blume	Shrub	W	Bidung-bidung	Leaves	4	Sg
Apiaceae						
<i>Apium graveolens</i> L.	Herb	C	Daun sop	Leaves	2	Do
Areaceae						
<i>Cocos nucifera</i> L.	Tree	C	Tualah	Fruit	26	Li, Me, Sg, Ja, Cr, and Do
Asteraceae						
<i>Enhydra fluctuans</i> Lour	Herb	W	Kurmak ndekah	Leaves	5	Cr and Do
Begoniaceae						
<i>Begonia laruei</i> M. Hughes	Herb	W	Riang-riang	Leaves	22	Li, Me, Sg, Ja, and Do
Euphorbiaceae						
<i>Manihot esculenta</i> Crantz	Shrub	C	Gadung	Leaves	29	Li, Me, Sg, Ja, Cr, and Do
<i>Aleurites moluccanus</i> (L.) Willd.	Tree	W	Kembiri	Fruits	27	Li, Sg, Ja, Cr, and Do
<i>Bischofia javanica</i> Blume.	Tree	W	Cingkam	Barks	29	Li, Me, Sg, Ja, Cr, and Do
Liliaceae						
<i>Allium fistulosum</i> L.	Herb	C	Daun pere	Leaves	17	Li, Cr, and Do
<i>Allium tuberosum</i> Rottler ex Spreng	Herb	C	Gundera mbelang	Leaves	1	Do
<i>Allium cepa</i> L.	Herb	C	Pia	Bulbs	29	Li, Me, Sg, Ja, Cr, and Do
<i>Allium sativum</i> L.	Herb	C	Lasuna	Bulbs	29	Li, Me, Sg, Ja, Cr, and Do
Melastomataceae						
<i>Medinilla speciosa</i> Blume	Herb	W	Kambing-kambing	Leaves	11	Me, Sg, and Ja
Moraceae						
<i>Ficus fistulosa</i> Reinw	Tree	W	Ober	Leaves	1	Do
Musaceae						
<i>Musa acuminata</i> Colla	Herb	C	Galuh si uncim	Flowers	4	Sg
Myrtaceae						
<i>Eugenia polyantha</i> Barb. Rord.	Tree	W	Salam	Leaves	13	Li and Cr
Poaceae						
<i>Cymbopogon citratus</i> (DC.) Stapf	Herb	C	Sere	Leave	29	Li, Me, Sg, Ja, Cr, and Do
Polygonaceae						
<i>Polygonum chinense</i> Linn	Herb	W	Siang-siang	Leaves	10	Sg and Ja
Rutaceae						
<i>Citrus hystrix</i> DC.	Shrub	C	Rimo mungkur (purut)	Leaves	13	Li and Cr
<i>Zanthoxylum acanthopodium</i> DC	Shrub	C	Tuba	Fruits	1	Do
Solanaceae						
<i>Solanum lycopersicum</i> L.	Herb	C	Tomat	Fruits	29	Li, Me, Sg, Ja, Cr, and Do
<i>Capsicum annuum</i> L.	Herb	C	Cina	Fruits	29	Li, Me, Sg, Ja, Cr, and Do
<i>Solanum torvum</i> Sw	Shrub	C	Rimbang	Fruits	17	Me, Sg, Ja, Cr, and Do
Urticaceae						
<i>Debregeasia longifolia</i> (Burm.f.) Wedd	Shrub	W	Cepira	Leaves	5	Cr and Do
Zingiberaceae						
<i>Etlingera elatior</i> (Jack.) R.M.Sm.	Herb	C	Asam patikala/ Asam cekala/Kincung	Fruits and flowers	29	Li, Me, Sg, Ja, Cr, and Do
<i>Alpinia galanga</i> L. (Willd.)	Herb	C	Kelawas	Rhizomes	27	Li, Sg, Ja, Cr, and Do
<i>Curcuma domestica</i> Valetton	Herb	C	Kuning gersing	Rhizomes	23	Li, Sg, Ja, and Do
<i>Kaempferia galanga</i> L.	Herb	C	Kaciwer	Rhizomes	14	Sg, Ja, and Do
<i>Zingiber</i> sp.	Herb	C	Alia	Rhizomes	14	Sg, Ja, and Do

C, cultivated; Cr, Cinta Rayat; Do, Doulu; Ja, Jaranguda; Li, Lingga; Sg, Semangat Gunung; W, wild.




Fig. 3. Number of species and families used for preparing *terites*. Zingiberaceae contributed the highest number of species to the cuisine. It is consisted of five species.




Fig. 4. Number of species and plant parts used for preparing *terites* soup. Leaves are more widely used than barks. The bark of *Bischofia javanica* is used not only as an ingredient of *terites* soup but also as a medicinal plant.

Table 2

Informant consensus on most commonly used wild edible plants.

Scientific name of wild edible plants	No. of informants	Percentage
<i>Bischofia javanica</i> Blume.	29	22.8
<i>Aleurites moluccanus</i> (L.) Willd.	27	21.3
<i>Begonia laruei</i> M. Hughes	22	17.3
<i>Eugenia polyantha</i> Barb. Rord.	13	10.2
<i>Medinilla speciosa</i> Blume	11	8.7
<i>Polygonum Chinense</i> Linn	10	7.9
<i>Enhydra fluctuans</i> Lour	5	3.9
<i>Debregeasia longifolia</i> (Burm.f.) Wedd.	5	3.9
<i>Artabotrys suaveolens</i> (Blume) Blume	4	3.1
<i>Ficus fistulosa</i> Reinw	1	0.8
	100	



Fig. 7. The *terites* soup. *Terites* is a Batak Karo soup. It is cooked using 29 herbs and meat. This soup is usually served with rice during a wedding or harvest festival. It is the favorite among the Batak Karo elders.

amounts of crude fiber. It is mixed with spices and vegetables which is good for the digestive system of the human body.

Traditions related to gathering and cooking food plants are very popular in the Karo area. Ten wild and 19 cultivated plants species were identified which Karo uses as ingredients of *terites* soup. The Karonese used to collect common food plants for cooking before returning home after spending the entire day working in the fields. Most of them are vegetables and spices such as *Apium graveolens*, *Manihot esculenta*, *Allium fistulosum*, *Cymbopogon citratus*, *Zanthoxylum acanthopodium*, *Solanum lycopersicum*, *Capsicum annuum*, *Etlingera elatior*, and *Curcuma domestica*. Generally, men collect a few wild plants from the woods located further from the village surrounding fields (*kerangen*). Today, some of the wild plants that are used as medicinal or food plants can be found in traditional markets in Karo Regency: these include *Begonia laruei*, *Aleurites moluccanus*, *Bischofia javanica*, and *Eugenia polyantha*.

The Zingiberaceae is the most represented family in the ingredients of *terites*. It is well known as the most important source of medicinal plants in Karo [20] and Indonesia [8,21–23]. A large number of informants cited *Etlingera elatior*. It is native to Sumatra, Indonesia, and has been found in many places throughout South-east Asia [24,25]. In Karo, *E. elatior* is locally known as *asam cekala* or *asam patikala* (fruit) and *kincung* (flower). Both are important spices in Karonese dishes including *arsik*. In Bali, the flower buds of the plant are used for chili sauce such as *sambal bongkot* and *sambal kecicang*. In Malaysia, they called it as *bunga kantan*, which is used as a spice in Malaysian dishes such as *laksa* and *nasi ulam* [26]. In Thailand, they eat the young shoots and flower buds as a vegetable [27]. *E. elatior* is also known to have medicinal uses in treating diseases. In Malaysia, fruits of *E. elatior* are used to treat earache; the leaves are used for cleaning wounds and during bathing to removing body odor [28,29]. Batak Simalungun people used the leaves and stems to cure cough and ulcer and the decoction to maintain health [7]. Batak Karo people used stems and leaves to prepare a decoction *tawar* that is used to cure weakness and have a steam bath by postpartum women [20,30]. Some results showed that flowers of *E. elatior* had high values of antimicrobial, antibacterial, antifungal, antioxidant, antibacterial [29], and anticancer [31] properties. Leaves have the highest level of antioxidants. It has great potential to be developed into natural preservatives that are applicable to be used in the food, and most are represented as nutraceuticals [29].



Fig. 5. Filtering the juice of partially digested grass from the rumen of ruminants. The filtered *terites* is used as a stock to prepare the *terites* soup. Water is added to the stock before cooking it.



Fig. 6. Cooking the *terites* soup. Meat, spices, and herbs are put into a pot to which coconut milk is added, and the whole thing is simmered until done.

The Indonesian Food and Drug Administration has recommended in its traditional health maintenance program the following nine species of medicinal plants: (1) *Piper retrofractum*, (2) *Curcuma xanthorrhiza*, (3) *Curcuma domestica*, (4) *Guazuma ulmifolia*, (5) *Andrographis paniculata*, (6) *Zingiber officinale*, (7) *Morinda citrifolia*, (8) *Eugenia polyantha*, and (9) *Psidium guajava* [32]. Among the nine species, two species are used in the preparation of *terites*—*Curcuma domestica* and *Eugenia polyantha*. Indonesians use these species for various purposes, from cooking to medicine. *Curcuma domestica* is known to have medicinal uses in treating abdominal pain [33]. This species has been extensively studied and has demonstrated several pharmacological activities e.g., antibacterial, antifungal [34], and antioxidant [35] properties. Meanwhile, *Eugenia polyantha* has various ethnobotanical uses (in treating hypertension, gastritis, diarrhea, diabetes mellitus, and high cholesterol) [8].

A large number of informants cited *Bischofia javanica* (Table 2). Batak Karo people use this species to minimize the strong aroma of the rumen. Some studies have been conducted to evaluate its antiulcer, anthelmintic, antidiarrheal [36], antiparasitic [37], antiinflammatory, antinociceptive [38], antileukemic [39], and antimicrobial [40] activities. These activities do not explain all the current uses for *terites*, so probably this species could have other pharmacological activities that are still unknown. The leaves and bark of *Bischofia javanica* are known to be rich in tannin, so it can be used as a food-coloring agent [41]. Fatty acid profile of *Bischofia javanica* seed oil indicates that it is an important source of omega-3 fatty acids and can be used as food supplement and for medicinal purposes in the future [30]. Some local people in Indonesia use *Bischofia javanica* to treat some diseases; for example, people in Lahat Regency, South Sumatra, used the leaves to treat diarrhea [42], and Batak Simalungun people of North Sumatra used the roots and barks to treat diarrhea, diabetes mellitus, gastritis, stomach ache, and injuries [7]. In India, local people use the stem bark of *Bischofia javanica* to cure nervous disorders, to stimulate hair growth [43], and to treat skin diseases [44].

Aleurites moluccanus (Euphorbiaceae) is also cited by a large number of informants. It is native to the Moluccas and widely found in Southeast Asia. It is commonly used in Indonesian dishes as curry pastes that are eaten with vegetables and rice and as a meat tenderizer. In Java, the bark is used to treat diarrhea and dysentery; meanwhile in Malaysia, the leaves are used to treat headaches, fevers, ulcers, gonorrhea, and joint inflammation [45]. The nuts of *Aleurites moluccanus* inhibit the growth of bacterial species associated with the onset of rheumatoid arthritis, ankylosing spondylitis, and rheumatic heart disease [46].

Both medicinal plants and foods are considered to have healthful properties, preventing disease by strengthening the body's defense system [47]. Several plants cited in this study are poorly investigated for their pharmacological activity or chemical profiles. Considering that many other plant species instead have already been researched to cure many diseases and health problems, we believe that these understudied plant species also could have interesting applications in pharmacology and as new food plants.

This study indicated that Batak Karo people in Karo Regency of North Sumatra have made efficient use of cow's stomach content within the indigenous food system. The local people perceive *terites*, an indigenous food, to be of high medicinal value, particularly because of the bitter flavor of the chyme and also because of the aromatic plants added to compensate the bitterness and unpleasant scent of the stomach fluid. In conjunction with the consumption of *terites*, the Batak Karo people still preserve traditional knowledge related to remarkable wild ingredients and processing of the chyme. The present study revealed a complex relationship

between a human food culture and biodiversity of the surrounding environment. A detailed ethnobiological view on one complex cultural food showed importance of this approach by discovering several uncommon food plant species. Some of the species would be interesting to study for their nutritional content and pharmacological properties, e.g., *Bischofia javanica*. In (sub-)tropical Asia, the consumption of animal stomach content seems to be unique to Batak Karo people in Sumatra, which is particularly interesting, considering the evergreen tropical environment with abundance of natural resources. Noteworthy is also the fact that this practice has been associated with hunter-gatherers and pastoralists, whereas Karo people are farmers combining both crop production and animal husbandry. Therefore, this is the first record of consuming the stomach content of cows. The reason behind the consumption of chyme among Karo lies rather on the medicinal perception of the chyme. Considering the scientific discussion whether the early hominins were using bitter-tasting plants for self-medication or as a food resource (condiments, or as part of herbivore's stomach content), the contemporary use of chyme among Batak Karo people may indicate that stomach content could have been consumed by our ancestors as food that had medicinal properties as well.

Conflicts of interest

The authors have no conflicts of interest to disclose.

Acknowledgments

The authors are grateful for the kindness and cooperation of people in Semangat Gunung, Jaranguda, Merdeka, Cinta Rayat, Doulu, and Lingga villages, Karo Regency, North Sumatra, Indonesia.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.jef.2018.06.002>.

References

- [1] Balick MJ and Cox PA. Ethnobotanical research and traditional health care in developing countries. In: Bodeker G, Bhat KKS, Burley J and Vantomme P, editors. Medicinal plants for forest conservation and health care, vol. 92. Rome: Food and Agriculture Organization of the United Nations; 1997. p. 12–23.
- [2] Fabricant DS and Farnsworth NR. The value of plants used in traditional medicine for drug discovery. Environ Health Perspect 2001;109(1):69–75.
- [3] Gu R, Wang Y, Long B, Kennely E, Wu S, Liu B, Li P and Long C. Prospecting for bioactive constituents from traditional medicinal plants through ethnobotanical approaches. Biol Pharm Bull 2014;37(6):903–15.
- [4] Pieroni A. Medicinal plants and food medicines in the folk traditions of the upper Lucca Province, Italy. J Ethnopharmacol 2000;70(3):235–73.
- [5] Harsono T and Restuati M. Terites sebagai makanan budaya Suku Batak Karo: suatu tinjauan etnobotani. In: Prosiding seminar nasional etnobotani III. Bogor: LIPI & Kehati Indonesia; 1998. p. 57–61.
- [6] Singarimbun M. Kutagambar: a village of the Karo. In: Koentjaraningrat, editor. Villages in Indonesia. Singapore: Equinox Publishing; 2007. p. 115–28.
- [7] Silalahi M, Supriatna J, Walujo EB and Nisyawati. Local knowledge of medicinal plants in sub-ethnic Batak Simalungun of North Sumatra, Indonesia. Biodiversitas 2015;16(1):44–54.
- [8] Silalahi M. The ethnobotany of the medicinal plants in sub-ethnic Batak, North Sumatra and the conservation perspective, dissertation. Indonesia: Universitas Indonesia; 2014. p. 140.
- [9] Rianti A, Novenia AE, Christopher A, Lestari D and Parassih EK. Ketupat as traditional food of Indonesian culture. J Ethnic Foods 2018;5:4–9.
- [10] Sukenti S, Hakim I, Indriyani S, Purwanto Y and Matthews PJ. Ethnobotanical study on local cuisine of the Sasak tribe in Lombok Island, Indonesia. J Ethnic Foods 2016;3:189–200.
- [11] Buck LT, Berbesque JC, Wood BM and Stringer CB. Tropical forager gastrophagy and implications for extinct hominin diets [Internet]. 2015. <https://doi.org/10.1016/j.jasrep.2015.09.025> [cited 2017 Sept 14]. Available from: .

- [12] Anderson J. Mission to the East Coast of Sumatra in 1823. Kuala Lumpur: Oxford University Press; 1971. p. 424.
- [13] Elmore JS and Mottram DS. The role of lipids in the flavor of cooked beef. *Dev Food Sci* 2006;43:375–8.
- [14] Alfaia CPM, Alves SP, Martins SIV, Costa ASH, Fontes CMGA, Lemos JPC, Bessa RJB and Prates JAM. Effect of the feeding system on intramuscular fatty acids and conjugated linoleic acid isomers of beef cattle, with emphasis on their nutritional value and discriminatory ability. *Food Chem* 2009;114: 939–46.
- [15] Daley CA, Abbot A, Doyle PS and Larson S. A review of fatty acid profiles and antioxidant content in grass-fed and grain-fed beef. *Nutr J* 2010;9(10): 1–12.
- [16] Krause DO, Denman SE, Mackie RI, Morrison M, Rae AL, Attwood GT and McSweeney CS. Opportunities to improve fiber degradation in the rumen: microbiology, ecology, and genomics. *FEMS Microbiol Rev* 2003;27: 663–93.
- [17] Choudhury PK, Salem AZM, Jena R, Kumar S, Singh R and Puniya AK. Rumen microbiology: an overview. In: Puniya AK, Singh R and Kamra DN, editors. *Rumen microbiology: from evolution to revolution*. India: Springer; 2015. p. 3–16.
- [18] Tjakradidjaja AS, Suryahadi, Bakrie B and Permana Z. Nutrient potential of biomineral supplement for dairy calf produced from rumen fluid-byproduct of slaughtered house. In: Nurdin E, Rusfidra, Zain M, Aritonang SN and Rusmana, editors. *Proceeding international seminar of dairy cattle 2009 "improving productivity of dairy cattle and dairy product using natural sources"*. Padang, Indonesia: Universitas Andalas; 2009. p. 65–9.
- [19] Suparthana IP, Perangin-angin RS and Agung IGN. Food safety and nutrients of Terites, a case study in Kabanjahe, the city of Karo Region, Sumatera Utara. *Media Ilmiah Teknologi Pangan* 2015;2(2):163–9.
- [20] Purba EC, Nisyawati and Silalahi M. The ethnomedicine of the Batak Karo people of Merdeka sub-district, North Sumatra, Indonesia. *Int J Biol Res* 2016;4(2):181–9.
- [21] Siagian MH and Sunaryo. Pemanfaatan suku Zingiberaceae sebagai obat tradisional oleh masyarakat Lembak Delapan, Bengkulu. In: Staf peneliti pusat penelitian biologi, editors. *Indeks beranotasi keanekaragaman hayati*. Bogor, Indonesia: LIPI; 1996. p. 237–46.
- [22] Kuntorini EM. Botani ekonomi suku Zingiberaceae sebagai obat tradisional oleh masyarakat di Kotamadya Banjarbaru. *Bioscientiae* 2005;2(1):25–36.
- [23] Susiarti S, Purwanto Y and Walujo EB. Medicinal plant diversity in The Tesso Nilo National Park, Riau, Sumatra, Indonesia. *Reinwardtia* 2008;12(5): 383–90.
- [24] Jaafar FM, Osman CP, Ismail NH and Awang K. Analysis of essential oils of leaves, stems, flowers and rhizomes of *Etlingera elatior* (Jack) R.M. Smith. *Malays J Anal Sci* 2007;11(1):269–73.
- [25] Juhari NH, Lasekan O, Kharidah M and Karim SAB. Optimization of hot-air drying conditions on the physicochemical characteristics of torch ginger (*Etlingera elatior*). *J Food Agric Environ* 2012;10(2):64–72.
- [26] Ng CY and Karim SAB. Historical and contemporary perspectives of the Nyonya food culture. *J Ethnic Foods* 2016;3:93–106.
- [27] Ghasemzadeh A, Jaafar HZE and Ashkani S. Secondary metabolites constituents and antioxidant, anticancer and antibacterial activities of *Etlingera elatior* (Jack) R.M.Sm grown in different locations of Malaysia. *BMC Compl Alternative Med* 2015;15:335–44.
- [28] Ibrahim H and Setyowati FM. *Etlingera*. In: de Guzman CC and Siemonsma JS, editors. *Plant resources of South-east Asia*. Wageningen: Pudoc; 1999. p. 123–6.
- [29] Chan E, Lim Y and Omar M. Antioxidant and antibacterial activity of leaves of *Etlingera* species (Zingiberaceae) in Peninsular Malaysia. *Food Chem* 2007;104(4):1586–93.
- [30] Indra R, Bachheti RK and Archana J. Chemical composition, mineral and nutritional value of wild *Bischofia javanica* seed. *Int Food Res J* 2013;20(4): 1747–51.
- [31] Ghasemzadeh A, Jaafar HZ, Rahmat A and Ashkani S. Secondary metabolites constituents and antioxidant, anticancer and antibacterial activities of *Etlingera elatior* (Jack) R.M.Sm grown in different locations of Malaysia. *BMC Compl Alternative Med* 2015;15(335):1–10.
- [32] Dewoto HR. Pengembangan obat tradisional Indonesia menjadi fitofarmaka. *Majalah Kedokt Indones* 2007;57(7):205–11.
- [33] Siriruga P. The Zingiberaceae: species diversity and their uses [Internet]. 1999 [cited 2017 June 17]. Available from: <https://www.scribd.com/document/190466707/Thai-Zingiberaceae-Species-Diversity-Andn-Their-Uses>.
- [34] Banerjeea and Nigam SS. Antimicrobial efficacy of the essential oil of *Curcuma longa*. *Indian J Med Res* 1978;68:864–6.
- [35] Phan TT, See P, Lee ST and Chan SY. Protective effects of cur-cumin against oxidative damage on skin cell in vitro: its implication for wound healing. *J Trauma* 2001;51:927–31.
- [36] Gupta DR, Dhiman RP, Naithani S and Ahmed B. Chemical investigation of *Bischofia javanica* Blume. *Pharmazie* 1988;43:222–3.
- [37] Alen Y, Nakajima S, Nitoda T, Baba N, Kanzaki H and Kawazu K. Antinematodal activity of some tropical rainforest plants against the pinewood nematode *Bursaphelenchus xylophilus*. *Zeitschrift Fur Naturforschung C* 2000;55(3–4): 295–9.
- [38] Sutharson L, Nath LK, Kar PK, Shila EB and Rajan JV. Anti-inflammatory and antinociceptive activities of methanolic extract of leaves of *Bischofia javanica* Blume on experimental animals. *Asian J Chem* 2007;7:5150–6.
- [39] Sutharson L, Roy S, Joseph RV and Nath LK. Antileukemic activity of the leaf extract of *Bischofia javanica* Blume on human leukemic cell lines. *Indian J Pharmacol* 2011;43(2):143–9.
- [40] Khan MR, Kihara M and Omoloso AD. Anti-microbial activity of *Bidens pilosa*, *Bischofia javanica*, *Elmerillia papuana* and *Sigesbekia orientalis*. *Fitoterapia* 2001;72:662–5.
- [41] Tanaka T, Nonaka GI, Nishioka I, Kouno I and Ho FC. Bischofinanin, a dimeric dehydroellagitannin from *Bischofia javanica*. *Phytochemistry* 1995;38(2): 509–13.
- [42] Harmida S and Yuni VF. Studi etnofitomedika di Desa Lawang Agung Kecamatan Mulak Ulu Kabupaten Lahat Sumatra Selatan. *J Penelitian Sains* 2011;14(1):1–5.
- [43] Ignacimuthu S, Ayyanar M and Sankarasivaraman K. Ethnobotanical investigations among tribes in Madurai District of Tamil Nadu (India). *J Ethnobiol Ethnomed* 2006;2(25):1–7.
- [44] Prasad PRC, Reddy CS, Raza SH and Dutt CBS. Folklore medicinal plants of North Andaman Islands, India. *Fitoterapia* 2008;79:458–64.
- [45] Krisnawati H, Kallio M and Kanninen M. *Aleurites moluccana* (L.) Willd. Ecology, silviculture and productivity. Bogor: Center for International Forestry Research; 2011. p. 11.
- [46] Mpala LN, Chikowe GR and Cock IE. *Aleurites moluccanus* (L.) Willd. Extracts inhibit the growth of bacterial triggers of selected autoimmune inflammatory diseases. *PharmacognosyCommun* 2017;7(2):83–90.
- [47] Etkin N and Ross P. Pharmacological implications of "wild" plants in the Hausa diet. In: Etkin NL, editor. *Eating on the Wild Side: the pharmacologic, ecologic, and social implications of using noncultigens*. Tucson: University of Arizona Press; 1994. p. 85–101.